

FINAL

BIOLOGICAL SURVEY REPORT

FOR

CONSTRUCTION, OPERATION, AND MAINTENANCE
OF VEHICLE FENCE AND RELATED
TACTICAL INFRASTRUCTURE
TUCSON SECTOR, ARIZONA

(b) (7)(E) STATION



U.S. DEPARTMENT OF HOMELAND SECURITY
U.S. CUSTOMS AND BORDER PROTECTION
U.S. BORDER PATROL TUCSON SECTOR, ARIZONA

Prepared by



SEPTEMBER 2008

Warning! This document, along with any attachments, contains NON-PUBLIC INFORMATION exempt from release to the public by federal law. It may contain confidential, legally privileged, proprietary or deliberative process inter-agency/intra-agency material. You are hereby notified that any dissemination, copying, or further distribution of this information to unauthorized individuals (including unauthorized members of the President-elect Transition Team) is strictly prohibited. Unauthorized disclosure or release of this information may result in loss of access to information, and civil and/or criminal fines and penalties.

BW FOIA CBP 002493

ABBREVIATIONS AND ACRONYMS

AZDA	Arizona Department of Agriculture
AZGFD	Arizona Game and Fish Department
BLM	Bureau of Land Management
BMP	Best Management Practices
BSR	Biological Survey Report
CBP	U.S. Customs and Border Protection
CFR	Code of Federal Regulations
cm	centimeter(s)
CWA	Clean Water Act of 1977
°F	Degrees Fahrenheit
e ² M	engineering-environmental Management, Inc.
ESP	Environmental Stewardship Plan
FE	Federally Endangered
GAP	Gap Analysis Program
GIS	Geographic Information System
GPS	Global Positioning System
HDMS	Heritage Data Management System
HS	Highly Safeguarded
m	meter(s)
m ²	square meters
MBTA	Migratory Bird Treaty Act of 1918, as amended
MJD	Multi-Jurisdictional Dataset
mph	Miles per hour
NVCS	National Vegetation Classification System
OHM	Ordinary high water mark
ROE	Right-of-Entry
(b) (7)(E)	
USACE	U.S. Army Corps of Engineers
USBP	U.S. Border Patrol
U.S.C	United States Code
USFWS	U.S. Fish and Wildlife Service
USGS	U.S. Geological Survey
UTM	Universal Transverse Mercator
WRCC	Western Regional Climate Center

FINAL BIOLOGICAL SURVEY REPORT
FOR
CONSTRUCTION, OPERATION, AND MAINTENANCE
OF VEHICLE FENCE AND RELATED TACTICAL INFRASTRUCTURE
TUCSON SECTOR, ARIZONA
(b) (7)(E) TATION

TABLE OF CONTENTS

ABBREVIATIONS AND ACRONYMS	INSIDE FRONT COVER
1. INTRODUCTION	1-1
2. PROJECT DESCRIPTION	2-1
2.1 SURVEY METHODS	2-2
2.2 ARIZONA GAME AND FISH DEPARTMENT; ARIZONA NATURAL HERITAGE PROGRAM, HERITAGE DATA MANAGEMENT SYSTEM.....	2-4
3. ENVIRONMENTAL SETTING	3-1
4. BIOLOGICAL RESOURCES.....	4-1
4.1 VEGETATION CLASSIFICATION	4-1
4.1.1 Madrean Pinyon-Juniper Woodland Ecological System (CES305.797)	4-4
4.1.2 North American Warm Desert Riparian Woodland and Shrubland Ecological System (CES302.753)	4-4
4.1.3 Apacherian-Chihuahuan Mesquite Upland Scrub Ecological System (CES302.733).....	4-9
4.1.4 Chihuahuan Mixed Desert and Thorn Scrub (CES302.734).....	4-11
4.1.5 Apacherian-Chihuahuan Mesquite Upland Scrub Ecological System (CES302.733).....	4-15
4.1.6 North American Warm Desert Riparian Mesquite Bosque Ecological System (CES302.752)	4-19
4.1.7 North American Warm Desert Wash Ecological System (CES302.755)	4-23
4.1.8 Apacherian-Chihuahuan Semi-Desert Grassland and Steppe Ecological System (CES302.735)	4-26
4.1.9 North American Warm Desert Cienega Ecological System (CES302.747)	4-27
4.1.10 North American Warm Desert Playa Ecological System (CES302.751)	4-28
4.1.11 Other Nonnative Herbaceous Vegetation Alliances and Associations	4-30
4.2 PLANT SPECIES IDENTIFIED.....	4-30
4.3 SURVEY CORRIDOR CHARACTERISTICS AND DESCRIPTION OF HABITAT QUALITY	4-36
4.4 WETLANDS AND WATERS OF THE UNITED STATES.....	4-38

4.4.1	Field Evaluation Summary.....	4-40
4.4.2	Wetlands and Other Waters of the United States Vegetation Summary.....	4-49
4.4.3	Forest and Woodland Palustrine Forested	4-50
4.4.4	Shrubland	4-50
4.4.5	Herbaceous Palustrine Emergent.....	4-50
4.4.6	Wetlands Soil Summary	4-50
4.5	NOXIOUS WEEDS AND INVASIVE NONNATIVE PLANT SPECIES	4-50
4.6	PROTECTED NATIVE PLANTS	4-51
4.7	WILDLIFE AND WILDLIFE HABITAT	4-51
4.7.1	Introduction.....	4-51
4.7.2	Wildlife and Habitat Overview.....	4-52
4.8	SPECIES GROUPS AND HABITAT AFFINITY	4-58
4.8.1	Mammals	4-58
4.8.2	Birds	4-58
4.8.3	Reptiles and Amphibians	4-60
4.8.4	Fish.....	4-60
4.8.5	Invertebrates	4-61
4.9	PREHISTORIC HUMANS, SPANISH SETTLEMENT, AND CURRENT LAND CONSERVATION	4-62
4.10	HABITAT RESTORATION AND MONITORING	4-64
5.	LIST OF PREPARERS.....	5-1
6.	REFERENCES	6-1

ATTACHMENTS

- A. Description of Federally Listed Species
- B. (b) (7)(E) Station Species Lists

FIGURES

4-1.	Representative Photographs of One-seed Juniper Habitat	4-4
4-2.	Representative Photographs of Fremont Cottonwood – Goodding Willow Habitat.....	4-6
4-3.	Representative Photographs of Fremont Cottonwood / Honey Mesquite Habitat.....	4-8
4-4.	Representative Photographs of Arizona Sycamore - Fremont Cottonwood / Honey Mesquite Habitat	4-8
4-5.	Representative Photographs of Ocotillo – Tarbush Habitat.....	4-10
4-6.	Representative Photographs of Limestone Ridge and Slope Habitat.....	4-11
4-7.	Representative Photographs of Steep Ridge and Slope Habitat	4-11
4-8.	Representative Photographs of Creosotebush - Mariola Desert Slopes and Plains Habitat.....	4-12
4-9.	Representative Photographs of Mariola - Volcanic Slope Habitat	4-13
4-10.	Representative Photographs of Creosotebush – Honey Mesquite Slope and Dissected Plain Habitat	4-14
4-11.	Representative Photographs of Creosotebush – Tarbush Desert Plain Habitat.....	4-15
4-12.	Representative Photographs of Tarbush Alluvial Fan Habitat.....	4-16
4-13.	Representative Photographs of Dwarf Shrub Dry Ridge Habitat	4-17
4-14.	Representative Photographs of Honey Mesquite – Whitethorn Bajada Habitat	4-18
4-15.	Representative Photographs of Honey Mesquite / Hook Threawn Ridge Habitat	4-18
4-16.	Representative Photographs of Honey Mesquite – Tarbush Slope and Braided Wash Habitat.....	4-19
4-17.	Representative Photographs of Honey Mesquite – Four-wing Saltbush Alkaline Flat, Volcanic Slope, and Bladed Habitat	4-20
4-18.	Representative Photographs of Honey Mesquite – Alkali Sacaton Creek and Wash Habitat.....	4-21
4-19.	Representative Photographs of Honey Mesquite with Little Understory Cover Habitat.....	4-22
4-20.	Representative Photograph of Honey Mesquite – Littleleaf Sumac Desert Wash Habitat.....	4-23
4-21.	Representative Photographs of Seepwillow – Burro Bush (b) (7)(E) Habitat	4-24
4-22.	Representative Photographs of Alkali Sacaton Desert Wash Habitat	4-25
4-23.	Representative Photographs of Wild Barley/ Honey Mesquite Small Drainage Habitat.....	4-25
4-24.	Representative Photographs of Hook Threawn Flats Habitat.....	4-26
4-25.	Representative Photographs of Desert Marigold Flats Habitat.....	4-27
4-26.	Representative Photographs of Slimleaf Bursage – Common Sunflower Cienega Habitat and Existing Vehicle Barrier	4-28
4-27.	Representative Photographs of Russian-thistle Playa Habitat	4-29
4-28.	Representative Photographs of Common Cocklebur Excavation Habitat.....	4-29
4-29.	Representative Photographs of Bermuda Grass Irrigated Pasture Habitat	4-30

TABLES

2-1.	Tactical Infrastructure Sections, Tucson Sector, (b) (7)(E) Station	2-1
2-2.	Federal Threatened and Endangered Species and Arizona Wildlife Species of Concern Occurring Within (b) (7)(E)	2-3
4-1.	Crosswalk Relationship of USGS GAP Map Units and USFWS Habitat Types with NVCS Ecological Systems and Vegetation Alliances	4-2
4-2.	Plant Species List, Relative Abundance in the Survey Corridor, and Habitat for (b) (7)(E) Station, FV-1b	4-31
4-3.	Delineated Acreages, General Locations and Potential Impact Acreage of Wetlands and Other Waters of the United States in (b) (7)(E) AZ section FV-1b	4-42
4-4.	Wildlife Species Observed Within the Survey Corridor, Staging Areas, and Associated Access Roads	4-52

1. INTRODUCTION

This Biological Survey Report (BSR) synthesizes information collected from a variety of literature sources and field surveys to describe the biological resources within the survey corridor; provides supporting information from the Project region; allows evaluation within the Project Environmental Stewardship Plan (ESP) of the potential effects of the Project on those biological resources; and provides the basis of recommendations for avoidance or reduction of those effects using mitigation including best management practices (BMPs). Information was gathered from publicly available literature, data provided by relevant land management agencies such as the U.S. Fish and Wildlife Service (USFWS) (b) (7)(E) review of aerial photography and U.S. Geological Survey (USGS) topographic maps, data from the State of Arizona, data from NatureServe, and field surveys of the survey corridor conducted in January and May 2008. Of particular importance were data from (b) (7)(E) whose entire southern boundary with Mexico comprises a portion of the survey corridor.

This BSR supports the Environmental Stewardship Plan by providing information on biological resources potentially affected by impacts resulting from the construction, operation, and maintenance of the tactical infrastructure. The BSR was prepared as an independent document that is an appendix to the Environmental Stewardship Plan developed for this Project. The survey corridor is approximately (b) (7)(E) in length, approximately 760.5 acres within (b) (7)(E) (b) (7)(E) area. In total, approximately 700.6 acres of mostly native vegetation providing wildlife habitat occurs in the survey corridor. The remaining area (59.9 acres) supports land use in the form of unvegetated desert wash bottoms, irrigated pasture, and roads and trails.

Herbaceous vegetation (i.e., desert grasslands, forblands, emergent wetlands) composes approximately 15.5 acres. Shrublands (i.e., dwarf, short, and tall) compose approximately 587.8 acres. Forests and woodlands comprise 97.3 acres of vegetation cover. The vegetation represents a combination of mostly native Chihuahuan Desert shrublands that have become established in sparse to dense stands on ridges, slopes, alluvial fans, outwash plains, and along desert washes, draws, creeks, and springs.

THIS PAGE INTENTIONALLY LEFT BLANK

2. PROJECT DESCRIPTION

U.S. Customs and Border Protection (CBP) proposes to construct, maintain, and operate tactical infrastructure (b) (7)(E) and associated access roads along the U.S./Mexico border in the U.S. Border Patrol (USBP), Tucson Sector, (b) (7)(E) Station, Arizona. The locations of tactical infrastructure are based on a USBP Tucson Sector assessment of local operational requirements (b) (7)(E). Tactical infrastructure will be constructed in Section FV-1b along the international border in (b) (7)(E) County, Arizona (see **Table 2-1**).

Table 2-1. Tactical Infrastructure Sections, Tucson Sector, (b) (7)(E) Station

General Location	Land Ownership	Type of Tactical Infrastructure	Length of New Fence Section/Length of Construction Access Roads
(b) (7)(E)	Private, Public, USWFS, BLM, Arizona State Lands	Primary vehicle fence, access roads	(b) (7)(E)
	Private, BLM	Primary vehicle fence, access roads	
Total			(b) (7)(E)

The vehicle barrier will be constructed in (b) (7)(E) sections (b) (7)(E) partially within the Roosevelt Reservation, land reserved in 1907 within 60 feet of the international boundary between the U.S. and Mexico along California, Arizona, and New Mexico. (b) (7)(E)

Constructing the vehicle barrier will require improving or building (b) (7)(E) access roads, totaling (b) (7)(E) miles on USFWS, Bureau of Land Management (BLM), and private land parcels.

The final design will be prepared by a design/build contractor with oversight from the U.S. Army Corps of Engineers (USACE). Design criteria that have been established based on USBP operational needs require minimum standards for vehicle barriers, as follows: (1) capable of (b) (7)(E); (2) capable of (b) (7)(E); (3) designed to (b) (7)(E); (4) designed to (b) (7)(E); (b) (7)(E) (5) not impede the natural flow of surface water; and (6) to be as aesthetically pleasing as possible.

The area of impact for barrier construction is approximately (b) (7)(E) wide along the entire survey corridor, with wider but temporary impacts occurring at staging areas for construction materials and vehicles. Vegetation removal and land clearing/grading activities may occur on an as-needed basis.

2.1 Survey Methods

To provide flexibility in placement of tactical infrastructure within the survey corridor and to ensure consideration of potential impacts due to construction, patrol, and maintenance, surveys were conducted in an area extending (b) (7)(E) (b) (7)(E) on the north side (i.e., the side away from the international border) of the individual tactical infrastructure sections and extending at least (b) (7)(E) past the ends of the section (a total of 760.5 acres). Along access roads, the survey was conducted (b) (7)(E) corridor. The areas thus defined are referred to hereafter as the "survey corridor."

Field investigations of the survey corridor were conducted by biologists of engineering-environmental Management, Inc. (e²M): (b) (6) (senior ecologist), (b) (6) (staff biologist), (b) (6) (staff botanist), and (b) (6) (senior wetlands biologist). The January, April/May, and June 2008 surveys examined the survey corridor on January 14, from April 28 through May 2, and from June 9 through 13, 2008. A Contractor Site Visit Request Form was approved by the USACE, with assistance from the USFWS, (b) (7)(E) Manager, (b) (6), and USBP escorts. A second field visit was conducted on August 13, 2008 to assess a modification that exceeded the original survey corridor. The field investigation for this smaller area was conducted by senior e²M biologists (b) (6) and (b) (6).

Due to the schedule requirements for acquiring field information, e²M assigned senior and staff ecologists/biologists familiar with the USBP Projects, reporting process, vegetation, wetlands/waters of the United States, wildlife habitat classification and mapping protocols, and field sampling methods to intuitively examine the landscape and survey corridor for the (b) (7)(E) length. Further, senior e²M natural resources staff used USFWS species lists and comprehensive conservation planning data (USFWS 1995) to ensure accurate identification of plant species and competent surveys for rare plants, wildlife, and potential habitat. The surveys were controlled, in that right-of-entry (ROE) was approved for the entire corridor and access road widths, and survey crews were in contact with USBP operations. While on (b) (7)(E), field biologists were accompanied by a USFWS law enforcement officer and were met by the Refuge Manager for sensitive site overviews. Investigations included preparing lists of observed plant and wildlife species; an assessment of habitat and surveys for rare plant and wildlife species; landscape photography points; observation points recording dominant species, location, cover, environmental conditions, and photodocumentation; determination of potential wetlands and other waters of the United States for future research; locations of major desert

washes; and general note taking of natural resources, cultural resources, and other Environmental Stewardship Plan reporting needs.

Biologists walked the entire survey corridor, including all the access road corridors and staging areas. The survey team conducted reconnaissance level surveys on areas of land use (irrigated pasture and sites devoid of vegetation including playas, desert wash bottoms, and access roads) and examined in detail areas containing unique species compositions or habitat that might be conducive to sensitive species (desert grasslands, shrublands, riparian woodlands and forests, emergent wetlands, etc.). Observation data (Universal Transverse Mercator [UTM] coordinates, photographs, field notes, environmental information, vegetation structure, and plant community composition) were recorded at regular intervals along the corridor where vegetation occurred as homogenous stands and also where plant communities presented substantial shifts in species composition. These data were used to generate a vegetation classification and map to facilitate delineation of habitat types, analyses of potential sensitive species occurrences, and analyses of potential Project impacts on biological resources. The botanist and wildlife biologist specifically examined habitats to determine the presence of state- and Federal-listed species (see **Table 2-2**). Descriptions of the federally listed species developed by NatureServe (2008) are provided in **Attachment A**.

Table 2-2. Federal Threatened and Endangered Species and Arizona Wildlife Species of Concern Occurring Within (b) (7)(E) County

Common Name	Scientific Name	Federal Status	State Status
FISH AND INVERTEBRATES			
Beautiful shiner	<i>Cyprinella Formosa</i>	LT	WSC
Desert pupfish	<i>Cyprinodon macularius</i>	LE	---
Gila chub	<i>Gila intermedia</i>	LE	WSC
Yaqui chub	<i>Gila purpurea</i>	LE	WSC
Yaqui catfish	<i>Ictalurus pricei</i>	LT	WSC
Spikedace	<i>Meda fulgida</i>	LT	---
Gila topminnow	<i>Poeciliopsis occidentalis occidentalis</i>	LE	---
Yaqui topminnow	<i>Poeciliopsis occidentalis sonoriensis</i>	LE	WSC
Loach minnow	<i>Tiaroga cobitis</i>	LT	---
Huachuca springsnail	<i>Pyrgulopsis thompsonii</i>	C	---
AMPHIBIANS			
Sonora tiger salamander	<i>Ambystoma tigrinum stebbinsi</i>	LE	---

Common Name	Scientific Name	Federal Status	State Status
Ramsey Canyon leopard frog	<i>Lithobates subaquavocalis</i>	CA	---
Chiricahua leopard frog	<i>Rana chiricahuensis</i>	LT	WSC
REPTILES			
New Mexico ridge-nosed rattlesnake	<i>Crotalus willardi obscurus</i>	LT	—
BIRDS			
Western yellow-billed cuckoo	<i>Coccyzus americanus occidentalis</i>	C	WSC
Southwestern willow flycatcher	<i>Empidonax traillii extimus</i>	LE	WSC
Northern aplomado falcon	<i>Falco femoralis septentrionalis</i>	LT, XN	---
American peregrine falcon	<i>Falco peregrinus anatum</i>	SC	WSC
California brown pelican	<i>Pelecanus occidentalis californicus</i>	PDL	---
Mexican spotted owl	<i>Strix occidentalis lucida</i>	LT	WSC
MAMMALS			
Ocelot	<i>Leopardus pardalis</i>	LE	---
Lesser long-nosed bat	<i>Leptonycteris curasoae</i>	LE	WSC
Jaguar	<i>Panthera onca</i>	LE	WSC
PLANTS			
Cochise pincushion cactus	<i>Coryphantha robbinsorum</i>	LT	HS
Lemmon fleabane	<i>Erigeron lemmonii</i>	FC	HS
Huachuca water umbel (Cienega false rush)	<i>Lilaeopsis schaffneriana</i> var. <i>recurva</i>	LE	HS
Madrean ladies'-tresses	<i>Spiranthes delitescens</i>	LE	HS

Source: AZFGD 2008b, USFWS 2008a

Notes: LE = Listed Endangered; LT = Listed Threatened; FC = Federal Candidate; C = Candidate; CA = Conservation Agreement; PDL = Proposed for Delisting; PR = Protected; SC = Species of Concern; XN: Experimental Population; WSC = Wildlife of Special Concern in Arizona; HS = Highly Safeguarded Protected Native Plants (no collection allowed)

2.2 Arizona Game and Fish Department; Arizona Natural Heritage Program, Heritage Data Management System

The Arizona Heritage Data Management System (HDMS) was established to collect, synthesize, and catalog information concerning the distribution and

occurrence of species and habitats in need of special attention (Arizona Game and Fish Department [AZGFD] 2008a, 2008b). It is part of a global network of 80 Natural Heritage Programs and Conservation Data Centers. The HDMS is Arizona's most comprehensive source of information related to rare, threatened, and endangered animals, plants, exemplary natural communities, and other significant features. The data are publicly available from which to make prudent decisions weighing future development, economic growth, and environmental integrity (AZGFD 2008a, 2008b). While these data are continually updated, there are gaps in coverage and species information due to lack of access to land for inventory, data from many sources, and a lack of staff and resources to collect and process data for all rare and significant resources. To request information from the HDMS online, access: http://www.azgfd.gov/w_c/edits/hdms_natural_heritage.shtml.

For the survey corridor, HDMS data were used to assist with the evaluation of environmental impacts of the vehicle barrier section under consideration. The interpretation and extrapolation of the data included consideration that: (1) data gaps possibly occur because of the availability of data extraction from public information sources, (2) species and geographic coverage focused on the most rare species and ecosystems, and (3) the potential lack of precise locality data in some secondary sources exists. Because of the large proportion of public land versus private land in Arizona, the HDMS includes a representative inventory of rare resources in the state. It is based on the best data available to the AZGFD in terms of rare species locations and distributions.

The (b) (7)(E) County list of rare species was acquired from HDMS and consolidated into **Table 2-2**. The county lists include wildlife species of special concern in Arizona and highly safeguarded plant species. In general, species that appear on county lists do not all share the same probability of occurrence within a county (e.g., some species are migrants or wintering residents and a few species might be historic or considered extirpated within a county).

THIS PAGE INTENTIONALLY LEFT BLANK

3. ENVIRONMENTAL SETTING

The survey corridor climate is semiarid within the Xeric Climatic Region as described in Robinson et al. (2006). This region is characterized by deep, northwest-trending, alluvium-filled basins separated by linear mountain ranges (basin and range lowlands). Relatively recent volcanic activity was evident with many slopes covered by gravel and cobble of volcanic origin. Land surface elevations range from approximately 3,720 feet above mean sea level to more than 5,000 feet above mean sea level in the immediate Project region. Precipitation typically increases and temperatures decrease with increasing altitude in the Xeric Climatic Region during all seasons of the year. Low rainfall and high temperatures are characteristic of the basin and range lowlands (e.g., summers are long and hot and winters are short, dry, and cold and can include brief periods when temperatures are below freezing) (Robinson et al. 2006, Bailey 1995). Many of the streams in the Xeric Climatic Region are intermittent or ephemeral (i.e., more than 250 days annually of no flow), but can have high flow in response to intense thunderstorms.

The general climatic summary records for (b) (7)(E), Arizona (Station (b) (7)(E)), have been prepared from 1948 to 2007 data (Western Regional Climate Center [WRCC] 2008). Average minimum temperatures (b) (7)(E) range from a low of 29 degrees Fahrenheit (°F) in December and January to 65°F in July, and average high temperatures range from 63°F in December and January to 96°F in June (WRCC 2008). The lowest temperature recorded was 2°F on December 8, 1978, and the highest temperature recorded was 109°F on June 26, 1990. The average annual precipitation is 14.2 inches, over half of which falls in July, August, and September during the summer monsoon season. A long growing season is experienced for the Project region, averaging 240 frost-free days (WRCC 2008). The evaporation rate during the summer season is high, about twice the precipitation amount, and averages about 70 inches annually in (b) (7)(E).

Upland soils within the survey corridor are classified within the Bonita-Sontag Association and valley floor soils have been classified within the Karro Association (USFWS 1995). Karro Association soils are typically deep and well-drained and formed in old alluvium from mixed igneous and sedimentary rocks on alluvial fans and uplands; they include (1) Karro Loam, (2) Bonita Clay, (3) Bonita Cobbly Clay, and (4) Riggs. The Bonita-Sontag Association soils are typically shallow and well-drained and formed in mixed slope alluvium from sedimentary and igneous rocks; they include (1) Stronghold, (2) Mabray, (3) Lampshire-Ridgelite, and (4) Gadwell-Caralampi Complex.

The vegetation of the basin and range lowlands of (b) (7)(E) Arizona has generally been classified under the Dry Domain (Map Unit 300), Tropical/Subtropical Desert Division (Map Unit 320) of Bailey (1995). The survey corridor is more finely classified by Bailey (1995) as the Chihuahuan Desert Province (Map Unit 321). The Arizona Gap Project (Bennett et al. 2004)

provided discussion and described plant geography to vegetation series using topographic features, climate, vegetation types, and terrestrial vertebrates. This system placed the survey corridor generally in the Nearctic Upland; Warm Temperate Desertland; Chihuahuan Desertscrub classification. Vegetation series that were described and are applicable to the survey corridor included (1) Creosotebush-Tarbush Series; (2) Mesquite Series; (3) Whitethorn Series; (4) Mixed Scrub Series, and (5) Scrub Grassland Series (Bennett et al. 2004).

4. BIOLOGICAL RESOURCES

4.1 Vegetation Classification

The USGS (Bennett et al. 2004) recognizes nine Nearctic Upland and Nearctic Wetland vegetation mapping units in the (b) (7)(E) Arizona, vicinity using a combination of plant species dominance, wildlife use, topography, hydrology, and geology. The vegetation series that are associated with the survey corridor include (1) Warm Temperate Grassland, Scrub-Grassland (Semidesert), Tobosa Grass-Scrub Series; (2) Warm Temperate Scrub-Grassland (Semidesert), Sacaton-Scrub Series; (3) Warm Temperate Desertland, Chihuahuan Desertscrub, Creosotebush-Tarbrush Series; (4) Warm Temperate Desertland, Chihuahuan Desertscrub, Whitethorn Series; (5) Warm Temperate Desertland, Chihuahuan Desertscrub, Mesquite Series; (6) Warm Temperate Desertland, Chihuahuan Desertscrub, Mixed Scrub Series; (7) Tropical-Subtropical Swamp Riparian and Oasis Forests, Sonoran Riparian and Oasis Forest, Cottonwood-Willow Series; (8) Tropical-Subtropical Swamp and Riparian Scrub, Sonoran Deciduous Swamp and Riparian Scrub, Mixed Scrub Series; and (9) Tropical-Subtropical Marshland, Sonoran Interior Marshland, Cattail Series. The entire corridor was predominantly characterized by Chihuahuan Desertscrub vegetation series.

NatureServe (2008) has defined ecological systems to represent recurring groups of biological communities that are found in similar physical environments and are influenced by similar dynamic ecological processes such as drought, fire, or flooding. Ecological systems represent classification units that are readily identifiable by conservation and resource managers in the field. The ensuing vegetation description for the survey corridor was prepared in the framework of ecological systems that include (1) North American Warm Desert Riparian Woodland and Shrubland (CES302.753), (2) Apacherian-Chihuahuan Mesquite Upland Scrub (CES302.733); (3) Chihuahuan Mixed Desert and Thorn Scrub (CES302.734), (4) Apacherian-Chihuahuan Mesquite Upland Scrub (CES302.733), (5) North American Warm Desert Riparian Mesquite Bosque (CES302.752), (6) North American Warm Desert Wash (CES302.755), (7) Apacherian-Chihuahuan Semi-Desert Grassland and Steppe (CES302.735), (8) North American Warm Desert Cienega (CES302.747), and (9) North American Warm Desert Playa (CES302.751). **Table 4-1** provides a crosswalk between the biotic communities described by the USGS and the ecological systems of NatureServe (2008).

Classification of existing vegetation within the survey corridor was achieved by accessing the survey corridor, access roads, and staging areas as planned, sampling observation points, and relating them to the NatureServe Explorer classification database directly or as provisional types (NatureServe 2008). At the coarsest level, the nine above-named ecological systems were determined and local vegetation types described using the national system.

Table 4-1. Crosswalk Relationship of USGS GAP Map Units and USFWS Habitat Types with NVCS Ecological Systems and Vegetation Alliances

Ecological System (NatureServe 2008) Provisional Vegetation Alliance	Vegetation Structure and Series	Habitat Types
Madrean Pinyon-Juniper Woodland - One-seed Juniper Wooded Herbaceous	Madrean Evergreen Forest - Oak-Pine Series	Chihuahuan Desert Scrub
North American Warm Desert Riparian Woodland and Shrubland - Fremont Cottonwood – Goodding Willow Forest - Fremont Cottonwood / Honey Mesquite Forest - Arizona Sycamore – Fremont Cottonwood / Honey Mesquite Woodland	Sonoran Riparian and Oasis Forest - Cottonwood-Willow Series Mogollon Mixed Broadleaf - Mixed Broadleaf Series Sonoran Deciduous Swamp and Riparian Scrub - Mixed Scrub Series Sonoran Interior Marshland - Cattail Series	Riparian Forest/Woodland Riparian Scrub Mesquite Bosque Marshland
Apacherian-Chihuahuan Mesquite Upland Scrub - Ocotillo – Tarbush Shrubland - Mortonia – Mariola Shrubland - Whitethorn – Mariola Shrubland	Chihuahuan Desertscrub - Whitethorn Series Chihuahuan Desertscrub - Mixed Scrub Series	Chihuahuan Desert Scrub
Chihuahuan Mixed Desert and Thorn Scrub - Creosotebush – Mariola Shrubland - Mariola Dwarf-shrubland - Creosotebush – Honey Mesquite Shrubland - Creosotebush – Tarbush Shrubland - Tarbush Shrubland - Shrubby Coldenia – Engelmann Prickly-pear Dwarf-shrubland	Chihuahuan Desertscrub - Creosotebush-Tarbush Series Chihuahuan Desertscrub - Mixed Scrub Series	Chihuahuan Desert Scrub
Apacherian-Chihuahuan Mesquite Upland Scrub - Honey Mesquite – Whitethorn Bajada Shrubland - Honey Mesquite / Hook Threeawn Shrubland - Honey Mesquite – Tarbush Shrubland	Chihuahuan Desertscrub - Mesquite Series	Chihuahuan Desert Scrub

Ecological System (NatureServe 2008) Provisional Vegetation Alliance	Vegetation Structure and Series	Habitat Types
North American Warm Desert Riparian Mesquite Bosque <ul style="list-style-type: none"> - Honey Mesquite – Four-wing Saltbush Shrubland - Honey Mesquite / Alkali Sacaton Woodland and Shrubland - Honey Mesquite Sparse Understory Woodland and Shrubland - Honey Mesquite – Littleleaf Sumac Shrubland 	Chihuahuan Desertscrub - Mesquite Series	Chihuahuan Desert Scrub Mesquite Bosque Riparian Scrub
North American Warm Desert Wash <ul style="list-style-type: none"> - Seepwillow – Burro Bush Shrubland - Alkali Sacaton Herbaceous Vegetation - Wild Barley / Honey Mesquite Shrub Herbaceous Vegetation 	Scrub-Grassland - Sacaton-Scrub Sonoran Deciduous Swamp and Riparian Scrub - Mixed Scrub Series	Chihuahuan Desert Scrub Desert Grassland Riparian Scrub
Apacherian-Chihuahuan Semi-Desert Grassland and Steppe <ul style="list-style-type: none"> - Hook Threawn Herbaceous Vegetation - Desert Marigold Herbaceous Vegetation 	Scrub Grassland	Desert Grassland

Note: NVCS = National Vegetation Classification System.

A finer level of classification equaling or approximating the vegetation alliance level of the National Vegetation Classification System (NVCS) (NatureServe 2008) was used to prepare the plant community discussions under each ecological system. Vegetation stands and patches that are generally unclassified in the current system and sampled within the Project area typically consisted of nonnative species including Bermuda Grass Herbaceous Vegetation, Russian-thistle Herbaceous Vegetation, and Common Cocklebur Herbaceous Vegetation.

Habitats observed, sampled, and photographed within the survey corridor range from upland mixed desert scrub and thorn-scrub throughout the alignment to riparian woodland and forest stands (b) (7)(E). Much of the vegetation cover along the (b) (7)(E) fence section consists of native shrublands characterized by honey mesquite, creosotebush, tarbush, whitethorn, shrubby coldenia, mortonia, and ocotillo; vegetation cover occupies approximately 88 percent of the corridor. Development is limited to one irrigated pasture and existing roads and trails; these land uses occupy approximately 12 percent of the corridor.

A brief description of each plant community observed within the section (FV-1b) is provided herein; they are distinguished using the NatureServe Vegetation Alliance level of classification or an approximation. Each community is illustrated and supported by representative ground photographs and foliar cover information for dominant and characteristic plant species.

4.1.1 Madrean Pinyon-Juniper Woodland Ecological System (CES305.797)

One-seed Juniper / Whitethorn Wooded Shrubland

The uplands and associated small drainages within (b) (7)(E) occupied 25.1 acres of the survey corridor and supported 2 to 4 meter (m) tall one-seed juniper that ranged in cover from 3 to 5 percent (see **Figure 4-1**). The tall shrub layer provided low to moderate cover and includes whitethorn, catclaw acacia, littleleaf sumac, and honey mesquite. The short and dwarf-shrub layers contribute low cover, up to 5 percent cover and include mortonia, creosotebush, prickly-pear cactus, and agave. The herbaceous layer provides low to moderate cover, predominantly from the bunchgrasses tobosa, black grama, bush muhly, and three-awn.



Figure 4-1. Representative Photographs of One-seed Juniper Habitat

4.1.2 North American Warm Desert Riparian Woodland and Shrubland Ecological System (CES302.753)

Fremont Cottonwood – Goodding Willow Forest

(b) (7)(E) supports a forest stand (b) (7)(E) at the international border which occupied 0.9 acre of the survey corridor. The draw was ponded at the international border and upstream approximately 50 m had been hardened using gabions filled with rocks to reduce erosion (see **Figure 4-2**). (b) (7)(E) measured approximately 60 m from bank-to-bank and had become incised from 5 to 8 m deep; the banks are nearly vertical. Debris lines on the trees suggest flows that could exceed 1.5 m in depth. The draw has perennial springs and near-to-surface ground water which provides habitat for

riparian and wetland plant species. Fremont cottonwood trees up to 30 m tall have become established on the banks and first terrace of the draw and provide approximately 90 percent cover (see **Figure 4-2**). These trees were mature with large diameters-at-breast-height (60 to 70 centimeters [cm]). An understory layer of Goodding willow trees provided approximately 20 percent cover and attained heights of approximately 8 m tall. Honey mesquite trees provided low to moderate cover (10 to 15 percent cover) on the first terrace adjacent to the Fremont cottonwood stand. In the densest portion of this linear stand the understory was composed of leaf litter, but where canopy openings occurred the emergent wetland species southern cattail, three-square bulrush, and mixed graminoids provided moderate cover (up to 25 percent cover). Approximately (b) (7)(E) was a small stand of Fremont cottonwood trees with nearly 100 percent cover by three-square bulrush in the understory that was fed by a spring. (b) (7)(E) provided rare and valuable wildlife habitat with the tallest structural component of any plant community along this portion of the international border. Its waters also support three endangered fish species. The small Fremont cottonwood stand to the west provided active nest sites for the gray hawk, a pair of which was in residence.



Figure 4-2. Representative Photographs of Fremont Cottonwood – Goodding Willow Habitat

Apparently, (b) (7)(E) did not exist in the 1850s, but by the 1890s it occurred as a creek lined with cottonwood trees (Lanning 1981 in USFWS 1994). Later, it became a ditch approximately 3 to 5 m deep, 5 to 25 m in width, and was typically dry.

Fremont Cottonwood / Honey Mesquite Forest

(b) (7)(E) supports a forest stand (b) (7)(E) at the international border that occupied 3.6 acres within the survey corridor. The wash has a sandy bottom, is approximately 40 m wide, has incised up to 4 m deep within nearly vertical banks, and has sufficient surface flows and near-to-surface groundwater to support riparian plant species (see **Figure 4-3**). Debris lines captured on the trees suggest flows that could exceed 1 m in depth. Fremont cottonwood trees up to 25 m tall have become established in the wash bottom and on banks and provide approximately 70 percent canopy cover. These trees are approximately 15 years of age and have diameters-at-breast-height of approximately 35 to 40 cm. Higher on the banks and on the first wash terrace, honey mesquite trees to 15 m tall provide moderate cover, up to 20 percent cover. Giant dropseed and alkali sacaton contributed low cover on the wash banks and first terrace. This is a rare and valuable wildlife habitat, with the tallest structural component of any plant community along this portion of the international border.

Arizona Sycamore - Fremont Cottonwood / Honey Mesquite Woodland

The wash located in (b) (7)(E) at the international border provides habitat for a sparse woodland community on the banks and terraces, occupying 14.8 acres within the survey corridor. The wash has incised approximately 4 m deep, contained a barren sandy or gravelly channel. Cobble was deposited on point bars and terraces; it ranged from 10 to 20 m wide (see **Figure 4-4**). Arizona sycamore and Fremont cottonwood trees up to 15 m tall have become established on the banks and first terraces of the moderately large desert wash and provide low to moderate cover, approximately 10 to 20 percent and 1 to 5 percent, respectively. The short-statured honey mesquite trees occurred as understory to the taller trees or formed monotypic stands or clumps on the desert wash banks and terraces; they provided low to moderate cover from 10 to 20 percent cover. Additional understory trees included hackberry, green ash, one-seed juniper, and oak which provided low cover. The tall shrub layer ranged from 2 to 5 m tall and was characterized by honey mesquite which provided moderate cover, from 10 to 30 percent cover. The short shrub layer provided sparse cover and was characterized by littleleaf sumac, wait-a-minute, wolfberry, and burro bush. The herbaceous layer was comprised of grasses providing low to moderate cover, from 10 to 25 percent cover and included sideoats grama, deer grass, sand dropseed, tobosa, and big dropseed.



Figure 4-3. Representative Photographs of Fremont Cottonwood / Honey Mesquite Habitat



Figure 4-4. Representative Photographs of Arizona Sycamore - Fremont Cottonwood / Honey Mesquite Habitat

4.1.3 Apacherian-Chihuahuan Mesquite Upland Scrub Ecological System (CES302.733)

Ocotillo – Tarbush Shrubland

(b) (7)(E) supported tall shrub stands on rocky outcrops, cobbly and gravelly colluvial deposits, and outwash fans on 32.8 acres of the survey corridor (see **Figure 4-5**). The tall shrub layer was 3 to 5 m high and characterized by ocotillo that provided low to moderate cover (5 to 25 percent cover), typically with a short shrub understory of tarbush that provided low cover (5 to 15 percent cover). Associated short and dwarf-shrubs provided sparse cover and included mariola, yucca, little-head snakeweed, rainbow cactus, and wait-a-minute. South-facing rock outcrops within this type supported dense patches of little bluestem and sparse cover of tobosa and threeawn.

Mortonia - Mariola Shrubland

Limestone outcrops and colluvial slopes in the vicinity of (b) (7)(E) and small drainage systems in (b) (7)(E) supported unique short shrub stands on 26.6 acres within the survey corridor (see **Figure 4-6**). The short shrub layer was characterized by mortonia that provided low to moderate cover (5 to 15 percent cover) in association with the dwarf-shrub mariola that contributed low to moderate cover (5 to 20 percent cover). The tall shrub layer was often present, provided sparse to low cover, and included ocotillo, whitethorn, squawbush, and one-seed juniper. The remaining short and dwarf-shrub layers were diverse, contributed sparse to moderate cover, and included creosotebush, tarbush, althorn, agave, yucca, sotol, Engelmann prickly-pear, and shrubby coldenia. The herbaceous layer contributed sparse cover and included hook threeawn, tobosa, and fluffgrass.





Figure 4-5. Representative Photographs of Ocotillo – Tarbush Habitat

Whitethorn - Mariola Shrubland

(b) (7)(E) supported short and dwarf-shrub stands on the driest exposures and steepest ridges (see **Figure 4-7**). The dwarf-shrub layer was characterized by mariola which provided low cover, up to 10 percent cover, and the tall shrub layer was characterized by low cover of whitethorn (5 percent cover). Associated tall and short shrubs that contributed sparse to low cover included ocotillo, yucca, and Palmer agave. The herbaceous layer provided sparse cover and was characterized by tobosa and fluffgrass. This type occurred on the edge of the survey corridor and occupied 0.7 acre.





Figure 4.6. Representative Photographs of Limestone Ridge and Slope Habitat



Figure 4-7. Representative Photographs of Steep Ridge and Slope Habitat

4.1.4 Chihuahuan Mixed Desert and Thorn Scrub (CES302.734)

Creosotebush – Mariola Shrubland

Gentle slopes and alluvial outwash plains, typically gravelly and sandy in texture, supported consistent dominance by short and dwarf shrubs of this type which occupied 234.7 acres of the survey corridor (see **Figure 4-8**). The short shrub creosotebush and dwarf-shrub mariola characterized this type and each provided from 5 to 25 percent cover within stands. The tall shrub layer contributed sparse to low cover (1 to 10 percent cover) and included ocotillo, whitethorn, and honey mesquite. The remaining short and dwarf-shrub layers contributed sparse to low cover (1 to 10 percent cover) and included tarbush, yucca, Engelmann prickly-pear, cane cholla, and shrubby coldenia. The herbaceous layer was low in terms of species diversity, contributed sparse cover or was absent, and was characterized by fluffgrass, tobosa, and sprangletop.




Figure 4-8. Representative Photographs of Creosotebush - Mariola Desert Slopes and Plains Habitat

Mariola Dwarf-shrubland

This type is unique and had become established on a south-facing slope armored by volcanic rocks, occupying 2.4 acres within the survey corridor (see **Figure 4-9**). The dwarf-shrub mariola characterized the site, providing 20 percent cover. Associated tall and short shrubs provided sparse cover and included creosotebush and honey mesquite. Herbaceous vegetation was nearly absent; the short bunchgrass tobosa contributed sparse cover.




Figure 4-9. Representative Photographs of Mariola - Volcanic Slope Habitat

Creosotebush – Honey Mesquite / Tobosa Shrubland

Gentle to steep slopes, volcanic cobble exposures, and dissected plains supported mixtures of tall and short shrubs that were relatively consistent on 73.0 acres throughout the survey corridor (see **Figure 4-10**). This type is characterized by creosotebush and honey mesquite tall shrubs that each range in cover from 5 to 15 percent. Associated tall shrubs provided sparse cover and included littleleaf sumac, tarbush, four-wing saltbush, whitethorn, and shrubby coldenia. The herbaceous layer was patchy in distribution, contributed sparse to low cover (2 to 12 percent cover), and included tobosa and black grama.

Creosotebush – Tarbush Shrubland

Broad, gravelly plains supported moderate stands of this type (b) (7)(E) where 48.1 acres of the survey corridor supported this type (see **Figure 4-11**). Creosotebush short shrubs provided low cover, up to 15 percent, and tarbush short shrubs provided sparse cover (up to 4 percent cover) in these open stands. In one stand, the tall shrub ocotillo contributed sparse cover. Associated short and dwarf-shrubs contributed sparse cover and included whitethorn, mariola, soaptree yucca, and shrubby coldenia. The grasses bush muhly and tobosa provided sparse cover in one stand.




Figure 4-10. Representative Photographs of Creosotebush – Honey Mesquite Slope and Dissected Plain Habitat

Tarbush Shrubland

An individual sandy and gravelly alluvial fan was characterized by moderate cover (35 percent cover) of tarbush that occupied 1.1 acres of the survey corridor (see **Figure 4-12**). The remaining short and dwarf-shrub layers contributed sparse to low cover (up to 10 percent cover) and included whitethorn, honey mesquite, yucca, prickly-pear, and small-headed snakeweed. The herbaceous layer was absent from this stand.

Shrubby Coldenia – Engelmann Prickly-pear Dwarf-shrubland

This type became established on a dry ridge off a hill with gravelly, thin soils and occupied 1.6 acres in the survey corridor (see **Figure 4-13**). The dwarf-shrubs shrubby coldenia (12 percent cover) and Engelmann prickly-pear (8 percent cover) characterized the site and provided low to moderate cover. Sparse cover was provided by the short shrub creosotebush. The herbaceous layer contributed sparse cover and was characterized by fluffgrass.




Figure 4-11. Representative Photographs of Creosotebush – Tarbush Desert Plain Habitat

4.1.5 Apacherian-Chihuahuan Mesquite Upland Scrub Ecological System (CES302.733)

Honey Mesquite – Whitethorn Bajada Shrubland

Several small, gravel and cobble-covered hilltops and ridges occur (b) (7)(E) (b) (7)(E) and continue to (b) (7)(E) occupying 10.8 acres within the survey corridor (see **Figure 4-14**). These exposed sites support low cover, between 15 to 20 percent cover of short and dwarf-shrubs including honey mesquite, whitethorn acacia, four-wing saltbush, creosotebush, Mormon-tea or jointfir, and little-head snakeweed. No one shrub contributed more than 5 percent cover in these stands.




Figure 4-12. Representative Photographs of Tarbush Alluvial Fan Habitat

Honey Mesquite / Hook Threeawn Shrubland

A single stand of this vegetation type occurred on a cobbly ridge and occupied 7.7 acres of the survey corridor (see **Figure 4-15**). The tall shrub layer was characterized by 2 to 4 m high honey mesquite that provided moderate cover, up to 30 percent cover, and the short bunchgrass hook threeawn also provided moderate cover (up to 40 percent cover). The short shrub layer contributed low cover (less than 10 percent cover) and included whitethorn, yucca, and Engelmann prickly-pear. Tobosa provided low cover (5 percent cover) in the herbaceous layer.



Figure 4-13. Representative Photographs of Dwarf Shrub Dry Ridge Habitat

Honey Mesquite - Tarbush Shrubland

This vegetation type occurred on a cobbly volcanic slope and on sandy soils within a braided desert wash (b) (7)(E), occupying 5.5 acres in the survey corridor (see **Figure 4-16**). The tall shrub (to 4 m high) honey mesquite provided moderate cover (20 to 35 percent cover) and the short shrub tarbush provided low cover (4 percent cover). Whitethorn provided sparse cover in the tall shrub layer of one stand, and the short and dwarf-shrubs creosotebush, four-wing saltbush, Engelmann prickly-pear, and small-headed snakeweed contributed sparse to low cover. The herbaceous layer provided sparse cover by fluffgrass and alkali sacaton.



Figure 4-14. Representative Photographs of Honey Mesquite – Whitethorn Bajada Habitat



Figure 4-15. Representative Photographs of Honey Mesquite / Hook Threeawn Ridge Habitat




Figure 4-16. Representative Photographs of Honey Mesquite – Tarbush Slope and Braided Wash Habitat

4.1.6 North American Warm Desert Riparian Mesquite Bosque Ecological System (CES302.752)

Honey Mesquite – Four-wing Saltbush Shrubland

The relatively flat alkaline soils (b) (7)(E) and the volcanic cobble slopes (b) (7)(E) supported this type on 25.8 acres of the survey corridor (see **Figure 4-17**). Honey mesquite tall shrubs to 4 m high provided low to high cover (15 to 60 percent cover) and four-wing saltbush short shrubs provided 5 to 12 percent cover in the understory. In one stand, creosotebush and soaptree yucca provided sparse cover in the short shrub layer and small-headed snakeweed provided sparse cover in the dwarf-shrub layer. The herbaceous layer was characterized by sparse cover of tobosa and alkali sacaton. Within the refuge a strip of land adjacent to the border appeared to have been bladed or similarly treated to remove honey mesquite shrubland habitat. The treated area occupied 4.1 acres within the survey corridor and had recovered to sparse cover of honey mesquite, four-wing saltbush, and tobosa that in total provided less than 10 percent cover (see **Figure 4-17**).




Figure 4-17. Representative Photographs of Honey Mesquite – Four-wing Saltbush Alkaline Flat, Volcanic Slope, and Bladed Habitat

Honey Mesquite – Alkali Sacaton Woodland and Shrubland

This type occurs in drainage bottoms that include upper (b) (7)(E), large desert washes, and broad swales of the eastern Project terminus and occupies (b) (7)(E) within the survey corridor (see **Figure 4-18**). Honey mesquite trees or tall shrubs to 5 m tall occurred on the wash and swale banks or elevated sediment bars and provided low to moderate cover (15 to 40 percent cover). The tall bunchgrass alkali sacaton provided low to moderate cover (5 to 25 percent cover) on the wash and swale bottoms. In the remaining shrub layer, sparse cover was contributed by netleaf hackberry, four-wing saltbush, desert broom, yerba de pasmo, burro bush, littleleaf sumac, tarbush, and little-head snakeweed. In the remaining herbaceous layer, sparse cover was contributed by scratchgrass and Dakota verbena. This type was similar to the Honey Mesquite / Four-wing Saltbush Shrubland; however, the cover by four-wing saltbush short shrubs was sparse, typically 1 percent or less in terms of foliar cover.



Figure 4-18. Representative Photographs of Honey Mesquite – Alkali Sacaton Creek and Wash Habitat

Honey Mesquite Sparse Understory Woodland and Shrubland

Stands of honey mesquite with little understory vegetation have become established on the terraces of (b) (7)(E), and a relatively flat plain where they form moderately dense woodlands and tall shrublands (see **Figure 4-19**). Honey mesquite trees and tall shrubs range from 3 to 10 m in height, provided 40 to 85 percent cover, and occupied 20.7 acres of the survey corridor. The short shrub layer provides sparse cover and may include four-wing saltbush, creosotebush, and burro bush. The herbaceous layer provides sparse cover and includes tobosa, giant dropseed, six weeks fescue, Russian-thistle, and London rocket. One stand occurred near a windmill and exhibited signs of cattle foraging and resting under the tree canopies, likely on an annual basis.



Figure 4-19. Representative Photographs of Honey Mesquite with Little Understory Cover Habitat

Honey Mesquite – Littleleaf Sumac Shrubland

Small desert washes, approximately 5 to 15 m wide, totaled 7.3 acres in the survey corridor and will be crossed by access roads (see **Figure 4-20**). The washes were coequally characterized by 2 to 5 m tall honey mesquite and littleleaf sumac tall shrubs that together provided 30 percent cover. Short shrubs occupied the understory and small openings on wash banks and were characterized by low cover of tarbush (10 percent cover) and four-wing saltbush (4 percent cover). The herbaceous layer contributed sparse cover and included giant dropseed and annual desert holly.



Figure 4-20. Representative Photograph of Honey Mesquite – Littleleaf Sumac Desert Wash Habitat

4.1.7 North American Warm Desert Wash Ecological System (CES302.755)

Seepwillow – Burro Bush Shrubland

This type occurs just outside the survey corridor on a meander terrace (b) (7)(E) and occasionally to rarely experiences overbank flooding following precipitation events. (b) (7)(E) is very broad at this location, up to 100 m wide from bank to bank, has a sandy to gravelly channel that is mostly devoid of vegetation, and is incised up to 10 m deep (see **Figure 4-21**). The tall shrub yerba de pasmo (seepwillow) provides low cover (up to 10 percent cover) along with a few individual desert willow tall shrubs, which occupy a slightly elevated second terrace. The first terrace lay slightly above the channel and supported the short shrub burro bush which provided low cover up to 15 percent cover. Alkali sacaton, the medium-tall bunchgrass, provided sparse cover in the herbaceous layer.




Figure 4-21. Representative Photographs of Seepwillow – Burro Bush (b) (7)(E)
(b) (7)(E) Habitat

Alkali Sacaton Herbaceous Vegetation

A few larger desert washes and large swales (from 40 to 75 m wide) were characterized by 45 to 65 percent cover of alkali sacaton, a coarse bunchgrass that may be 1.5 m tall (see **Figure 4-22**). Additional grass and forb species occurred in sparse cover and included vine mesquite, Dakota verbena, and crimson sage. The tall and short shrub layer provided low to moderate cover (up to 15 percent cover) and included honey mesquite, littleleaf sumac, creosotebush, and four-wing saltbush. The shrub layer ranged from 1 to 4 m tall and contributed valuable wildlife habitat structure in the 6.9 acres examined in the survey corridor.



Figure 4-22. Representative Photographs of Alkali Sacaton Desert Wash Habitat

Wild Barley / Honey Mesquite Shrub Herbaceous Vegetation

A narrow drainage, up to 5 m wide, occurs [REDACTED] (b) (7)(E) and carries flows across the international border into Mexico (see **Figure 4-23**). The drainage, which occupies 0.3 acre within the survey corridor, is cobble-lined, appears to be spring-fed, and has a high groundwater table. The drainage was characterized by low cover of the annual wild barley, which provided up to 15 percent cover in addition to sparse cover provided by the grasses foxtail barley, rescue grass, and rabbitfoot grass. The forbs common sunflower and yellow sweetclover provided sparse cover. Along the drainage banks, honey mesquite shrubs to 4 m tall and netleaf hackberry trees to 7 m tall contributed low cover (up to 10 percent cover) and provided structural value as wildlife habitat.



Figure 4-23. Representative Photographs of Wild Barley/ Honey Mesquite Small Drainage Habitat

4.1.8 Apacherian-Chihuahuan Semi-Desert Grassland and Steppe Ecological System (CES302.735)

Hook Threeawn Herbaceous Vegetation

A moderately large stand of hook threeawn, providing approximately 15 percent cover, has become established on clay soils deposited in the vicinity of a livestock corral (see **Figure 4-24**). Sparse shrub cover (up to 2 percent cover) of creosotebush also occurs in this otherwise monotypic stand which occupies 1.8 acres within the survey corridor.



Figure 4-24. Representative Photographs of Hook Threeawn Flats Habitat

Desert Marigold Herbaceous Vegetation

An area that had been graded and cleared of creosotebush and mariola shrubs currently supported short-stature desert marigold forbs that provided low cover, (up to 8 percent cover) (see **Figure 4-25**). The disturbance covered 0.9 acre of the survey corridor and was maintained by ground squirrel and kangaroo rat burrowing and foraging activity, which was extensive across the site. The low-growing fluffgrass provided sparse grass cover, as did the dwarf-shrub little-head snakeweed.



Figure 4-25. Representative Photographs of Desert Marigold Flats Habitat

4.1.9 North American Warm Desert Cienega Ecological System (CES302.747)

Slimleaf Bursage – Common Sunflower Herbaceous Vegetation

A large old field, formerly an extensive cienega drained historically to support farming, occurs adjacent to (b) (7)(E) (see **Figure 4-26**). It holds water following large precipitation events and the old field/former cienega bottom is well-vegetated with forbs and grasses. The forbs slimleaf bursage, common sunflower, and prostrate saltbush provide moderate cover (up to 30 percent cover) within the 2.2 acres of the stand within the survey corridor. Common sunflower stalks from the previous year's growth attained heights to 3 m tall. Grasses provide low to moderate cover and include inland saltgrass, giant dropseed, and the nonnative Bermuda grass. The dwarf-shrub seep weed provides sparse cover. A (b) (7)(E) is illustrated in

Figure 4-26.

According to the USFWS (2008), the (b) (7)(E) once supported permanently flowing creeks, springs, and marshy wetlands composing this cienega. Giant sacaton grasslands occupied portions of the valley and were described as a luxuriant meadow some 8 or 10 miles wide. The dependable water sources and herbaceous vegetation cover made the area invaluable to fish and wildlife and to humans.



Figure 4-26. Representative Photographs of Slimleaf Bursage – Common Sunflower Cienega Habitat (b) (7)(E)

4.1.10 North American Warm Desert Playa Ecological System (CES302.751)

Russian-thistle Semi-natural Herbaceous Vegetation

A depression or playa that was nearly devoid of vegetation occurred (b) (7)(E) and formed across the border into Mexico. This depression apparently fills and ponds with water following large precipitation events. At the time of site visit (April 2008), sparse cover (less than 5 percent cover) of the annual forb Russian-thistle or tumbleweed and the perennial forb narrowleaf globemallow was emerging on the 2.8 acres examined in the survey corridor (see **Figure 4-27**). Dead stems from the previous year's growth indicated that low to moderate cover (up to 15 to 20 percent cover) by Russian-thistle could occur on this site. A single honey mesquite shrub provided sparse cover within this depression. The soils of this site are highly erosive with a texture of fine clay and silt and are apparently quite alkaline; water and wind erosion of these soils during construction could occur and could affect adjacent drainages with fine sediments. Equipment would have difficult access across this area when the soils are saturated and would damage the playa bottom with tire ruts.



Figure 4-27. Representative Photographs of Russian-thistle Playa Habitat

Common Cocklebur Semi-natural Herbaceous Vegetation

An (b) (7)(E) occurs on 0.3 acre across an access road and ponds water on both sides of the road when sufficient runoff is collected (see **Figure 4-28**). The annual forb common cocklebur had become established and provided moderate cover (35 percent) across the (b) (7)(E) bottom. Associated mesic grasses included Johnsongrass and scratchgrass that provided low cover (5 percent cover) along the (b) (7)(E) margin. The dwarf-shrub little-head snakeweed occurred on the (b) (7)(E) margin above the level of inundation and contributed sparse cover. This site could have difficult access when ponded water is present following precipitation events.



Figure 4-28. Representative Photographs of Common Cocklebur (b) (7)(E) Habitat

4.1.11 Other Nonnative Herbaceous Vegetation Alliances and Associations

Bermuda Grass Semi-natural Herbaceous Vegetation

A small irrigated pasture characterized by the nonnative Bermuda grass was maintained at the (b) (7)(E), located adjacent to (b) (7)(E) the international border (see **Figure 4-29**). The pasture occupied 0.3 acre in the survey corridor, provided 40 to 80 percent cover of Bermuda grass, and supported annual forbs along its edges and in bare patches. The annual forbs provided sparse cover and included Russian-thistle or tumbleweed, horsenettle, whitetop, and London rocket. A sprinkler irrigation system was installed that used rotating Rain-Bird style heads for water distribution.



**Figure 4-29. Representative Photographs of
Bermuda Grass Irrigated Pasture Habitat**

4.2 Plant Species Identified

A list of plant species prepared during the field surveys and annotated for nonnative and Arizona protected status is provided in **Table 4-2**. An early spring survey identified 125 taxa.

Table 4-2. Plant Species List, Relative Abundance in the Survey Corridor, and Habitat for (b) (7)(E) Station, FV-1b

Species / Common Name	Distribution	Location / Habitat
Trees and Tall Shrubs		
<i>Baccharis pteronoides</i> / Yerba de Pasmó	Rare	(b) (7)(E)
<i>Baccharis sarothroides</i> / Desert broom	Uncommon	Desert washes, roadsides
<i>Celtis reticulata</i> / Nettleleaf hackberry	Rare	Desert washes, springs
<i>Chilopsis linearis</i> / Desert willow ⁴	Rare	(b) (7)(E)
<i>Fouquieria splendens</i> / Ocotillo ³	Common	Rocky slopes, alluvial plains
<i>Fraxinus velutina</i> / Green ash	Rare	(b) (7)(E)
<i>Juniperus monosperma</i> / One-seed juniper	Rare	Limestone outcrops, (b) (7)(E)
<i>Populus fremontii</i> / Fremont cottonwood	Rare	(b) (7)(E), springs
<i>Prosopis glandulosa</i> / Honey mesquite ^{4,5}	Abundant	Rocky slopes, alluvial plains, desert washes, swales, cienegas, playas
<i>Salix gooddingii</i> / Goodding willow	Rare	(b) (7)(E)
Short and Dwarf Shrubs		
<i>Acacia constricta</i> / Whitethorn	Abundant	Most upland habitats
<i>Acacia millefolia</i> / Acacia	Uncommon	Rocky slopes
<i>Agave palmeri</i> / Century plant ³	Uncommon	Rocky slopes
<i>Agave parryi</i> / Parry agave ³	Uncommon	Rocky slopes
<i>Agave</i> sp. / Agave ³	Rare	Limestone bedrock
<i>Anisacanthus thurberi</i> = <i>Justicia californica</i> / Chuparosa	Uncommon	Desert washes
<i>Atriplex canescens</i> / Fourwing saltbush	Common	Rocky slopes, alluvial fans, desert washes
<i>Bebbia juncea</i> / Rush bebbia	Rare	Desert washes
<i>Brickellia californica</i> / Brickelbush, Pachaba	Rare	Rocky slopes, Desert washes
<i>Bumelia lanuginosa</i> / Buckthorn	Rare	Rocky slopes
<i>Clematis drummondii</i> / Texas virgin's bower	Rare	Desert wash
<i>Condalia spathulata</i> / Squawbush	Uncommon	Rocky slopes, alluvial fans
<i>Dalea formosa</i> / Feather peabush	Uncommon	Rocky slopes
<i>Dasyilirion wheeleri</i> / Sotol ³	Uncommon	Rocky slopes, limestone bedrock

Species / Common Name	Distribution	Location / Habitat
Short and Dwarf Shrubs (continued)		
<i>Dyssodia acerosa</i> / Spiny dogweed, Prickly fetid marigold	Uncommon	Rocky slopes
<i>Echinocereus engelmannii</i> / Strawberry hedgehog ³	Rare	Rocky slopes
<i>Echinocereus pectinatus</i> / Rainbow cactus ³	Rare	Rocky slopes, alluvial fans
<i>Encelia farinosa</i> / Brittlebush	Rare	Rocky slopes
<i>Ephedra trifurca</i> / Long-leaved jointfir, Mormon-tea	Rare	Rocky slopes
<i>Flourensia cernua</i> / Tarbush	Abundant	Rocky slopes, alluvial fans, plains, desert washes
<i>Gutierrezia microcephala</i> / Little-head snakeweed	Common	Rocky slopes, alluvial fans, plains, desert washes
<i>Hymenoclea monogyra</i> / Burro bush	Uncommon	Creek bed
<i>Hymenoclea salsola</i> / Cheesebush	Uncommon	Desert washes
<i>Koeberlinia spinosa</i> / Junco, Allthorn	Rare	Limestone outcrops
<i>Larrea tridentata</i> / Creosotebush	Abundant	Most upland habitats
<i>Mammillaria</i> sp. / Fishhook cactus ³	Rare	Rocky slopes, alluvial fans
<i>Menodora scabra</i> / Rough menodora	Uncommon	Limestone outcrops
<i>Mimosa biuncifera</i> / Wait-a-minute	Uncommon	Rocky slopes
<i>Mortonia scabrella</i> / Mortonia	Uncommon	Limestone outcrops and colluvium
<i>Opuntia leptocaulis</i> / Desert Christmas cactus ³	Uncommon	Rocky slopes
<i>Opuntia phaeacantha</i> / Engelmann prickly pear ³	Common	Rocky slopes, alluvial fans
<i>Opuntia ramosissima</i> / Diamond cholla ³	Uncommon	Rocky slopes
<i>Opuntia spinosior</i> / Cane cholla ³	Uncommon	Rocky slopes
<i>Opuntia violacea</i> / Purple prickly pear ³	Uncommon	Rocky slopes
<i>Parthenium incanum</i> / Mariola	Abundant	Most upland habitats
<i>Peniocereus greggii</i> / Deerhorn cactus ³	Rare	Volcanic cobble slope
<i>Platanus wrightii</i> / Arizona sycamore	Rare	(b) (7)(E) bottom
<i>Quercus</i> sp. / Oak	Rare	Limestone outcrop
<i>Rhus microphylla</i> / Littleleaf sumac	Uncommon	Desert washes, swales, creeks, draws
<i>Suaeda torreyana</i> / Seepweed	Rare	Cienega

Species / Common Name	Distribution	Location / Habitat
Short and Dwarf Shrubs (continued)		
<i>Tiquilia canescens</i> = <i>Coldenia canescens</i> / Dog's ear	Abundant	Most upland habitats
<i>Yucca elata</i> / Soap tree yucca ³	Uncommon	Sandy plains
<i>Yucca schottii</i> / Yucca ³	Uncommon	Rocky slopes, alluvial fans
<i>Ziziphus obtusifolia</i> = <i>Condalia lycioides</i> / Graythorn	Uncommon	Rocky slopes
Graminoids		
<i>Aristida adscensionis</i> / Annual threeawn	Uncommon	Alluvial fans, plains, disturbed sites
<i>Aristida purpurea</i> / Purple threeawn	Uncommon	Rocky slopes
<i>Aristida hamulosa</i> / Hook threeawn	Common	Rocky slopes, alluvial fans, plains
<i>Bouteloua eriopoda</i> / Black grama	Uncommon	Rocky slopes
<i>Bromus willdenowii</i> = <i>Bromus catharticus</i> / Rescue grass	Rare	Spring-fed drainage
<i>Carex</i> sp. / Sedge	Rare	Spring-fed drainage
<i>Chloris virgata</i> / Windmill grass ¹	Rare	Spring-fed drainage
<i>Cynodon dactylon</i> / Bermudagrass ¹	Rare	Irrigated pasture
<i>Distichlis spicata</i> / Saltgrass	Rare	Cienega
<i>Erioneuron pulchellum</i> = <i>Tridens pulchellus</i> / Fluffgrass	Abundant	Most upland habitats
<i>Hilaria mutica</i> / Tobosa	Common	Most upland habitats
<i>Hordeum jubatum</i> / Foxtail barley ¹	Rare	Spring-fed drainage
<i>Hordeum leporinum</i> / Wild barley ¹	Rare	Spring-fed drainage
<i>Juncus balticus</i> / Baltic rush	Rare	Spring-fed drainage
<i>Leptochloa dubia</i> / Sprangletop	Uncommon	Rocky slopes
<i>Muhlenbergia asperifolia</i> / Scratchgrass	Rare	Excavated site
<i>Muhlenbergia porteri</i> / Bush muhly	Uncommon	Rocky slopes
<i>Polypogon monspeliensis</i> / Rabbitsfoot grass ¹	Rare	Spring-fed drainage
<i>Schismus barbatus</i> / Mediterranean grass ¹	Uncommon	Alluvial fans, plains
<i>Scirpus americanus</i> = <i>Schoenoplectus pungens</i> / Three-square bulrush	Rare	(b) (7)(E) spring-fed sites, ponds

Species / Common Name	Distribution	Location / Habitat
Graminoids (continued)		
<i>Schizachyrium scoparium</i> / Little bluestem	Rare	rock outcrops
<i>Sorghum halepense</i> / Johnsongrass ¹	Rare	Spring-fed drainage
<i>Sporobolus airoides</i> / Alkali sacaton	Common	Desert washes, swales
<i>Sporobolus cryptandrus</i> / Sand dropseed	Uncommon	Desert washes, sandy areas
<i>Sporobolus giganteus</i> / Giant dropseed	Common	Desert washes
<i>Typha domingensis</i> / Southern cattail	Rare	(b) (7)(E) springs, ponds
Forbs		
<i>Acourtia nana</i> = <i>Perezia nana</i> / Dwarf desert holly	Common	Most upland habitats
<i>Allionia incarnata</i> / Trailing windmills	Uncommon	Rocky slopes, desert washes
<i>Amaranthus palmeri</i> / Palmer's amaranth ¹	Rare	Cienega
<i>Ambrosia confertiflora</i> = <i>Franseria confertiflora</i> / Slimleaf bursage	Rare	Cienega
<i>Argemone</i> sp. / Prickly poppy	Rare	(b) (7)(E)
<i>Astragalus</i> sp. / Milkvetch	Rare	Desert wash
<i>Atriplex</i> sp. / Prostrate saltbush	Rare	Cienega, playa, irrigated pasture
<i>Baileyia multiradiata</i> / Desert marigold	Uncommon	Alluvial fans, plains, bladed site
<i>Cardaria</i> sp. / Whitetop ¹	Uncommon	Irrigated pasture
<i>Centaurea melitensis</i> / Malta starthistle	Rare	Spring-fed drainage
<i>Chenopodium fremontii</i> / Fremont goosefoot	Rare	Cienega, Irrigated pasture
<i>Cryptantha</i> sp. / Cryptantha	Common	Most upland habitats
<i>Cucurbita foetidissima</i> / Buffalo gourd	Rare	Desert wash
<i>Cymopterus multinervatus</i> / Purple cymopterus	Rare	Rocky slopes
<i>Datura meteloides</i> / Sacred datura	Rare	Desert washes
<i>Descurainia pinnata</i> / Tansy mustard ¹	Rare	Irrigated pasture, disturbed roadside
<i>Eriastrum diffusum</i> / Miniature wool-star	Uncommon	Rocky slopes, alluvial fans
<i>Erodium cicutarium</i> / Filaree ¹	Rare	Irrigated pasture

Species / Common Name	Distribution	Location / Habitat
Forbs (continued)		
<i>Euphorbia albomarginata</i> / Rattlesnake weed	Rare	Spring-fed drainage
<i>Haplopappus spinosus</i> / Golden aster	Uncommon	Rocky slopes
<i>Helianthus annuus</i> / Common sunflower	Uncommon	Cienega, playa, spring-fed drainage
<i>Lappula redowskii</i> / Stickweed	Uncommon	Rocky slopes, cienega, disturbed roadsides
<i>Lepidium thurberi</i> / Thurber's peppergrass	Rare	Plains
<i>Medicago lupulina</i> / Black medic ¹	Rare	Spring-fed drainage
<i>Melilotus officianalis</i> / Yellow sweetclover ¹	Rare	Spring-fed drainage
<i>Mentzelia albicaulis</i> / White stem stickleaf	Uncommon	Rocky slopes, desert washes
<i>Mentzelia pumila</i> / Stickleaf	Uncommon	Rocky slopes
<i>Notholaena</i> sp. / Cloak fern	Rare	Limestone bedrock
<i>Pectis fillipes</i> / Threadstem cinchweed	Rare	Rocky slopes
<i>Penstemon superbus</i> / Superb penstemon	Rare	Desert washes
<i>Penstemon</i> sp. / Beardtongue	Rare	Limestone bedrock
<i>Phacelia coerulea</i> / Blue scorpionweed	Uncommon	Desert washes
<i>Phoradendron californicum</i> / Mistletoe	Uncommon	Honey mesquite trees and shrubs, many habitats
<i>Phoradendron flavescens</i> / Mistletoe	Rare	Arizona sycamore trees, desert washes
<i>Plantago patagonica</i> / Plantain	Common	Rocky slopes, alluvial fans, plains
<i>Proboscidea parviflora</i> / Devil's claw	Rare	Alluvial plains, desert washes
<i>Rumex hymenosepalus</i> / Canaigre	Rare	Cienega
<i>Salsola iberica</i> / Russian-thistle ¹	Rare	Playa, disturbed roadsides
<i>Salvia henryi</i> / Crimson sage	Uncommon	Desert washes
<i>Selaginella</i> sp. / Club moss	Rare	Limestone bedrock
<i>Silene antirrhinum</i> / Sleepy catchfly ¹	Uncommon	Most upland habitats
<i>Sisymbrium irio</i> / London rocket ¹	Rare	Irrigated pasture, disturbed roadsides
Forbs (continued)		

Species / Common Name	Distribution	Location / Habitat
<i>Solanum eleagnifolium</i> / Horse nettle ¹	Rare	Desert washes, irrigated pasture, cienega
<i>Sphaeralcea angustifolia</i> / Narrowleaf globemallow	Uncommon	Rocky slopes
<i>Sphaeralcea laxa</i> / Caliche globemallow	Rare	Playa
<i>Verbena gooddingii</i> / Goodding (Dakota) verbena	Uncommon	Desert washes
<i>Verbesina encelioides</i> / Cowpen daisy	Rare	Irrigated pasture
<i>Xanthium strumarium</i> / Common cocklebur ¹	Rare	Excavated depression

Notes:

¹ Nonnative species (noxious weeds were not identified within the corridor).

² Highly Safeguarded Protected Native Plants (this category was not identified within the corridor): species of native plants whose prospects for survival in Arizona are in jeopardy or are in danger of extinction).

³ Salvage Restricted Protected Native Plants (species of native plants that are subject to damage by theft or vandalism).

⁴ Salvage Assessed Protected Native Plants (species of native plants that have a sufficient value if salvaged to support the cost of salvage).

⁵ Harvest Restricted Protected Native Plants (species of native plants that are subject to excessive harvesting or overcutting because of their intrinsic value).

4.3 Survey Corridor Characteristics and Description of Habitat Quality

To ensure the most recent data were acquired for rare species analyses, e²M requested Element Occurrence Data from NatureServe Central Databases in Arlington, Virginia, through a referral from the USFWS (NatureServe and e²M 2007a). Additionally, rare species data were acquired from AZGFD and USFWS at Project inception. General descriptions of the habitat quality as it relates to rare plant species and the landscape characteristics of the survey corridor are provided herein and are based on field observations, personal communications, and the literature.

SECTION FV-1b

County: (b) (7)(E)

Potential Listed

Plant Occurrence: *Spiranthes delitescens* (Canelo Hills or Madrean ladies'-tresses) (Federal [FE], state [HS])
Coryphantha robbinsorum (Cochise pincushion cactus) (Federal [FE], state [HS])
Lilaeopsis schaffneriana ssp. *recurva* (Huachuca water umbel) (Federal [FE], state [HS])

Erigeron lemmonii (Lemmon fleabane) (Federal [FC], state [HS])

Listed Plants Observed: None

Suitable Listed Plant Habitat Present: Possible habitat (perennial low gradient streams and wetlands) for the Huachuca water umbel occurs adjacent to (b) (7)(E).

If so, Habitat Quality: Fair to Good

Section Habitat Description: This section includes approximately (b) (7)(E)

(b) (7)(E)

(b) (7)(E) The western portion of section FV-1b occurs on steep slopes dominated by ocotillo, transitioning to alluvial fans and plains characterized by creosotebush, mariola, tarbush, and honey mesquite shrubs. A few slopes are armored by volcanic gravel and cobble where patches of tobosa occur amid shrublands dominated by honey mesquite and/or creosotebush. Gullies and desert washes commonly occur in this terrain. Within (b) (7)(E) lowland habitats including irrigated pasture, go-back fields, and extensive stands of honey mesquite with four-wing saltbush in the understory, and riparian forests and woodlands occur. Particularly good riparian forest habitats occupy (b) (7)(E) and thick honey mesquite woodlands occur along (b) (7)(E). The easternmost portion of the FV-1b survey corridor consists of ridges and drainages supporting creosotebush and mariola shrublands, honey-mesquite-dominated lower slopes and drainage bottoms, and stands of alkali sacaton in narrow drainages. The road providing access between (b) (7)(E) and (b) (7)(E) crosses exposed Permian limestone, and the shrubland is dominated by mortonia, tarbush, and mariola. This access road traverses the only small trees of one-seed juniper and shrubs of oak and junco observed in the survey corridor.

- Madrean ladies'-tresses occur in the (b) (7)(E) in cienegas with finely grained, highly organic, saturated soils. This habitat does not occur within the survey corridor.

Cochise pincushion cactus occurs on gray (Permian) limestone hills that support semi-desert grassland with small shrubs, agave, cacti, and grama grass. It does not co-occur with mortonia shrubs on lower slopes (USFWS 1993). Permian limestone outcrops exposed along an access road on the eastern end of FV-1b survey corridor were searched for Cochise pincushion cactus, but none were observed and mortonia was a dominant shrub. There are no limestone outcrops within the Project area. The hill or small mountain (b) (7)(E) (b) (7)(E) does have appropriate Permian limestone habitat and does support a semi-desert grassland community on its upper one-fourth; however, there will be no construction and hence no impact to this potential habitat for the Cochise pincushion cactus.

Huachuca water umbel occurs in cienegas, perennial low gradient streams, and wetlands. (b) (7)(E) and one nearby spring (b) (7)(E) provide this habitat and represent potential sites for Huachuca water umbel establishment.

Lemmon fleabane occurs in pine-oak woodlands in rock crevices, on ledges, and among boulders in canyon bottoms. This habitat does not occur within the survey corridor.

4.4 Wetlands and Waters of the United States

Wetlands and waters of the United States can be confusing terms and are defined here for the convenience of document users. The USACE has jurisdiction to protect wetlands under section 404 of the Clean Water Act using the following definition:

... areas that are inundated or saturated by surface or ground water at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions (33 Code of Federal Regulations [CFR] 328.3[b]). Wetlands generally include swamps, marshes, bogs, and similar areas.

Wetlands have three diagnostic characteristics that include (1) more than 50 percent of the dominant species present must be classified as obligate, facultative wetland; or facultative, (2) the soils must be classified as hydric; and (3) the area is either permanently or seasonally inundated (Environmental Laboratory 1987).

Waters of the United States are defined under 33 *United States Code* (U.S.C.) 1344, as follows:

- a. The term “waters of the United States” means
 1. All waters which are currently used, or were used in the past, or may be susceptible to use in interstate or foreign commerce, including all waters which are subject to the ebb and flow of the tide;
 2. All interstate waters including interstate wetlands;
 3. All other waters such as intrastate lakes, rivers, streams (including intermittent streams), mudflats, sandflats, wetlands, sloughs, prairie potholes, wet meadows, playa lakes, or natural ponds, the use, degradation or destruction of which could affect interstate or foreign commerce including any such waters:
 - i. Which are or could be used by interstate or foreign travelers for recreational or other purposes; or
 - ii. From which fish or shellfish are or could be taken and sold in interstate or foreign commerce; or

- iii. Which are used or could be used for industrial purpose by industries in interstate commerce;
- 4. All impoundments of waters otherwise defined as waters of the United States under the definition;
- 5. Tributaries of waters identified in paragraphs (a) (1)-(4) of this section;
- 6. The territorial seas;
- 7. Wetlands adjacent to waters (other than waters that are themselves wetlands) identified in paragraphs (a) (1)-(6) of this section.
- 8. Waters of the United States do not include prior converted cropland. Notwithstanding the determination of an area's status as prior converted cropland by any other federal agency, for the purposes of the Clean Water Act (CWA), the final authority regarding CWA jurisdiction remains with the EPA.
- 9. Waste treatment systems, including treatment ponds or lagoons designed to meet the requirements of the CWA (other than cooling ponds as defined in 40 CFR 123.11(m) which also meet the criteria of this definition) are not waters of the United States.
- b. The term "wetlands" means those areas that are inundated or saturated by surface or ground water at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions. Wetlands generally include swamps, marshes, bogs, and similar areas.
- c. The term "adjacent" means bordering, contiguous, or neighboring. Wetlands separated from other waters of the United States by man-made dikes or barriers, natural river berms, beach dunes and the like are "adjacent wetlands."
- d. The term "high tide line" means the line of intersection of the land with the water's surface at the maximum height reached by a rising tide. The high tide line may be determined, in the absence of actual data, by a line of oil or scum along shore objects, a more or less continuous deposit of fine shell or debris on the foreshore or berm, other physical markings or characteristics, vegetation lines, tidal gages, or other suitable means that delineate the general height reached by a rising tide. The line encompasses spring high tides and other high tides that occur with periodic frequency but does not include storm surges in which there is a departure from the normal or predicted reach of the tide due to the piling up of water against a coast by strong winds such as those accompanying a hurricane or other intense storm.
- e. The term "ordinary high water mark" means that line on the shore established by the fluctuations of water and indicated by physical characteristics such as clear, natural line impressed on the bank, shelving, changes in the character of soil, destruction of terrestrial vegetation, the

presence of litter and debris, or other appropriate means that consider the characteristics of the surrounding areas.

The term "tidal waters" means those waters that rise and fall in a predictable and measurable rhythm or cycle due to the gravitational pulls of the moon and sun. Tidal waters end where the rise and fall of the water surface can no longer be practically measured in a predictable rhythm due to masking by hydrologic, wind, or other effects.

The survey corridor lies within (b) (7)(E), which includes approximately 250,000 acres (USEPA 2008; NRCS 2008). The watershed occurs in the (b) (7)(E). The (b) (7)(E) and artesian aquifer cover approximately (b) (7)(E).

Generally, the (b) (7)(E) an asymmetric extensional basin that experienced extensive mafic volcanism in the Cenozoic Period (Earman et al. 2002). On a large scale, it lies within the Rio Yaqui Watershed (USFWS 1994a). In Arizona, the basin's major fault lies on the western side of the basin, and in Sonora, the Pitaycachi Fault occurs on the eastern side of the basin (Earman et al. 2002). In Arizona, the bedrock forms a half graben down-dropped on the western side by antithetic normal faults with structural highs formed by transfer faults. The heterogeneous basin fill, which contains numerous basalt interbeds, was deposited predominantly in alluvial fan and alluvial slope environments. Generally, the basin structure does not have a large effect on groundwater flow as most water is transmitted through paleochannel deposits that make up a relatively small portion of the aquifer.

Arizona water rights (b) (7)(E) date from 1882 for unlitigated surface water rights and from 1903 for groundwater permits (USFWS 1995). Surface water rights include (1) (b) (7)(E) (100 acre-feet per year); (2) (b) (7)(E) (19 gallons per minute); (3) (b) (7)(E) (15 gallons per minute); (4) (b) (7)(E) (9.5 gallons per minute); (5) (b) (7)(E) (49 gallons per minute); and (6) (b) (7)(E) (3.8 gallons per minute). There are ten permitted groundwater wells on the refuge that each are adjudicated for 250 acre-feet of water production annually.

4.4.1 Field Evaluation Summary

Observations and initial identification of potential wetlands and waters of the United States for the survey corridor were recorded during the April/May 2008 field inventory.

Field surveys were conducted on June 10 through 13, 2008, to delineate jurisdictional wetlands and other Waters of the United States within the survey corridor. Delineations were also conducted along access roads and staging

areas associated with the fence alignments. Formal delineations were conducted within a 150-foot corridor associated with the fence alignments, 60 feet to either side of the center line of access roads, and within staging areas.

Determination of the occurrence and extent of jurisdictional wetlands and other Waters of the United States were based on the application of procedures established in the USACE *Wetlands Delineation Manual*, Technical Report Y-87-1 (USACE 1987) and the *Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Arid West Region*, Technical Report ERDC/EL TR-06-16 (USACE 2006). Determination of the occurrence of jurisdictional wetlands was based on the presence or absence of hydrophytic (wetland) vegetation, hydric (wetland) soils, and wetland hydrology. The presence of all three of the criteria is necessary for an area to be designated as a jurisdictional wetland under normal conditions.

Determination of the extent of jurisdictional washes and other Waters of the United States in the survey corridor was based on characterization of the landward extent of the ordinary high water mark (OHM). Indicators used to determine the occurrence and extent of jurisdictional washes included the presence of developed channels, typically 2 feet or greater in width; the occurrence of an OHM; the absence of fine sediments along flow paths; distinct changes in the vegetative assemblage or larger or more dense vegetation than surrounding areas; the presence of cut banks; the presence of litter, debris, or wrack lines; occurrence of desiccation cracks or other indicators of hydrology; and other indicators of the occurrence of intermittent water flow regimes.

All wetlands and other Waters of the United States within the survey corridor were delineated.

Table 4-3 provides wetland and other Waters of the United States types and delineated acreages within a 150-foot corridor associated with the fence alignments, 60 feet to either side of the center line of access roads, or within planned staging areas; and potential impact acreages in Section FV-1b. A 60-foot impact corridor to the north of the fence alignment or adjacent to access roads is considered the maximum width of potential impact associated with implementing the Project. All wetland or other Waters of the United States acreages within staging areas are included as potential impact areas. The following text provides general descriptions of wetlands and other Waters of the United States identified within the Project assessment areas in Section FV-1b.

Table 4-3. Delineated Acreages, General Locations and Potential Impact Acreage of Wetlands and Other Waters of the United States in (b) (7)(E) AZ Section FV-1b

WL ID	Wetland or Other WOUS Type	General Location	Delineated Area (acres)	Potential Impacts (acres)
(b) (7)(E)	Wash	(b) (7)(E)	5.14	1.75
	Wash		0.27	0.07
	Wash		0.18	0.07
	Wash		0.28	0.09
	Wash		0.26	0.07
	Wash		0.98	0.31
	Wash		0.43	0.20
	Wash		4.88	1.66
	Wash		0.17	0.08
	Wash		0.29	0.17
	Wash		0.46	0.15
	Wash		0.25	0.01
	Wash		6.07	0.42
	Wash		0.55	0.29
	Emergent wetland		0.11	0.01
(b) (7)(E)	Riverine and palustrine forested wetland		1.74	0.35
	Wash		1.92	1.31

WL ID	Wetland or Other WOUS Type	General Location	Delineated Area (acres)	Potential Impacts (acres)
(b) (7)(E)	Wash	(b) (7)(E)	1.33	0.36
	Wash		1.07	0.25
	Wash		0.27	0.11
	Wash		0.44	0.15
	Wash		0.61	0.18
	Wash		0.30	0.15
	Wash		0.21	0.12
	Wash		0.46	0.09
	Wash		0.49	0.12
	Wash		3.68	2.54
	Wash		0.26	0.10
	Wash		0.45	0.22
	Wash		0.38	0.05
	Wash		0.32	0.16
	Wash		0.17	0.03
	Wash		0.19	0.08
	Wash		0.14	0.07
	Wash		0.50	0.10

WL ID	Wetland or Other WOUS Type	General Location	Delineated Area (acres)	Potential Impacts (acres)
(b) (7)(E)	Wash	(b) (7)(E)	0.55	0.08
	Wash		0.69	0.34
	Wash		1.24	0.84
	Wash		0.13	0.13
	Wash		0.15	0.04
Totals			38.01	13.32

Based on the field surveys, 37 ephemeral wash channels and 2 vegetated wetlands occur within the (b) (7)(E) corridor associated with the fence alignments, (b) (7)(E) to either side of the center line of access roads, or within staging areas. Of the 37.87 total delineated acres of wetlands and other Waters of the United States, 13.28 acres occur within the potential impact areas. Wetlands and other Waters of the United States delineated in Section FV-1b were designated as (b) (7)(E).

(b) (7)(E) is characterized by two ephemeral wash channels ((b) (7)(E)) that flow (b) (7)(E) along the U.S./Mexico international border. (b) (7)(E) is located primarily (b) (7)(E).

Channel width at base in the two channels ranges from 2 feet to 5 feet with near-vertical banks. Vegetation characterizing the banks of the two channels is characterized by creosotebush (*Larrea tridentate*), honey mesquite (*Prosopis glandulosa*) and littleleaf sumac (*Rhus microphylla*).

(b) (7)(E) are ephemeral washes that cross the (b) (7)(E). The access road is located approximately (b) (7)(E). (b) (7)(E) is the downstream component of (b) (7)(E). (b) (7)(E) join (b) (7)(E) and then join with (b) (7)(E). (b) (7)(E) have wide shallow channels (b) (7)(E) that range from 12 to 15 feet in width at their base. The banks range from approximately 1 to 2 feet in height with bank slopes ranging from 45 percent to near vertical. (b) (7)(E) has a narrow channel that ranges from 2 to 4 feet in width at base with approximately 2-foot-high banks and slopes ranging from 45 percent to near vertical. Vegetation characterizing the banks of the three wash channels includes littleleaf sumac, honey mesquite, desert broom (*Baccharis sarothroides*), whitethorn acacia (*Acacia constricta*), tar bush (*Flourensia cernua*), little-head snakeweed (*Gutierrezia microcephala*), and Engelmann prickly-pear (*Opuntia engelmannii*).

(b) (7)(E) are ephemeral washes that cross (b) (7)(E). (b) (7)(E) is a narrow wash with an approximately 2-foot-wide channel at base and near-vertical 2-foot-high banks (b) (7)(E). The channel directs flows to the southeast. Vegetation on the wash banks is characterized by honey mesquite and Engelmann prickly-pear. (b) (7)(E) is a wide, incised ephemeral wash with an approximately 25- to 35-foot-wide channel at base. (b) (7)(E) the channel is characterized by near vertical 6 to 8 foot high banks on its south side and 2 foot high banks with 45 percent slopes on the north side. The channel flows (b) (7)(E). (b) (7)(E) Vegetation occurring on the wash banks in proximity to the access road and fence alignment is characterized by honey mesquite.

(b) (7)(E) is a narrow ephemeral wash that (b) (7)(E). The wash is characterized by a 4-foot-wide channel at base and near-vertical 3-foot banks. Vegetation of the banks of the wash is characterized by honey mesquite.

(b) (7)(E) is an ephemeral wash that drains (b) (7)(E). The wash crosses (b) (7)(E). Channel width at base ranges from approximately 15 feet at the confluence with (b) (7)(E) to approximately 3 feet to 8 feet at the (b) (7)(E). (b) (7)(E). Channel banks are near vertical and range from approximately 3 feet to 8 feet in height at Project alignment crossing. Vegetation on the banks of (b) (7)(E) is characterized by honey mesquite, littleleaf sumac, desert broom, and creosotebush.

(b) (7)(E) are tributary washes that drain into (b) (7)(E). (b) (7)(E) is a narrow ephemeral wash that directs flows (b) (7)(E) discharging into (b) (7)(E) direct flows (b) (7)(E) where they discharge into (b) (7)(E) directs flows (b) (7)(E) discharges into (b) (7)(E). (b) (7)(E) Channel widths at base range from 3 feet to 5 feet in the survey corridor and the channel banks range from 1 to 3 feet in height with bank slopes ranging from 45 percent to near vertical. Vegetation characterizing the wash banks is characterized by white thorn acacia, littleleaf sumac, and creosotebush.

(b) (7)(E) is (b) (7)(E) a large, wide, ephemeral wash with high, near-vertical banks on outside bends and well-developed alluvial bars on inside bends. The wash directs flows (b) (7)(E). The channel width at base ranges from approximately 400 feet (b) (7)(E) to approximately 160 feet (b) (7)(E). Bank

heights on outside bends exceed 20 feet in places. Honey mesquite and desert broom characterize the vegetation of the wash banks and alluvial bar deposits.

(b) (7)(E) is an ephemeral wash that is the downstream segment of a wash that crosses through (b) (7)(E). Most of the historical natural flow in the wash has probably been cut off as a result of an upstream impoundment constructed (b) (7)(E). The wash within the Project alignment has been disturbed and is characterized an approximately 30-foot-wide channel at base, with 2-foot-high, non-distinct banks having 25 percent slopes. Vegetation on the banks of the wash is characterized by honey mesquite. Several species of grasses occur in the wash channel.

(b) (7)(E) is a palustrine emergent wetland located along the Project alignment in the (b) (7)(E). The small emergent wetland is characterized by a near monotypic stand of three square (*Scirpus olneyi*). A few Goodding's willows (*Salix nigra* var. *goodingii*) and Fremont cottonwoods (*Populus fremontii*) occur on the perimeter of the wetland. A vehicle fence has been placed within the boundaries of the wetland by the USFWS.

(b) (7)(E) is a riverine and palustrine forested wetland associated with (b) (7)(E). (b) (7)(E) a perennial stream that directs flows (b) (7)(E). (b) (7)(E) has a deeply incised central channel that is approximately 60 feet wide at base. The central channel was maintaining flow at the time of the site survey. Base flow in the channel is supported by upstream springs. The riverine component of (b) (7)(E) occurs in association with the central channel. Vegetation occurring in association with the riverine wetland is characterized by southern cattail (*Typha domingensis*) and three square. The channel is bordered by a low terrace. The width of the central channel and terrace is approximately 100 feet with near-vertical 10-foot banks bordering the terrace. The palustrine forested component of (b) (7)(E) occurs in association with the terrace. Vegetation occurring in association with the palustrine forested wetland includes Fremont cottonwood and Goodding's willows with some honey mesquite. Vegetation on the upper banks of (b) (7)(E) is characterized by honey mesquite. An impoundment has been constructed across the channel and terrace with gabions. (b) (7)(E)

(b) (7)(E) is (b) (7)(E) in proximity to the Project alignment is characterized by a vegetated ephemeral wash with an approximately 75-foot channel width at base. Channel banks are approximately 8 feet in height and vertical. Fremont cottonwood and Goodding's willow occur in association with alluvial bars in the wash and some giant dropseed (*Sporobolus giganteus*) occurs along the base of the channel banks. Vegetation on the wash banks is characterized by honey mesquite. The wash channel and banks have been disturbed at the border as a result of border crossings.

(b) (7)(E) is a deeply incised ephemeral wash that directs flows (b) (7)(E). The channel width at base is approximately 7 feet. Channel banks are

near vertical and are approximately 8 feet in height. Vegetation on the banks of the wash is characterized by white thorn acacia and creosotebush.

(b) (7)(E) is a disturbed ephemeral wash that historically directed flows (b) (7)(E). Flow to the wash has been cut off by an impoundment located approximately 300 feet upstream from the border. (b) (7)(E), the channel base is approximately 20 feet wide. Channel banks have been disturbed and are approximately 3 feet high with 35 percent slopes. Giant dropseed and little-head snakeweed occur in the wash channel and vegetation on the banks is characterized by Russian thistle (*Salsola kali*), four-wing saltbush (*Atriplex canescens*), and honey mesquite.

(b) (7)(E) are ephemeral washes that direct flows (b) (7)(E). The two washes join approximately (b) (7)(E). (b) (7)(E) discharges into the main wash channel (b) (7)(E). (b) (7)(E) has an incised channel that is approximately 5 feet wide at base with 5-foot-high banks in the survey corridor. The channel banks slope at approximately 60 percent. (b) (7)(E) has an approximately 20-foot-wide channel at base with 3-foot banks in the survey corridor. The channel banks have 35 percent slopes. Vegetation on the banks of both washes is characterized by honey mesquite, creosotebush, four-wing saltbush, and tar bush.

(b) (7)(E) is an ephemeral wash that directs flows (b) (7)(E). The channel width at base is approximately 4 feet. Channel banks are approximately 4 feet high with 45 percent slopes at the border and become near vertical downstream (b) (7)(E). Evidence of active headwall cutting is present in places along the channel. Vegetation on the banks of the wash is characterized by honey mesquite and white thorn acacia.

(b) (7)(E) is an ephemeral wash that directs flows (b) (7)(E)s. The channel width at base is approximately 8 feet. Channel banks are near vertical and approximately 5 feet in height. Vegetation on the banks of the wash is characterized by honey mesquite, four-wing saltbush, white thorn acacia, and little-head snakeweed.

(b) (7)(E) is an ephemeral wash that directs flows (b) (7)(E). The channel width at base is approximately 2 feet. Channel banks are approximately 4 feet high with 55 percent slopes. The wash in the survey corridor is characterized by two channels with an alluvial island in the center. The channels join (b) (7)(E). Vegetation on the banks of the channels is characterized by honey mesquite, four-wing saltbush, and little-head snakeweed.

(b) (7)(E) is an ephemeral wash that directs flows (b) (7)(E). The channel width at base is approximately 4 feet. Channel banks are approximately 8 feet high with 45 percent slopes. Vegetation on the banks of the wash is characterized by honey mesquite and little-head snakeweed.

(b) (7)(E) is an ephemeral wash that directs flows (b) (7)(E). The channel widths at base range from approximately 3 to 8 feet. Channel bank heights range from 3 feet to 8 feet with bank slopes ranging from approximately 75 percent to near vertical. Vegetation on the banks of the wash is characterized by honey mesquite, giant dropseed, four-wing saltbush, and little-head snakeweed.

(b) (7)(E) is a narrow ephemeral wash that (b) (7)(E). The channel width at base is approximately 3 feet. Channel banks are approximately 2 feet high with 45 percent slopes. Vegetation on the banks of the wash is characterized by honey mesquite, littleleaf sumac, and creosotebush.

(b) (7)(E) is an ephemeral wash that (b) (7)(E). The channel width at base in the staging area is approximately 5 feet. Channel banks are approximately 2 feet high with 45 percent slopes. The channel becomes indistinct and braided where it crosses the border. Vegetation on the banks of the wash is characterized by white thorn acacia, tar bush, creosotebush, giant dropseed, and littleleaf sumac.

(b) (7)(E) is an ephemeral wash that (b) (7)(E). The channel width at base is approximately 3 feet. Channel banks are approximately 5 feet high with 55 percent slopes. Vegetation on the banks of the wash is characterized by tar bush, littleleaf sumac, white thorn acacia, purple three awn (*Aristida purpurea*), and little-head snakeweed.

(b) (7)(E) is an ephemeral wash that directs flows (b) (7)(E). The channel base is approximately 2 feet wide (b) (7)(E). The channel banks are near vertical and approximately 2 feet high. Approximately 60 feet downstream of the border the channel width at base is approximately 4 feet with 1-foot near-vertical banks. Vegetation on the banks of the wash is characterized by honey mesquite, littleleaf sumac, tar bush, creosotebush, and giant dropseed.

(b) (7)(E) is an ephemeral wash that directs flows (b) (7)(E). The channel width at base is approximately 4 feet. Channel banks are approximately 2 feet high with 45 percent slopes. Vegetation on the banks of the wash is characterized by creosotebush, white thorn acacia, littleleaf sumac, and tar bush.

(b) (7)(E) is a narrow ephemeral wash that (b) (7)(E). The channel width at base is approximately 3 feet. Channel banks are approximately 2 feet high with 35 percent slopes. Vegetation on the banks of the wash is characterized by tar bush, littleleaf sumac, white thorn acacia, and creosotebush.

(b) (7)(E) is a narrow ephemeral wash that (b) (7)(E). The channel width at base is approximately 3 feet. Channel banks are approximately 2 feet high with 45 percent slopes. Vegetation on the banks of the wash is characterized by creosotebush, tar bush, littleleaf sumac, little-head snakeweed, and white thorn acacia.

(b) (7)(E) is a narrow ephemeral wash that directs flows (b) (7)(E). The channel width at base is approximately 2 to 3 feet. Channel banks are approximately 4 feet high with 60 percent slopes. Vegetation on the banks of the wash is characterized by creosotebush, littleleaf sumac, white thorn acacia, little-head snakeweed, honey mesquite, and giant dropseed.

(b) (7)(E) is a narrow ephemeral wash that directs flows (b) (7)(E). (b) (7)(E) is comprised of two channels (b) (7)(E). The western channel directs flows to (b) (7)(E). The western channel's width at base is approximately 2 feet. Channel banks in the western channel are approximately 4 feet high with 45 percent slopes. The eastern channel directs flows (b) (7)(E). The channel width at base is approximately 2 feet. Channel banks in the eastern channel are near vertical and approximately 4 feet in height. Vegetation on the banks of both wash channels is characterized by white thorn acacia, creosotebush, and little-head snakeweed.

(b) (7)(E) is a narrow ephemeral wash downstream of an impoundment. The channel directs flows (b) (7)(E). Historical flow in the wash has been cut off by the impoundment and the channel has been disturbed. The channel width at base is approximately 20 feet. Channel banks are approximately 3 feet high with 35 percent slopes. Vegetation on the banks of the wash is characterized by Russian thistle, creosotebush, four-wing saltbush, honey mesquite, little-head snakeweed, and giant dropseed.

(b) (7)(E) is a wide ephemeral wash (b) (7)(E). The channel width at base is approximately 40 feet. Channel banks are near vertical and 3 feet to 6 feet in height. Vegetation on the banks of the wash is characterized by burrobush (*Hymenoclea monogyra*), little-head snakeweed, western sycamore (*Platanus wrightii*), velvet ash (*Fraxinus velutina*), western hackberry (*Celtis reticulata*), mule fat (*Baccharis salicifolia*), and honey mesquite.

(b) (7)(E) is an ephemeral wash that directs flows (b) (7)(E). The wash was cut to direct storm water flows (b) (7)(E) into (b) (7)(E). The channel width at base is approximately 12 to 15 feet. The channel banks are near vertical and approximately 3 feet in height. Vegetation on the banks of the wash is characterized by western hackberry, western sycamore, burrobush, and little-head snakeweed.

(b) (7)(E) is (b) (7)(E) extension of (b) (7)(E) that was evaluated independently due to a change in alignment. (b) (7)(E) has similar vegetation as W38 and ranges from 12 to 20 feet in width. (b) (7)(E) is north of the gated dirt road in (b) (7)(E); (b) (7)(E) is the area south.

4.4.2 Wetlands and Other Waters of the United States Vegetation Summary

Wetlands and other Waters of the United States delineated within the survey corridor included one palustrine emergent habitat, one palustrine

forested/riverine, a palustrine emergent habitat, and 37 ephemeral washes. The characteristic vegetation species for each wetland type sampled and delineated during the April/May 2008 field inventory are presented below by stand physiognomy.

4.4.3 Forest and Woodland Palustrine Forested

Four forested and woodland palustrine forested plant communities are found in the survey corridor. They include (1) Fremont Cottonwood – Goodding Willow Forest; (2) Fremont Cottonwood – Honey Mesquite Forest; (3) Arizona Sycamore – Fremont Cottonwood / Honey Mesquite Woodland; and (4) Honey Mesquite / Alkali Sacaton Woodland.

4.4.4 Shrubland

Two shrubland plant communities are found in the survey corridor. They include (1) Seepwillow – Burro Bush Shrubland; and (2) Honey Mesquite – Littleleaf Sumac Shrubland.

4.4.5 Herbaceous Palustrine Emergent

Three herbaceous palustrine emergent plant communities are found in the survey corridor. They include (1) Alkali Sacaton Herbaceous Vegetation; (2) Wild Barley / Honey Mesquite Shrub Herbaceous Vegetation; and (3) Bermuda Grass Herbaceous Vegetation.

4.4.6 Wetlands Soil Summary

Soils identified within vegetated wetlands in FV-1b exhibited hydric soil characteristics.

4.5 Noxious Weeds and Invasive Nonnative Plant Species

Noxious weeds have been addressed nationally under Public Law 108-412 (U.S.C. 2004) “Subtitle E – Noxious Weed Control and Eradication.” The Arizona legislature addressed noxious weeds under Title 3 – Agriculture; Chapter 2 – Regulatory Provisions; Article 1 – Dangerous Plant Pests and Diseases; Section 3-205.01 – Summary abatement of noxious weeds, crop pests, or diseases under preapproved programs (AZDA 2008). The survey corridor does not support Federal-listed (USDA 2006) noxious weeds. One state-listed noxious weed, a species of whitetop, occurred on the edges of the irrigated pasture of the (b) (7)(E) (AZDA 2008a). Eighteen nonnative plant species were observed on-site (see **Table 4-2**); thirteen were annuals and five were biennial or perennial. All nonnative species occurred on disturbed sites receiving higher moisture amounts than normally occur in this region; the sites included roadsides, excavated areas, sandy desert wash bottoms, and irrigated pasture.

In general, nonnative noxious and invasive plant species represent a serious management concern and their inventory, monitoring, and control can be expensive for land managers. Nonnative species usually lower the value of wildlife habitat and they increase with disturbance, including livestock grazing and road maintenance. Once inventoried, methods commonly used to control nonnative species include biological, mechanical, and chemical. Controls must be ongoing to be effective in reducing, but only rarely eliminating, nonnative plant species.

4.6 Protected Native Plants

The Arizona Department of Agriculture (AZDA) oversees rules associated with the use and harvest of native plants, including protected native plant species (see **Table 4-2**) (AZDA 2008b, 2008c). Four categories of protected native plants have been established by the AZDA (2008c):

1. Highly Safeguarded – prospects for survival in Arizona are in jeopardy or are in danger of extinction.
2. Salvage Restricted – subject to damage by theft or vandalism.
3. Salvage Assessed – have sufficient value if salvaged to support the cost of salvage.
4. Harvest Restricted – subject to excessive harvesting or overcutting because of their intrinsic value.

There were no highly safeguarded protected native plants observed within the (b) (7)(E) survey corridor. Fifteen species of “salvage restricted” protected native plants were observed (see **Table 4-2**); the most common of these were pencil cholla, species of yucca, and species of agave. Honey mesquite and desert willow represented the species of “salvage assessed” protected native plants to occur on-site (see **Table 4-2**). Honey mesquite was the single “harvest restricted” protected native plant observed (see **Table 4-2**).

In general, landowners have the right to destroy or remove plants growing on their land, but 20 to 60 days prior to the destruction of any protected native plants landowners are required to notify the AZDA (AZDA 2008). The landowner also has the right to sell or give away any plant growing on the land; however, protected native plants may not be legally possessed, taken, or transported from the growing site without a permit from the AZDA.

4.7 Wildlife and Wildlife Habitat

4.7.1 Introduction

Wildlife habitats of the survey corridor are predominantly (b) (7)(E) shrublands that at the highest elevations are characterized by ocotillo, tarbush, and mortonia. In the middle elevations, creosotebush, tarbush, and honey

mesquite compose the shrubland canopy. The lowest elevations support extensive honey mesquite shrublands and woodlands, gallery forests of Fremont cottonwood and Goodding willow, small stands of grasslands, and forb-dominated go-back agricultural fields.

The entire survey corridor occurs within the (b) (7)(E)

Historically, the basin bottom was a large cienega (marshy wetland) composed of herbaceous vegetation with a few honey mesquite trees and shrubs. Due to ground water pumping, surface water diversion, and farming and ranching pursuits, the basin bottom has become invaded by extensive stands of honey mesquite trees and shrubs. Riparian and wetland plant communities have become established along draws and washes with adequate surface and groundwater flows, on seeps, and adjacent to springs. Limited open water and aquatic habitat occurs. The semi-arid desert uplands contrast sharply with the lowland artesian wells, associated ponds, and mesic habitats.

Recreation (b) (7)(E) and its adjacent environs is centered on wildlife (USFWS 2008). Typical forms include birdwatching, landscape and wildlife photography, and hiking. In season, hunting for species of dove, quail, and desert cottontail rabbits is permitted on designated sites in the refuge. Within the survey corridor, hunting for mule and whitetail deer and collared peccary also occurs.

4.7.2 Wildlife and Habitat Overview

The survey corridor supports diverse populations and individuals of vertebrate wildlife species (see **Attachment C**) and unique-to-common native and nonnative wildlife habitats, described as vegetation alliances, plant associations, or land-use types in this BSR. **Table 4-4** lists wildlife observed during the field surveys that were conducted in early spring (April to May) of 2008. Along the international border, climate, geology, soils, land forms, geography, precipitation, and plant communities combine to provide moderate habitat diversity.

Table 4-4. Wildlife Species Observed Within the Survey Corridor, Staging Areas, and Associated Access Roads

Group / Scientific Name	Common Name	Relative Abundance
BIRDS		
<i>Falco sparverius</i>	American kestrel	Rare
<i>Corvus corax</i>	Common raven	Uncommon
<i>Corvus cryptoleucus</i>	Chihuahuan raven	Common
<i>Geococcyx californianus</i>	Greater roadrunner	Rare

Group / Scientific Name	Common Name	Relative Abundance
<i>Zenaida asiatica</i>	White-winged dove	Rare
<i>Zenaida macroura</i>	Mourning dove	Common
<i>Chordeiles acutipennis</i>	Lesser nighthawk	Uncommon
<i>Buteo jamaicensis</i>	Red-tailed hawk	Uncommon
<i>Buteo nitidus</i>	Gray hawk	Rare
<i>Eremophila alpestris</i>	Horned lark	Uncommon
<i>Callipepla gambelii</i>	Gambel's quail	Common
<i>Fulica americana</i>	American coot	Uncommon
<i>Zonotrichia leucophrys</i>	White-crowned sparrow	Common
<i>Tyrannus verticalis</i>	Western kingbird	Uncommon
<i>Cathartes aura</i>	Turkey vulture	Common
MAMMALS		
Unknown bat	Bat	Rare
<i>Canis latrans</i>	Coyote	Uncommon
<i>Odocoileus hemionus</i>	Mule deer	Uncommon
<i>Lepus californicus</i>	Black-tailed jackrabbit	Common
<i>Sylvilagus audubonii</i>	Desert cottontail	Abundant
REPTILES AND AMPHIBIANS		
<i>Elgaria kingii</i>	Madrean alligator lizard	Uncommon
<i>Heloderma suspectum</i>	Gila monster	Rare
<i>Phrynosoma</i> sp.	Horned lizard	Unknown
<i>Masticophis flagellum</i>	Coachwhip	Common
<i>Pituophis catenifer</i>	Gopher snake	Common
<i>Crotalus atrox</i>	Western diamondback rattlesnake	Common

Within the survey corridor, the broad habitat types available to resident and migrating wildlife species include herbaceous vegetation, shrubland, woodland, and forest. Most of the available wildlife habitat consists of semi-arid desert shrubland communities that have become established on ridges, slopes, alluvial fans and plains, and along arroyos, gullies, and desert washes. This section provides a brief summary of wildlife habitats observed and sampled in 2008 during Environmental Stewardship Plan preparation, categorized as follows:

1. Herbaceous Vegetation: This class of wildlife habitat includes annual and perennial species of grasses, forbs, and graminoids, which typically are characterized by no less than 15 percent cover by shrubs or trees. Stands of herbaceous vegetation range from less than 0.5 m to 2.0 m tall and low to dense in terms of cover. Herbaceous wildlife habitat occurs in small bunchgrass patches on ridges and slopes, small stands on disturbed sites of alluvial plains, and large stands within the cienega bottom.
 - a. *Grasslands* – On ridges and slopes of upper elevations, patches of tobosa occur within larger stands of ocotillo, creosotebush, and honey mesquite shrublands. Rock outcrops occasionally supported patches of little bluestem. One heavily grazed alluvial plain site near a livestock corral supported hook threeawn almost exclusively. Desert wash bottoms, particularly the broader ones, often supported stands of alkali and giant sacaton and scattered shrubs of honey mesquite and littleleaf sumac. One irrigated pasture was planted to Bermuda grass. Grassland dominated habitats occur on approximately 9.1 acres within the survey corridor and provide forage, escape cover, and breeding/nesting sites for several species of wildlife. Species common to grassland habitats include desert cottontail, pocket gophers, pocket mice, harvest mice, deer mouse, grasshopper mice, coyote, mule deer, falcons, hawks, turkey vulture, ravens, quail, doves, loggerhead shrike, sparrows, meadowlarks, toads, lizards, and snakes.
 - b. *Forblands* – Forbs, including baileya, slimleaf bursage, Russian-thistle, common cocklebur, and annual sunflowers are rare dominants within the survey corridor, becoming established on one disturbed alluvial plain site, in an excavation used for road fill material, and in the large fallow agricultural field that was once part of a cienega bottom. Forb-dominated habitats occur on approximately 6.2 acres within the survey corridor and provide forage, escape cover, and breeding/nesting sites for several species of wildlife. Species common to forbland habitats include desert cottontail, ground squirrels, pocket gophers, pocket mice, harvest mice, deer mouse, grasshopper mice, coyote, collared peccary, mule and whitetail deer, falcons, hawks, turkey vulture, ravens, quail, doves, loggerhead shrike, sparrows, goldfinch, meadowlarks, toads, lizards, and snakes.
 - c. *Emergent Wetlands* – Narrowleaf cattail, three-square bulrush, saltgrass, alkali sacaton, and sedges occur on the margins of ponds, seeps, springs, and on the banks of (b) (7)(E) occupying approximately 0.1 acre within the survey corridor. Emergent wetlands can be from 0.5 m to 3 m in height, dense, and along with associated aquatic habitat supporting diverse birds, mammals, reptiles, amphibians, fishes, and many invertebrates. Species common to emergent wetland and associated aquatic habitats include desert shrew, bats, pocket gophers, mice, raccoon, skunks, collared peccary, coyote, whitetail deer, American coot, ring-necked duck, gadwall, Mexican duck, mallard, great blue heron, pied-billed grebe, falcons,

hawks, quail, killdeer, doves, owls, flycatchers, vireos, swallows, wrens, northern mockingbird, warblers, sparrows, blackbirds, toads, leopard frogs, bullfrogs, Sonoran mud turtle, garter snakes, and fishes (minnows, suckers, catfish, topminnows).

2. Shrublands: This habitat class is dominant within the survey corridor, occupying approximately 404.2 acres. The characteristic upland shrubs range from 0.5 m to 5 m tall and include ocotillo, creosotebush, honey mesquite, tarbush, whitethorn, four-wing saltbush, shrubby coldenia, and mortonia. Characteristic shrubs of desert washes, creeks, and draws include honey mesquite, seepwillow, burro bush, and littleleaf sumac. Shrublands provide sparse to dense cover and are common on the ridges and hills of the western Project terminus.
 - a. *Dwarf-shrublands* – Dwarf-shrub stands occur on approximately 8.2 acres of exposed hilltop along access roads and bladed sites (b) (7)(E). This habitat is characterized by shrubby coldenia, four-wing saltbush, honey mesquite, and creosotebush that provided limited wildlife habitat. Common wildlife species likely to use this habitat include desert cottontail, black-tailed jackrabbit, ground squirrels, pocket mice, coyote, mule deer, turkey vulture, ravens, falcons, hawks, quail, doves, loggerhead shrike, sparrows, lizards, and snakes.
 - b. *Short Shrublands* – Stands of short shrubs occur throughout the survey corridor on approximately 385.2 acres of gravelly to cobbly ridges, hills, and slopes, on exposed bedrock of ridges, on alluvial fans and plains, and along desert washes and gullies. Short shrub stands are characterized by creosotebush, honey mesquite, whitethorn, tarbush, and mortonia primarily. Stands range from 1 m to 3 m tall and provide low to moderately high foliar cover. Nearly all wildlife species within the survey corridor use the short shrub habitats for forage, escape cover, breeding/nesting, and resting. The most common species include bats, desert cottontail, black-tailed jackrabbit, ground squirrels, pocket mice, deer mouse, other mice, kangaroo rats, coyote, gray fox, badger, bobcat, collared peccary, mule deer, Swainson's hawk, red-tailed hawk, American kestrel, turkey vulture, quail, greater roadrunner, owls, nighthawks, flycatchers, kingbirds, Chihuahuan raven, wrens, thrashers, sparrows, Madrean alligator lizard, collared lizard, horned lizards, colubrid snakes, and western diamondback rattlesnake.
 - c. *Tall Shrublands* – Stands of tall shrubs occur on ridges and slopes characterized by ocotillo and on slopes, along desert washes, and on flats characterized by honey mesquite. Tall shrubs typically range from 3 m to 6 m tall, this habitat type ranges from low to dense in terms of foliar cover, and approximately 90.1 acres occur in the survey corridor. Tall shrubs provide important perching, breeding, nesting, brood rearing, and escape cover for a variety of wildlife including bats, desert cottontail, ground squirrels, pocket mice, deer mouse, other mice,

kangaroo rats, coyote, gray fox, badger, bobcat, collared peccary, mule deer, Swainson's hawk, red-tailed hawk, American kestrel, turkey vulture, quail, owls, nighthawks, flycatchers, kingbirds, Chihuahuan raven, warblers, wrens, thrashers, towhees, sparrows, colubrid snakes, and western diamondback rattlesnake.

3. Woodlands and Forests: Open- to closed-canopy stands of trees occupy approximately 19.3 acres of (b) (7)(E) drainage banks and terraces. Fremont cottonwood, Arizona sycamore, and Goodding willow trees have become established as a gallery forest on the banks of (b) (7)(E). (b) (7)(E) supports Fremont cottonwood and honey mesquite trees (b) (7)(E) and honey mesquite trees updrainage. (b) (7)(E) is dominated by honey mesquite trees on an elevated terrace. Other drainages, terraces, and depressions support woodlands dominated by honey mesquite that cover approximately 78 acres. Woodlands typically provide moderate canopy cover and range between 4 m to 10 m tall and forest stands range between 10 m to 35 m tall, provide dense canopy cover, and usually have a subcanopy layer, which enhances the wildlife habitat value in terms of structure.
 - a. *Drainage Banks, Floodplain Terraces, and Springs* – The riparian gallery forest and woodland habitats of (b) (7)(E) support moderately open- to close-canopied stands dominated by Fremont cottonwood, Arizona sycamore, and honey mesquite. A moderately well-developed subcanopy in (b) (7)(E) stands provide additional wildlife habitat values. Numerous avifauna use the bank and terrace woodland habitat for foraging, breeding, nesting, brood rearing, perching, and escape cover, including the Swainson's hawk, red-tailed hawk, American kestrel, doves, owls, nighthawks, hummingbirds, Gila woodpecker, northern flicker, flycatchers, kingbirds, vireos, verdin, northern mockingbird, warblers, common yellowthroat, yellow-breasted chat, summer tanager, towhees, sparrows, northern cardinal, pyrrhuloxia, grosbeaks, and Bullock's oriole. Mammal use is high in these mesic habitats with common species and groups including bats, raccoon, desert cottontail, pocket gophers, skunks, mice, coyote, bobcat, collared peccary, and mule and white-tailed deer. Reptiles and amphibians common to the riparian habitats of these drainages include species of toads, lizards, colubrid snakes, and rattlesnakes. Moderate to high diversity of invertebrates occurs within these terrace woodlands and forests.
4. Open Water: Occupying approximately 0.1 acre within the Project area, open water habitats are species-rich in terms of wildlife use and as habituated for threatened and endangered amphibians and fishes. Water bodies occurred as a pond in (b) (7)(E) and a nearby pond supported by a flowing artesian well. Most water sources are ephemeral, flowing

following precipitation events of sufficient size to produce runoff and typically during the monsoon months of July through September. The bottom substrate of water bodies and ephemeral drainages is typically sand and fine sediments.

- a. *Creeks, Draws, and Desert Washes* – Flowing water habitat was not present during the early spring survey corridor survey and unvegetated dry washes occurred on approximately 2.8 acres. In addition to many unnamed desert washes, arroyos, and gullies are (b) (7)(E) which are ephemeral and flow primarily during the monsoon season. Because of active seeps and springs, (b) (7)(E) maintains ponded water throughout the year. These open water habitats and their associated riparian and wetland vegetation are extremely valuable to local and seasonal vertebrate and invertebrate wildlife species of the survey corridor.
 - b. *Lakes and Ponds* – Ponds occur within (b) (7)(E) on the alignment and where artesian wells discharge north of the alignment (b) (7)(E) (b) (7)(E). The wetland and riparian vegetation surrounding the shoreline and the size of the water body can dictate the species using still open water, which include the American coot, a variety of ducks, passerine birds, Sonoran mud turtle, leopard frogs, bullfrogs, Mexican and checkered garter snakes, endangered and threatened fishes, and insects.
5. Land Use: Small acreages in the survey corridor are maintained on a regular basis, ranging from monthly to yearly maintenance of (b) (7)(E) (b) (7)(E) to less periodic maintenance on secondary access roads and trails. Even though subject to disturbance, these habitats are somewhat important to many species of resident and migratory wildlife which use them as movement corridors, foraging sites, and sunning sites.
- a. *Irrigated Agriculture* – One small irrigated pasture of Bermuda grass occurs on approximately 0.3 acre in the survey corridor. It is grazed by horses and mules but also provides habitat for local wildlife that forage and seek water in this nonnative shortgrass habitat.
 - b. *Fallow or Go-Back Agriculture* – (b) (7)(E) Approximately 2.2 acres of this habitat support a large number of perennial grasses and annual forbs which range to 3 m tall and provide quantities of seed used by foraging wildlife. Seeds present on the go-back fields attract mule deer, desert cottontail rabbit, other small mammals, species of sparrows, and species of doves. Raptors and other predators regularly forage over or in this habitat.
 - c. *Highways, Roads, and Trails* – Wildlife species use established transportation corridors to move and disperse rapidly across the

landscape. As a result, low to moderately high death rates can be experienced depending on adjacent habitat importance to wildlife, population levels, and design speed and safety features of transportation corridors. The western diamondback rattlesnake and other snake species were observed sunning on (b) (7)(E) in the survey corridor. Wildlife that forage on carrion or are omnivorous, including the turkey vulture, other raptors, and coyote, can benefit from the presence of road-killed animals. Transportation structures such as bridges can provide hiding and roosting cover for species including owls and bats or nesting sites for swallows. Approximately 47.1 acres of this land use type occur in the survey corridor.

4.8 Species Groups and Habitat Affinity

4.8.1 Mammals

Fifty-five species of mammals have been recorded (b) (7)(E) habitats and also use adjacent landscapes of the survey corridor (see **Attachment C**). The largest species groups include bats (14) and mice, including pocket mice (10). Most of the mammals are nocturnal (night-active) or crepuscular (dusk- and dawn-active), and with the exception of the bat species are year-round residents. Black bear may traverse the survey corridor in search of forage. The rugged mountains that surround the (b) (7)(E) may support the very rare and federally endangered jaguar and ocelot, which have been recorded southeast and west of the area.

4.8.2 Birds

(b) (7)(E) the survey corridor support at least 268 bird species (see **Attachment C**) (USFWS 2008). Raptors that commonly use area habitats include American kestrel, peregrine falcon, red-tailed hawk, sharp-shinned hawk, Swainson's hawk, gray hawk, zone-tailed hawk, golden eagle, turkey vulture, and Chihuahuan raven (USFWS 2008). Aquatic birds and shorebirds that have been observed in the refuge include American coot, great blue heron, green-backed heron, Virginia rail, ringneck duck, Mexican duck, and sandhill crane. Other species and groups of birds common to the survey corridor include doves, greater roadrunner, owls, nighthawks, hummingbirds, flycatchers, loggerhead shrike, vireos, swallows, verdin, wrens, northern mockingbird, thrashers, warblers, tanagers, towhees, sparrows, grosbeaks, eastern meadowlark, and Bullock's oriole.

Large numbers of birds migrate seasonally through the (b) (7)(E) using the natural and managed habitats for forage, roosting, and cover. The drainages and linear mountain ranges can serve as leading lines to guide raptors and neotropical migrants during migration.

The establishment of the (b) (7)(E) in addition to other Federal, state, and private lands is important to migratory bird management. The primary function of lands managed under the National Wildlife Refuge System is to provide habitat for waterfowl and shorebirds in addition to other wildlife-related benefits. A focused list for species occurring in the survey corridor is presented in **Attachment C**.

The American peregrine falcon, a subspecies of the peregrine falcon, currently proposed for de-listing, is reported as a rare migratory visitor to the Project area (USFWS 1995). Under the *Peregrine Falcon Recovery Plan* the general goal is to restore a self-sustaining population of peregrine falcons in the western United States. The (b) (7)(E) and private landowners contribute towards restoration goals by conserving wintering and migratory habitats, protecting peregrine falcons through law enforcement efforts, and promoting public support and understanding through education.

The bald eagle has been de-listed from federally endangered to the threatened status throughout the United States except in the Sonoran Desert. Bald eagles are still protected by the Bald Eagle Protection Act. The *Bald Eagle Recovery Plan* efforts are undertaken to recover the species and in the Project region are considered significant efforts (USFWS 1995). The (b) (7)(E), and private landowners contribute towards restoration goals by ensuring that bald eagle habitats are protected and possibly enhanced. The refuge protects bald eagles through law enforcement efforts and promoting public support and understanding through education.

The northern aplomado falcon was a former resident of desert grasslands of southeastern Arizona that has been extirpated from the United States and contaminated by pesticides in Mexico (USFWS 1995). The *Aplomado Falcon Recovery Plan* included six objectives: (1) evaluate, monitor, and minimize all threats, including pesticides, to extant populations; (2) identify, maintain, and improve habitat; (3) re-establish the northern aplomado falcon in the United States and Mexico; (4) conduct studies of habitat requirements, physiological ecology, and behavior; (5) enhance public support for this recovery effort through educational programs; and (6) encourage national and international cooperation and coordination in carrying out these objectives. (b) (7)(E) was considered a possible re-introduction site for the northern aplomado falcon but further restoration of native grasslands and riparian woodlands/forests would be required. Northern aplomado falcons were planned to be released in Chihuahuan desert grassland habitats of southwestern New Mexico and were expected to spread into southeastern Arizona if releases were successful (Federal Register 2006, USFWS 2006). The goal of re-introduction would be to maintain a self-sustaining resident population of 60 breeding pairs between the years 2010 to 2030.

4.8.3 Reptiles and Amphibians

A species list of 39 reptiles and amphibians was compiled (b) (7)(E) (see **Attachment C**). During early spring wildlife surveys of the survey corridor, western diamondback rattlesnakes, coachwhips, the Gila monster, and lizard species were observed. Other reptile and amphibian species that could occur include the black-tailed rattlesnake, desert kingsnake, ringneck snake, Madrean alligator lizard, collared lizard, and horned toad in uplands and rock outcrops, while the wetland habitats support the Chiricahua leopard frog, Sonoran mud turtle, and checkered and Mexican garter snakes (USFWS 2008).

The federally threatened Chiricahua leopard frog population continues to decline due to habitat degradation, predation by nonnative bullfrogs and other wildlife species, and exposure to a lethal skin fungus (USFWS 2008). Management and protection of this rare leopard frog species in the Project area include efforts by the USFWS, AZGFD, University of Arizona, (b) (7)(E) High School, and private landowners.

The rare Mexican garter snake is one of three species of garter snake that inhabit riparian and wetland habitat in the survey corridor. The population on (b) (7)(E) has been reduced by predation by the nonnative bullfrog, which forages on young snakes. Restoration of densely vegetated cienega wetland habitats on (b) (7)(E) and the (b) (7)(E) assist with the recovery of the Mexican garter snake.

4.8.4 Fish

(b) (7)(E) is one of the few units within the refuge system created specifically to protect native fish; there are eight fish species (b) (7)(E) see **Attachment C**. (b) (7)(E) is focused on preserving the remaining fisheries habitat, restoring degraded habitats, and maintaining native fish populations in appropriate habitats. In addition to physical effects due to construction, (b) (7)(E) and construction staging could result in sedimentation into the refuge drainages. Soils of the former cienega are extremely fine silt and clay and are highly erodible both by water and wind action. The preferred crossing of (b) (7)(E) from a fisheries perspective would be a concrete low water crossing with a perpendicular downstream face high enough to impede fish from downstream from entering the refuge (Radke pers. comm. 2008).

Prior to drainage of wetlands and loss of permanent surface flows, (b) (7)(E) drainages and cienega supported approximately one-fourth of the fish species native to Arizona (USFWS 2008). The more common species were Mexican stoneroller, longfin dace, roundtail chub, and Yaqui sucker. Endangered and threatened fish species that occurred historically through present include the Yaqui chub, Yaqui topminnow, beautiful shiner, and Yaqui catfish.

The federally threatened beautiful shiner was eliminated from the United States by 1970 due to the loss of suitable wetland and aquatic habitat (USFWS 2008).

The current populations (b) (7)(E) were reintroduced from stock captured in the Sierra Madre of Mexico. The federally endangered Yaqui chub is confined to the upper Rio Yaqui Basin (b) (7)(E) and populations occur in (b) (7)(E) Arizona, and (b) (7)(E) in Mexico. Most of the populations are threatened due to infestations of nonnative Asian tapeworms which are parasites of the digestive system. The federally threatened Yaqui catfish was eliminated from U.S. waters and reestablished with Mexican stocks in 1996. The species now occurs in (b) (7)(E) Arizona. The federally endangered Yaqui topminnow was severely affected by the loss of wetland and aquatic habitat by vegetation encroachment and by the introduction of nonnative mosquitofish. It now occurs in (b) (7)(E), and in Mexico (b) (7)(E).

The *Fishes of the Rio Yaqui Recovery Plan* (USFWS 1994a) outlined the objectives required to recover the Yaqui chub, Yaqui topminnow, Yaqui catfish, and beautiful shiner as secure and stable elements of the native fish fauna of the river system where they once occurred. A combination of refuge protection strategies and habitat protection in Mexico is necessary for these rare fishes to be down-listed. There must be compliance with the following conditions for a period of five years before down-listing of the federally endangered Yaqui chub and Yaqui topminnow to federally threatened status can be considered: (1) all nonnative fish species must be eradicated from critical habitat; (2) secure and protect the San Bernardino aquifer so that all artesian flows maintain themselves year-round, secure and protect Leslie Creek watershed to ensure adequate flows for Leslie Creek; and (3) protect critical habitat from detrimental human disturbance including mining, introduction of nonnative fishes, water diversion, and removal.

4.8.5 Invertebrates

Southeastern Arizona has been described as an ecological crossroads due to intersecting geographies including the Chihuahuan and Sonoran deserts and southern Rocky Mountains and Sierra Madres. As a result, invertebrates are diverse, e.g., over 100 butterfly species occur in (b) (7)(E) and adjacent survey corridor support an abundance of butterflies, damselflies, and other invertebrates, including several unique tropical species. Some species of invertebrates have been documented in the United States only within the refuge (USFWS 2008).

The (b) (7)(E) springsnail occurs within a single small spring on (b) (7)(E) (b) (7)(E) and at two locations (b) (7)(E) in Mexico. Research is currently being conducted to determine the species habitat requirements or its tolerances.

4.9 Prehistoric Humans, Spanish Settlement, and Current Land Conservation

This section briefly summarizes human use of the survey corridor. Generally, the survey corridor uplands were used sparingly by prehistoric humans and historically for grazing livestock and farming. However, the (b) (7)(E) (b) (7)(E) with several permanent water sources, has attracted humans both prehistorically and historically resulting in the basis for much of the discussion herein. Forms of farming within the valley occurred over the past 800 years, reaching an apex and level of disturbance to the natural landscape between about 1900 to 1979 (USFWS 1995).

The upper Rio Yaqui watershed has supported humans since prehistoric times, with evidence of use dating over 10,000 years ago to the Clovis culture (USFWS 1994a). The principal prehistoric periods represented in the survey corridor were the late Archaic (approximately 1500 BC–500 BC) and the late prehistoric pueblo occupation (approximately AD 1200–AD 1400) (USFWS 1995).

Archaic humans practiced a hunting-gathering lifestyle throughout the desert southwest; in the Project area artifacts included projectile points (Pinto, Chiricahua, and San Pedro styles), ground stone artifacts, hearths, and roasting pits with the absence of ceramics (USFWS 1995). (b) (7)(E) supported a large semipermanent campsite during the Archaic Period.

Pueblo-dwelling humans that occupied permanent settlements and practiced agriculture settled in the Project area from approximately 600 to 800 years ago (USFWS 1995). This occupation has been named by various researchers as the Animas Phase, Casas Grandes, or Salado cultures, people who possessed architectural and material cultures with strong ties to similar advanced societies in Chihuahua. The largest pueblo period site or pit house village in the Project area is located on the historic (b) (7)(E); it consisted of approximately 100 rooms and one or more plazas located on a bench above (b) (7)(E) and (b) (7)(E) (USFWS 1995).

In approximately the early 1600s, the Apache Tribe frequented the Project region to hunt, gather food, and conduct raids (USFWS 2008). They frequented this region until 1886 when Geronimo and his forces surrendered in (b) (7)(E) (b) (7)(E).

The Coronado-led Spanish expedition passed near the survey corridor in search of the Seven Cities of Cibola during 1540 (USFWS 2008). European presence in the survey corridor dates to around 1694 when Jesuit Padre Eusabio Francisco Kino (an Italian priest) and Captain Juan Mateo Manje traveled through the (b) (7)(E) (Lanning 1981, NPS 2008). At that time there was an Opatá Indian village on the location of the (b) (7)(E).

Padre Kino established good relations with the indigenous Piman groups and assisted them in resisting the Apache tribes. He was also credited with introducing agriculture and animal husbandry including wheat and domestic livestock, particularly cattle and sheep. Jesuit priests established a mission in the (b) (7)(E) during the 1700s and a Spanish Presidio was established there in 1774 (USFWS 2008).

Feral livestock were abundant by 1822, when the 73,000-acre (b) (7)(E), created by decree of the Spanish Crown, was acquired by Lieutenant Ignacio de Perez (USFWS 1994). Large-scale cattle, mule, and horse grazing occurred for ten years (up to 100,000 head), until 1832, when Perez was driven from the valley by the Apache Tribe (USFWS 2008). This land grant included much of the survey corridor.

In 1846, the Mormon Battalion under Lieutenant Colonel Philip St. George Cooke passed through the (b) (7)(E) enroute to California. They encountered many wild cattle and one of the battalion noted that the grass was two feet high as far as the eye could see and there was plenty of water, but there was no wood barring mesquite (USFWS 1993). (b) (7)(E) near the entrance to the (b) (7)(E)

In 1853, the Gadsden Purchase placed the international boundary such that 2,383 acres of the original land grant/ranch lay within the United States and the remainder in Mexico. A total of 65,000 acres of the land grant were purchased by "Texas" John Slaughter (John Horton Slaughter, a Texas cattle rancher) in 1884 and was used to raise longhorn cattle (up to 50,000 head) and for farming (hay, barley, wheat, and vegetables) until 1937 (USFWS 1994). The center of a cattle ranching empire that straddled the U.S.-Mexico border, this ranch illustrated the continuity of Spanish and American cattle ranching in the Southwest. Until late in the nineteenth century, the (b) (7)(E), a well-watered area occupying southern Arizona and northern Mexico, was not successfully occupied by Europeans due to the threat of Apache attack; in 1884, however, Slaughter leased a portion of the Mexican land grant and began the development of a ranch that would span up to 100,000 acres, supplying beef, fruits, and vegetables to the surrounding settlements and military posts.

In 1915, the Mexican Revolution included this area of the border and General Pancho Villa and his troops fought in nearby Agua Prieta (USFWS 2008). Between 1914 to 1919, U.S. cavalry encampments were established in the valley to protect settlers from raids conducted by General Villa. U.S. troops were stationed at the (b) (7)(E) during this period and remnants of the rock fortifications remain on the (b) (7)(E)

Between 1937 and 1979, there were a number of owners who conducted cattle ranching and farming in the valley, until the (b) (7)(E) was purchased by The Nature Conservancy, transferred to the USFWS in 1982, and established as the (b) (7)(E) (USFWS 1994). Properties adjacent to the refuge and composing

the remainder of the survey corridor are primarily privately owned ranch lands and lands managed by the State of Arizona (USFWS 1995).

Farming, mineral extraction, fire control, and livestock production altered and eliminated much of the natural wildlife habitat in the Rio Yaqui Basin over the past 100-plus years (USFWS 2008). Some wetlands were drained to increase cropland acreage and streams were diverted to irrigate fields and fill water impoundments. Grasslands were diminished by unsustainable grazing practices and many surface water flows were eliminated. The very large (b) (7)(E) (b) (7)(E) (described as marshy and spring-fed) once persisted on both sides of the international border, but has been reduced to isolated artesian wells and artificial ponds and it supports old field vegetation. Stands of honey mesquite trees and shrubs have become established across the former cienega.

4.10 Habitat Restoration and monitoring

Extensive habitat restoration has been undertaken on the (b) (7)(E) and on adjacent private lands (USFWS 2008). Eroded and incised stream channels are being elevated and armored using wire-basket gabions filled with rocks and by planting Fremont cottonwood and Goodding willow trees. Upland habitats are being replanted to native grass species and old farm fields are being returned to former cienega wetland conditions. Invasive, nonnative species are being controlled or removed from the habitats. Controlled fires are being used to burn the litter from native grasses, to return nutrients to the soil, and to control the spread of honey mesquite trees and shrubs across the former cienega.

Cooperative conservation between U.S. government and environmentally sensitive landowners in Mexico and the United States is protecting additional habitat and water sources, providing additional scientific research, and allowing introductions and maintenance of fish and wildlife populations. Restored wetlands support waterfowl, riparian gallery forests support raptors and migrating passerine bird species, and measures are being enacted to reduce erosion, protect groundwater levels, and to reclaim honey mesquite-dominated lowlands. Monitoring is being designed and conducted to record the results of management actions, guide future management decisions, and to learn more about the complex ecological relationships.

5. LIST OF PREPARERS

1

2

(b) (6)

3

Graduate Studies, Natural Resources, USDA Graduate School, 1989-1992

4

Graduate Studies, Geology, 1983-86

5

B.S., Geology, 1976

6

Years of Experience: 25

7

8

(b) (6)

9

B.S. Ecology, Behavior and Evolution

10

Years of Experience: 15

11

12

(b) (6)

13

B.S., Biology, 2003

14

Years of experience: 5

15

16

(b) (6)

17

M.S. Masters of Engineering

18

Years of Experience: 5

19

(b) (6)

20

B.S. Biology

21

M.S. Environmental Science and Education

22

Years of Experience: 9

23

(b) (6)

24

B.S. Biology

25

M.S. Biology

26

Years of Experience: 32

1

THIS PAGE INTENTIONALLY LEFT BLANK

6. REFERENCES

- AZDA 2008a Arizona Department of Agriculture (AZDA). 2008a. *Article 1 – Dangerous Plant Pests and Diseases*. Accessed online at: <http://www.azleg.state.az.us/ArizonaRevisedStatutes.asp>.
- AZDA 2008b AZDA. 2008b. *New Native Plant Rules (May 3, 2008)*. Accessed online at: <http://www.azda.gov/Main/NewNativePlantRules.html>.
- AZDA 2008c AZDA. 2008c. *Protected Native Plant Lists A-D*. Accessed online at: <http://www.azda.gov/ESD/protplantlst.htm>.
- AZGFD 2008a Arizona Game and Fish Department (AZGFD). 2008a. *Status Definitions*. Accessed online at: <http://www.azgfd.gov>.
- AZGFD 2008b AZGFD. 2008b. *Special Status Species by County, Taxon, Scientific Name*. Heritage Data Management System. Phoenix, AZ. Available online at: http://www.azgfd.gov/w_c/edits/documents/ssspecies_bycounty_001.pdf.
- Bailey 1995 Bailey, Robert F. 1995. *Ecoregions of the United States*. U.S. Department of Agriculture - Forest Service. Available online at: <http://www.fs.fed.us/colormap/ecoreg>.
- Bennett et al. 2004 Bennett, Peter S., Michael R. Kunzmann, and Lee A. Graham. 2004. *Descriptions of Arizona Vegetation Represented on the GAP Vegetation Map*. USGS, Biological Resources Division. Phoenix, AZ.
- Burgess 1995 Burgess, Tony. 1995. *Desert Grassland, Mixed Shrub Savannah, or Semidesert Scrub?* In: McClaran, M. and T. Van Devender, eds. *The Desert Grassland*. University of Arizona Press. Tucson, AZ.
- Dimmitt 2000 Dimmitt, Mark. 2000. *Biomes and Communities of the Sonoran Desert Region*. In: Phillips, S. and P. Comus, eds., *A Natural History of the Sonoran Desert*. Arizona-Sonoran Desert Museum Press. Tucson, AZ.
- Earman 2002 Earman, S., B.J.O.L. McPherson, F.M. Phillips, S. Ralser, and J.M. Herrin. 2002. *Hydrogeologic Framework and Groundwater Characteristics, (b) (7)(E) [REDACTED], Arizona and Sonora*. Earth and Environmental Science Department, New Mexico Institute of Mining and Technology. Socorro, NM.

- Environmental Laboratory 1987 Environmental Laboratory. 1987. *Technical Report Y-87-1, Corps of Engineers Wetlands Delineation Manual*. Available online at: <http://el.erdc.usace.army.mil/wetlands/pdfs/wlman87.pdf>. Accessed 20 August 2008.
- Epple 1995 Epple, Anne E. 1995. *A Field Guide to the Plants of Arizona*. The Globe Pequot Press. Guilford, CT.
- Federal Register 2006 Federal Register. 2006. *Establishment of a Nonessential Experimental Population of Northern Aplomado Falcon in New Mexico and Arizona*. Volume 71, Number 143, July 26, 2006 - Rules and Regulations. 50 CFR, Part 17. RIN 1018-A180. Washington DC.
- Kearney 1960 Kearney, Thomas H. and Robert H. Peebles. 1960. *Arizona Flora*. Second Edition with Supplement by J.T. Howell and E. McClintock. University of California Press. Berkely, CA.
- Lanning 1981 Lanning, Dirk B. 1981. *The Vertebrates of* [REDACTED] (b) (7)(E) County, Arizona. Arizona Natural Heritage Program. Tucson, AZ.
- NPS 2008 National Park Service (NPS). 2008. *Tumacacori National Historic Park: Father Eusabio Francisco Kino*. Accessed online at: http://www.nps.gov/archive/tuma/Father_Kino.html.
- NatureServe and e²M 2007a NatureServe and e²M. 2007a. *Element Occurrence Data Request Form, NatureServe Central Databases*. Arlington, VA and Englewood, CO.
- NatureServe and e²M 2007b NatureServe and e²M. 2007b. *License Agreement between NatureServe and engineering-environmental Management, Incorporated*. Arlington, VA and Englewood, CO.
- NatureServe 2008 NatureServe Explorer. 2008. An online encyclopedia of life [web application]. Version 7.0. NatureServe, Arlington, Virginia. Available online: <http://www.natureserve.org/explorer>. Accessed: June 3, 2008.
- Neily and Beckwith 1985 Neily, Robert and Ronald Beckwith. 1985. *A Cultural Resource Inventory of the* (b) (7)(E) [REDACTED] Arizona State Museum. University of Arizona.
- New Mexico State University 2008 New Mexico State University. 2008. *The Chihuahuan Desert*. Accessed online at: <http://horizon.nmsu.edu/chihuahua.html>.

- NRCS 2008 Natural Resources Conservation Service (NRCS). 2008. *Electronic Field Office Technical Guide*, (b) (7)(E) County, Arizona. Section II- Water Resources- Arizona Watershed- 15080302 (b) (7)(E). Available online at: <http://efotg.nrcs.usda.gov/treemenuFS.aspx>. Accessed 4 September 2008.
- Robinson et al. 2006 Robinson, Anthony T., Nick V. Paretti, and Gail E. Cordy. 2006. *Ecological Assessment of Arizona's Streams and Rivers*. AGFD and USGS, Arizona Water Science Center. Phoenix and Tucson, AZ.
- U.S.C. 2004 United States Code (U.S.C.). 2004. Noxious Weed Control and Eradication Act of 2004. 118 Statute 2320; Public Law 108-412; 108th Congress. Washington. DC.
- USACE 1987 U.S Army Corps of Engineers (USACE). 1987. *Wetland Delineation Manual*. Technical Report Y-87-1. 1987.
- USDA 2008 U.S. Department of Agriculture (USDA). 2008a. *Natural Resources Conservation Service, PLANTS Database*. Accessed online at: <http://plants.usda.gov/>.
- USDA 2008c USDA. 2008c. *Federal Noxious Weed List and Arizona State-listed Noxious Weeds*. Accessed online at: <http://www.aphis.usda.gov/ppq/weeds/>.
- USEPA 2008 U.S. Environmental Protection Agency (USEPA). 2008. (b) (7)(E) Watershed – 15080302. Available online at: http://cfpub.epa.gov/surf/huc.cfm?huc_code=15080302.
- USFWS 1993 U.S. Fish and Wildlife Service (USFWS). 1993. (b) (7)(E) Pincushion Cactus (*Coryphantha robbinsorum*) Recovery Plan. USFWS. Albuquerque, NM.
- USFWS 1994a USFWS. 1994a. *Fishes of the Rio Yaqui Recovery Plan: 1995*. Kevin S. Cobble. (b) (7)(E) AZ. USDI Fish and Wildlife Service. Albuquerque, NM.
- USFWS 1994b USFWS. 1994b. *Lesser Long-nosed Bat Recovery Plan*. Theodore H. Fleming. USFWS Arizona Environmental Services State Office. Phoenix, AZ.
- USFWS 1995 USFWS. 1995. (b) (7)(E) Comprehensive Management Plan, 1995-2015. USFWS Region 2. Albuquerque, NM.
- USFWS 2003 USFWS. 2003. (b) (7)(E) Watchable Wildlife List. (b) (7)(E) AZ.

- USFWS 2006 USFWS. 2006. *Final Environmental Assessment for Reestablishment of the Northern Aplomado Falcon in New Mexico and Arizona*. USFWS, Region 2, New Mexico Ecological Field Office. Albuquerque, NM.
- USFWS 2008a USFWS. 2008a. *Threatened and Endangered Species System (TESS)*. Accessed online at: http://ecos.fws.gov/tess_public/StateListing.do.
- USFWS 2008b USFWS. 2008b. (b) (7)(E) Pamphlet. (b) (7)(E) AZ.
- WRCC 2008 Western Regional Climate Center (WRCC). 2008. *Climate Summaries for Yuma WSO AP, Arizona (0296660)*. Available online at: <http://www.wrcc.dri.edu/cgi-bin/>.
- WWF 2008 World Wildlife Fund (WWF). 2008. *Terrestrial Ecoregions of the World*. Accessed online at: <http://www.worldwildlife.org>.

**BIOLOGICAL SURVEY
ATTACHMENT A**

DESCRIPTION OF FEDERALLY LISTED SPECIES

Beautiful Shiner (*Cyprinella formosa*)

The beautiful shiner was designated as a federally Threatened species on August 31, 1984.

Historic range: Northern Mexico (Sonora, Chihuahua), southeastern Arizona (b) (7)(E) Creek and associated artesian wells and cienegas; extirpated by 1970), and southwestern New Mexico (Mimbres River; disappeared after 1951). Current range in Mexico: Guzman basin (including Rios Casas Grandes, Santa Maria, and del Carmen), and Yaqui, Bavicora, and Sauz basins (current status in Sauz Basin is unknown). Elevations in Mexico from 800-1700 m (2625-5580 ft), previously in Arizona at approximately 1,158 m (3,800 ft) (Arizona Game and Fish Department 1994). Reintroduced and thriving on the (b) (7)(E), Arizona (USFWS 1994). Stocks occur also at the Dexter National Fish Hatchery & Technology Center, Dexter, New Mexico. See USFWS (1994).

Basic Description: A 3-inch fish.

Reproduction Comments: Spawns probably in late spring.

Habitat Type: Freshwater

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): CREEK, Moderate gradient, Pool, Riffle, SPRING/SPRING BROOK

Lacustrine Habitat(s): Shallow water

Special Habitat Factors: Benthic

Habitat Comments: A mid-water-column species; remains near but rarely within beds of plants or other cover along pond margins (USFWS 1994). Occupies riffles in small streams or pools of creeks with riffles during high water; also in large streams in rapids and in small tanks and spring-fed ditches. Streams typically are intermittent and subject to seasonal drying and sudden flooding; survives dry periods in permanent pools. Uncommon in large rivers. Small turbid pools over sand, gravel, or boulder substrates (Miller and Simon 1943). Thriving in pond habitats on the (b) (7)(E) in Arizona (USFWS 1994). Eggs are laid in a nest scooped out of gravel by male in shallow, fast-flowing water.

Adult Food Habits: Herbivore, Invertivore

Immature Food Habits: Herbivore, Invertivore

Food Comments: Feeds mostly on terrestrial and aquatic insects; also eats algae and other plant matter.

Length: 7 centimeters

Management Requirements: Securing habitat and water sources is a major management need (USFWS 1994).

1 **References:**

- 2 • Arizona Game and Fish Department. 1994. *Cyprinella formosa mearnsi*. Unpublished
3 abstract, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- 4 • Chernoff, B., and R. R. Miller. 1982. *Notropis bocagrande*, a new cyprinid fish from
5 Chihuahua, Mexico, with comments on *Notropis formosus*. *Copeia* 1982:514-522.
- 6 • Lee, D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer,
7 Jr. 1980. Atlas of North American Freshwater Fishes. North Carolina State Museum of
8 Natural History. 867 pp.
- 9 • Matthews, J.R. and C.J. Moseley (eds.). 1990. The Official World Wildlife Fund Guide to
10 Endangered Species of North America. Volume 1. Plants, Mammals. xxiii + pp 1-560 +
11 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians,
12 Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180.
13 Beacham Publications, Inc., Washington, D.C.
- 14 • Mayden, R. L. 1989. Phylogenetic studies of North American minnows, with emphasis on
15 the genus *Cyprinella* (Teleostei: Cypriniformes). Univ. Kansas Museum Natural History
16 Miscellaneous Publication (80):1-189.
- 17 • Miller, R. R., and J. R. Simon. 1943. *Notropis mearnsi* from Arizona, an addition to the
18 known fish fauna of the United States. *Copeia* 1943:253.
- 19 • Nelson, J. S., E. J. Crossman, H. Espinosa-Perez, L. T. Findley, C. R. Gilbert, R. N. Lea,
20 and J. D. Williams. 2004. Common and scientific names of fishes from the United States,
21 Canada, and Mexico. American Fisheries Society, Special Publication 29, Bethesda, MD.
22 386 pp.
- 23 • Page, L. M., and B. M. Burr. 1991. A field guide to freshwater fishes: North America north
24 of Mexico. Houghton Mifflin Company, Boston, MA. 432 pp.
- 25 • Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B.
26 Scott. 1991. Common and scientific names of fishes from the United States and Canada.
27 American Fisheries Society, Special Publishing 20. 183 pp.
- 28 • Sublette, J. E., M. D Hatch, and M. Sublette. 1990. The fishes of New Mexico. University
29 New Mexico Press, Albuquerque, NM. 393 pp.
- 30 • Texas Natural History Collections [University of Texas at Austin]. 1997. February 7-last
31 update. North American Freshwater Fishes Index (Images, Maps and Information).
32 Online. Available: [http://www.utexas.edu/depts/tnhc/www.fish.tnhc/na/naindex.ht ml](http://www.utexas.edu/depts/tnhc/www.fish.tnhc/na/naindex.html).
33 Accessed 1997, April 4.
- 34 • U.S. Fish and Wildlife Service. 1990. Endangered and threatened species recovery
35 program: report to Congress. 406 pp.
- 36 • U.S. Fish and Wildlife Service. 1994. Yaqui fishes recovery plan. USDI Fish and Wildlife
37 Service, Albuquerque, NM. iv + 48 pp.
- 38 • U.S. Fish and Wildlife Service. 1995. San Bernardino and Leslie Canyon National Wildlife
39 Refuges. Comprehensive management plan, 1995-2015. Includes the environmental
40 assessment under seprate cover. Albuquerque, NM.

Desert Pupfish (*Cyprinodon macularius macularius*)

The desert pupfish was designated as a federally Endangered species on March 31, 1986.

Historic Range: Formerly in lower Colorado and Gila river drainages, southern Arizona to southeastern California, and in the Salton Sea and Laguna Salada basins, California and Mexico. Currently occurs in California in San Felipe Creek (lower reaches and associated wetlands; best non-native habitat in California) and San Sebastian Marsh (Imperial County) and Salt Creek (Riverside County; population there may not be viable); also in shoreline pools and irrigation drains in the Salton Sea area, where the species is scarce (Miller and Fuiman 1987). No native populations of *C. macularius* remain in Arizona (Minckley et al. 1991), but several reintroduced populations exist, and the species has been introduced in areas outside the native range. See Hendrickson and Varela (1989) for information on the status of several introduced populations in Arizona. Currently occurs in Sonora, Mexico, in Santa Clara Slough and several locations extending southeast from there, and in northeastern Baja California (notable is an apparently large population found at Cerro Prieto, Baja California) (Hendrickson and Varela 1989, Echelle et al. 2000). Several populations exist in artificial refugia.

Basic Description: A small chunky fish.

Reproduction Comments: Spawning: spring and summer, or year-round in warm constant temperature environments. Each female may lay 50-800 eggs or more/season, depending on her size (Moyle 1976). Males defend eggs. Eggs hatch in 10 days at 20 C (within about 3 days according to Matthews and Moseley 1990). Reproduces at age 2-3 months in constant warm temperatures; first breeds at about 1 year in variable temperatures. Up to 2-3 generations per year (Matthews and Moseley 1990).

Ecology Comments: Typically swims in loose schools, often in groups of similar size and age (Moyle 1976).

Habitat Type: Freshwater

Non-Migrant: Yes

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): CREEK, Low gradient, MEDIUM RIVER, Pool, SPRING/SPRING BROOK

Lacustrine Habitat(s): Shallow water

Palustrine Habitat(s): HERBACEOUS WETLAND

Habitat Comments: Desert springs and outflow marshes, river-edge marshes, backwaters, saline pools, and streams. Original habitat probably was marshes and flood plain pools along the lower Colorado River and springs throughout the Salton Sink. Prefers areas with sand/silt substrates and aquatic plant life, limited surface flow, water less than 1 m in depth. Tolerates low oxygen levels, high temperatures, and high salinity. May forage in shallows in early morning, deeper water most of day. Often rests on bottom, especially at night. May dive into anoxic bottom mud. Male establishes territory prior to spawning, usually in water less than 1 meter deep (sometimes deeper). Territory is typically 1 to 2 square meters or more (Moyle 1976). Eggs are laid on substrate of sand, mud, or perhaps preferentially on algal mat (Schoenherr 1988).

Adult Food Habits: Herbivore, Invertivore

Immature Food Habits: Herbivore, Invertivore

Food Comments: Opportunistic. Feeds on algae, detritus, and small invertebrates. In the Salton Sea eats ostracods, copepods, and some insects and pile worms. In other areas feeds on aquatic crustaceans, aquatic insect larvae, and molluscs (Moyle 1976).

Phenology Comments: May burrow into loose substrate and become dormant in winter when temperatures are extreme.

Length: 6 cm

Management Requirements: Introductions into marginal, semi-natural, relatively stable habitats have not been especially successful; recovery planners should consider use of riverine habitat and manipulations of flows or other disturbances (Hendrickson and Varela 1989). See Meffe and Vrijenhoek (1988) for a discussion of conservation genetics.

Monitoring Requirements: San Felipe Creek and Salt Creek populations are regularly monitored by California Dept. of Fish and Game (California Department of Fish and Game 1990).

References:

- California Department of Fish and Game (CDF&G). 1990. 1989 annual report on the status of California's state listed threatened and endangered plants and animals. 188 pp.
- Echelle, A. A., R. A. Van Den Bussche, T. P. Malloy, Jr., M. L. Haynie, and C. O. Minckley. 2000. Mitochondrial DNA variation in pupfishes assigned to the species *Cyprinodon macularius* (Atherinomorpha: Cyprinodontidae): taxonomic implications and conservation genetics. *Copeia* 2000:353-364.
- Echelle, A. A., and A. F. Echelle. 1993. Allozyme perspective on mitochondrial DNA variation and evolution of the Death Valley pupfishes (Cyprinodontidae: Cyprinodon). *Copeia* 1993:275-287.
- Echelle, A. A., and T. E. Dowling. 1992. Mitochondrial DNA variation and evolution of the Death Valley pupfishes (Cyprinodon, Cyprinodontidae). *Evolution* 46:193-206.
- Hendrickson, D. A., and A. Varela Romero. 1989. Conservation status of the desert pupfish, *Cyprinodon macularius*, in Mexico and Arizona. *Copeia* 1989:478-483.
- Hendrickson, D. A., and J. E. Brooks. 1991. Transplanting short-lived fishes in North American deserts: review, assessment, and recommendations. Pages 283-98 in W. L. Minckley and J. E. Deacon (editors). *Battle Against Extinction: Native Fish Management in the American West*. University of Arizona Press, Tucson, AZ.
- Lee, D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer, Jr. 1980. *Atlas of North American Freshwater Fishes*. North Carolina State Museum of Natural History. 867 pp.
- Matthews, J.R. and C.J. Moseley (eds.). 1990. *The Official World Wildlife Fund Guide to Endangered Species of North America*. Volume 1. Plants, Mammals. xxiii + pp 1-560 + 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians, Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180. Beacham Publications, Inc., Washington, D.C.
- Meffe, G. K., and R. C. Vrijenhoek. 1988. Conservation genetics in the management of desert fishes. *Conservation Biology* 2:157-169.
- Miller, R. R., and L. A. Fuiman. 1987. Description and conservation status of *Cyprinodon macularius eremus*, a new subspecies of pupfish from Organ Pipe Cactus National Monument, Arizona. *Copeia* 1987:593-609.

- 1 • Minckley, W. L. 1973. Fishes of Arizona. Arizona Game and Fish Department, Phoenix,
2 AZ.
- 3 • Minckley, W. L., G. K. Meffe, and D. L. Soltz. 1991. Conservation and management of
4 short-lived fishes: the cyprinodontoids. Pages 247-82 in W. L. Minckley and J. E. Deacon
5 (editors). Battle Against Extinction: Native Fish Management in the American West.
6 University of Arizona Press, Tucson, AZ.
- 7 • Minckley, W. L., and J. E. Deacon. 1991. Battle Against Extinction: Native Fish
8 Management in the American West. University of Arizona Press, Tucson. xviii + 517 pp.
- 9 • Moyle, P. B. 1976. Inland fishes of California. University of California Press, Berkeley,
10 CA. 405 pp.
- 11 • Nelson, J. S., E. J. Crossman, H. Espinosa-Perez, L. T. Findley, C. R. Gilbert, R. N. Lea,
12 and J. D. Williams. 2004. Common and scientific names of fishes from the United States,
13 Canada, and Mexico. American Fisheries Society, Special Publication 29, Bethesda, MD.
14 386 pp.
- 15 • Page, L. M., and B. M. Burr. 1991. A field guide to freshwater fishes: North America north
16 of Mexico. Houghton Mifflin Company, Boston, MA. 432 pp.
- 17 • Pister, E.P. (ed.). 1987. Proceedings of the Desert Fishes Council, Volume XIX, the
18 Nineteenth Annual Symposium. Desert Fishes Council, Bishop, CA.
- 19 • Pister, E.P. (ed.). 1991. Proceedings of the Desert Fishes Council, Volumes XX and XXI,
20 the Twentieth & Twenty-first Annual Symposia. Desert Fishes Council, Bishop, CA.
- 21 • Platania, S. P., K. R. Bestgen, M. A. Moretti, D. L. Propst, and J. E. Brooks. 1991. Status
22 of Colorado squawfish and razorback sucker in the San Juan River, Colorado, New
23 Mexico, and Utah. Southwestern Naturalist 36:147-150.
- 24 • Platania, S. P., K. R. Bestgen, M. A. Moretti, D. L. Propst, and J. E. Brooks. 1991. Status
25 of Colorado Squawfish and razorback sucker in the San Juan River, Colorado, New
26 Mexico, and Utah. Southwestern Naturalist 36:147-150.
- 27 • Platania, S.P. 1990. Biological summary of the 1987 to 1989 New Mexico-Utah
28 ichthyofaunal study of the San Juan River. Prepared by Department of Biology, University
29 of New Mexico, Albuquerque, for New Mexico Department of Game and Fish, Santa Fe,
30 under contract 78-516.6-01, and U.S. Bureau of Reclamation, Salt Lake City, UT, under
31 cooperative agreement 7-FC-40-05060.
- 32 • Roberts, B. and M. Moretti. 1989. Fisheries survey of the San Juan River, Utah, 1988.
33 Final. Prepared by Utah Division of Wildlife Resources, Salt Lake City, UT, under U.S.
34 Bureau of Reclamation Cooperative Agreement 7-FC-40-05050.
- 35 • Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B.
36 Scott. 1991. Common and scientific names of fishes from the United States and Canada.
37 American Fisheries Society, Special Publishing 20. 183 pp.
- 38 • Schoenherr, A. A. 1988. A review of the life history and status of the desert pupfish,
39 *Cyprinodon macularius*. Bull. Southern California Academy Science 87:104-134.
- 40 • Turner, B. J. 1984. Evolutionary genetics of artificial refugium populations of an
41 endangered species, the desert pupfish. Copeia 1984:364-369.
- 42 • U.S. Fish and Wildlife Service (USFWS). 1986. Determination of endangered status and
43 critical habitat for the desert pupfish. Federal Register 51(61):10642-50.
- 44 • U.S. Fish and Wildlife Service (USFWS). 1990. Endangered and threatened species
45 recovery program: report to Congress. 406 pp.

46

Gila Chub (*Gila intermedia*)

The Gila chub was designated as a federally Endangered species on August 9, 2002.

Historic Range: Historically occurred in springs and small streams in the upper Gila River basin in southern Arizona, southwestern New Mexico, and northeastern Sonora, Mexico (Miller and Lowe 1964, Minckley 1973, USFWS 2002). In Arizona, Gila chub are known to have occupied portions of the Salt, Verde, Santa Cruz, San Pedro, San Carlos, San Simon, San Francisco, and Agua Fria drainages and smaller tributaries of the mainstem Gila River. Small remnant populations remain in most of these drainages with the exception of the Salt and San Simon Rivers, where all known populations have been extirpated. An observation of a Gila chub in Turkey Creek in the upper Gila River Basin in New Mexico was made in 2001 (Telles, pers. comm., 2001, cited by USFWS 2002). The current known distribution of Gila chub in Mexico has been reduced to two small spring areas, Cienega los Fresnos and Cienega la Cienegita, adjacent to the Arroyo los Fresnos (tributary of the San Pedro River), within 2 km (1.2 mi) of the Arizona-Mexico border (Varela-Romero et al. 1992). No Gila chubs remain in the Mexican portion of the Santa Cruz River basin (Weedman et al. 1996).

Basic Description: A fish (chub) that typically is about 15 cm long.

General Description: Fishes of the genus *Gila* that occur in the Colorado River basin range from the streamlined *Gila elegans* of large rivers, through *G. robusta* of intermediate-sized rivers, to the thick-bodied *G. intermedia* of creeks and marshes (cienegas) (Minckley 1973, DeMarais 1986). The following description of Gila chub is mainly from Minckley (1969, 1973) and Rinne (1976). The Gila chub is a robust, darkly colored minnow. A typical Gila chub would be approximately 150 mm in length. Gila chub from Redfield Canyon ranged in size from 45-222 mm TL (n=113) (Griffith and Tiersch 1989). At ages 1-4 years, based on scale analysis, calculated lengths averaged 90, 135, 160, and 183 mm TL. Minckley (1969) reported that males are typically smaller than females. Gila chubs usually have eight dorsal, anal, and pelvic fin rays. Scales are large and number less than 80 and more than 61 in the lateral line. Scales are also thick and broadly imbricate, and basal radii are usually present. Vertebrae number from 38 to 45. Barbels are absent and pharyngeal teeth are in two rows (2,5-4,2 with some variation). Head length divided by caudal peduncle depth is 3.0 or less. Both sexes possess breeding tubercles, although their distribution is less extensive on females. Minckley (1969) gave the following description of breeding coloration: "Breeding coloration in this fish may be far more intense than in other forms of the genus in Arizona. The axial and inguinal regions become a deep orange-red, which may develop further into a broken, orange-red band along the lower sides and caudal peduncle, extending forward to include the branchiostegal rays and cheeks. The eyes of males become yellow to yellow-orange and the body is blue-black dorsally. Fins of some individuals, especially the larger ones, may be washed with lemon yellow." Larvae were described by Winn and Miller (1954).

Diagnostic Characteristics: The Gila chub is most similar morphologically to the roundtail chub. The latter usually is lighter colored, less robust, and with scales that are relatively smaller, thinner, and only slightly embedded; basal radii on scales are absent to weakly developed; the number of dorsal, anal and pelvic fin rays in roundtail chubs usually is nine; there are usually 81 or more scales in the lateral line and 43 to 49 total vertebrae; the length of the head divided by the depth of the caudal peduncle is typically 3.3 to 4.3, rarely greater than 4.0. The Yaqui chub, *Gila purpurea*, and the Sonora chub, *Gila ditaenia*, have radii strongly developed on all fields of scales, the mouth is horizontal to oblique, and a basicaudal spot is present albeit possibly discrete or diffuse. *Gila elegans* is distinctive as adults and may be distinguished from the Gila chub using characteristics described by Douglas et al. (1989). *Gila elegans* has been extirpated from areas where the Gila chub occurs and, unless reintroductions of these species occur, these three species will not be taken in the same collections.

- 1 **Reproduction Comments:** In Monkey Spring, a relatively-constant spring-fed pond,
2 reproduction may have last throughout late winter, spring, and summer, and perhaps into autumn
3 (Minckley 1969, 1985). In other areas it occurs mostly in late spring and summer (Minckley
4 1973). Most Gila chub probably mature in their second or third year of life (Griffith and Tiersch
5 1989).
- 6 **Ecology Comments:** The Gila chub is associated with a native fish fauna that includes loach
7 minnow (*Tiaroga cobitis*), spikedace (*Meda fulgida*), speckled dace (*Rhinichthys osculus*), longfin
8 dace (*Agosia chrysogaster*), Sonora sucker (*Catostomus insignis*) and desert sucker (*Pantosteus*
9 *clarki*). Historically, it also was associated with the woundfin (*Plagopterus argentissimus*),
10 bonytail (*Gila elegans*), squawfish (*Ptychocheilus lucius*), razorback sucker (*Xyrauchen texanus*),
11 and Gila topminnow (*Poeciliopsis occidentalis*), all of which are now extirpated from the Gila River
12 basin. Gila chub and roundtail chub are sometimes found in the same stream systems,
13 separated by only tens of meters; however, the two species have never been collected together
14 at the same site (DeMarais 1990; Minckley 1985, 1990).
- 15 **Habitat Type:** Freshwater
- 16 **Non-Migrant:** No
- 17 **Locally Migrant:** No
- 18 **Long Distance Migrant:** No
- 19 **Riverine Habitat(s):** CREEK, MEDIUM RIVER, Moderate gradient, Pool, Riffle,
20 SPRING/SPRING BROOK
- 21 **Palustrine Habitat(s):** HERBACEOUS WETLAND
- 22 **Habitat Comments:** Gila chubs commonly inhabit pools in smaller streams, springs, and
23 cienegas, and they can survive in small artificial impoundments (Miller 1946, Minckley 1973,
24 Rinne 1975). They are highly secretive, preferring quiet, deeper waters, especially pools, or
25 remaining near cover including terrestrial vegetation, boulders, and fallen logs (Minckley 1973,
26 Rinne and Minckley 1991). Minckley (1973) suggested that spawning may occur over beds of
27 aquatic plants. Specific habitat associations are known to vary ontogenetically and likely vary
28 seasonally and geographically. Young in Monkey Spring, Arizona (from which the species is now
29 extirpated), 25-75 mm total length (TL), were found in swifter areas than were adults, which
30 utilized undercut banks and heavily vegetated margins of the spring run (Minckley 1969). Griffith
31 and Tiersch (1989) collected Gila chubs from both riffles and pools in Redfield Canyon, Arizona.
- 32 **Adult Food Habits:** Herbivore, Invertivore
- 33 **Immature Food Habits:** Herbivore, Invertivore
- 34 **Food Comments:** Feeds mainly on aquatic and terrestrial insects and filamentous and
35 diatomaceous algae (Minckley 1973, Griffith and Tiersch 1989). Of 27 specimens examined for
36 stomach contents in Redfield Canyon, four contained remains of fishes; three contained
37 *Rhinichthys osculus* (Griffith and Tiersch 1989). Gila chubs were observed chasing Gila
38 topminnows in Monkey Spring (Minckley 1969). No information is available on dietary differences
39 between size or age classes. Larger individuals feed during evening and early morning hours,
40 whereas young chubs feed during all daylight hours (Minckley 1973, Griffith and Tiersch 1989).
- 41 **Phenology Comments:** Young are active throughout the day; larger individuals tend to be most
42 active in evening and early morning.

Stewardship Overview: Existing Gila chub populations need to be identified and carefully monitored. Protection would be enhanced by the elimination of detrimental water and land use practices and the removal of non-native fishes. Degraded habitats should be reclaimed and enhanced, and chubs should be reintroduced where chances for success are judged good. Research is needed to identify specific threats.

Restoration Potential: Lack of knowledge of the biology of Gila chub clearly is a deterrent to its recovery. Recovery potential is good only if critical habitat is vigorously protected. Remaining populations continue to be threatened by habitat modification and interactions with non-native fishes. Reestablishment in former range is problematic until the causes of the decline are corrected.

Preserve Selection & Design Considerations: Habitat in the form of headwater cienegas or spring-fed streams are critical for the continued existence of the Gila chub. Gila chub also does well in spring-fed ponds if non-native fish are excluded (Minckley 1969).

Management Requirements: Existing populations not infected by non-native fishes should be protected through the establishment of fish barriers if such is judged not to be detrimental to the Gila chub. Necessary habitat and landscape improvements (including removal of non-native fishes) need to be determined and implemented. Stream flows and temperatures should not be modified by activities such as damming or diversion that substantially alter natural regimes. State or other fish management agencies and private entities should discontinue stockings of non-native, warmwater sport, forage, or bait fishes into streams occupied by Gila chub; this protection should extend downstream at least to the first absolute barrier to upstream fish movement. Proper management and maintenance of riparian zones are essential to native fish populations. Changes in the riparian zone can affect leaf fall and energy flow, stream flow, natural cover, temperature, and deposition of eroded materials (Baltz and Moyle 1984). Of five riparian systems studied in Arizona, only Aravaipa Creek, where cattle have been excluded since 1973, showed successful reproduction and dominance of the broadleaf riparian community (Rucks 1984). Cattle-browsing is a major factor in the replacement of a broadleaf riparian community by a riparian scrub community (Rucks 1984). A change from a broadleaf to scrub riparian community can change energy flow, tree-fall cover and amount of shade, and temperature profiles of a stream. Fire would be a preferred method of watershed management when necessary. However, the choice of fire as a management tool must take into account the fuel levels present. A crown fire ("hot fire") can lead to increased runoff and result in the filling of riffle or other spawning areas. The effects of a crown fire and subsequent runoff were reversed in three years in the upper Carmel River, California (Hecht 1984). If watershed management is necessary, controlled burns, frequent enough to prevent build-up of high fuel levels, set during nonspawning periods or periods of decreased spawning activity (winter), should be employed. Populations should be reintroduced into selected streams within the historic range. Potential dispersal routes should be closed to preclude reinvasion of non-native fishes. Barrier design should not significantly alter stream flow and the potential impact on natural upstream and downstream movements of native fishes should be assessed. Habitat improvement should be implemented, which may include removal of non-native fishes by piscicide. Reintroduced stocks should have a genetic affinity with those formerly occupying target streams. Stockings should be done according to guidelines set up by the American Fisheries Society (Williams et al. 1988), consultants familiar with GILA taxonomy, and the U.S. Fish and Wildlife Service. Reintroduced populations should be monitored for success or failure. Populations that are rapidly declining should be secured in a hatchery facility such as the Dexter National Fish Hatchery, Dexter, New Mexico. Techniques for spawning and rearing *Gila* spp. are available (Hamman 1981, 1982, 1982, Muth et al. 1985).

Monitoring Requirements: Known populations should be monitored biannually in the spring during the breeding season and in late autumn to check recruitment. Standardized techniques should be adopted so that data will be comparable over locations and time. Data are needed to distinguish between natural fluctuations in abundance and population decline due to human-

caused perturbation. Monitoring locations for Gila chub should be chosen so that all drainages and morphological variants represented. Techniques available for determination of absolute abundance for fishes are depletion sampling, mark-and-recapture, underwater censusing, and passive capture devices. These may be modified or others developed specifically for application to the Gila chub. Such techniques should be adjusted as dictated by experience, and uniformly applied. *Gila* species can be difficult to collect as they will flee when approached. They are often located under or next to objects, making seining difficult; as a result, electroshocking devices may provide more efficient sampling. Large areas must be sampled to determine presence/absence of Gila chub because certain areas are used, sometimes consistently over time, and others, which may be similar, are not (Minckley 1990). If resources are limited, a better strategy is to sample an entire headspring-cienega-stream system thoroughly every two to three years rather than sampling annually small areas of a stream or cienega system. DeMarais (1990) and Minckley (1990) stated that Gila chub occurrences are extremely spotty and localized. When chub populations are located, these data could be recorded on aerial photographs, and these photographs used to relocate chub populations. Recording these data on aerial photographs might also reveal clues about other stream reaches that have appropriate Gila chub habitat.

Biological Research Needs: The impact of flooding on nutrient cycling, substrate renewal, and availability of cover, with respect to native fishes, needs to be examined.

References:

- Arizona Game and Fish Commission. 1988. Threatened native wildlife in Arizona. Arizona Game and Fish Department Publication, Phoenix, AZ. 32 pp.
- Baltz, D. M., and P. B. Moyle. 1984. The influence of riparian vegetation on steam fish communities of California. Pp. 183-7 in R. E. Warner and K. M. Hendrix (eds.). California riparian systems. University of California Press, Berkeley, CA.
- Bestgen, K. R., and D. L. Propst. 1989. Distribution, status, and notes on the ecology of *Gila robusta* (Cyprinidae) in the Gila River drainage, New Mexico. Southwestern Naturalist 34:402-412.
- DeMarais, B. D. 1986. Morphological variation in Gila (Pisces: Cyprinidae) and geologic history: lower Colorado River Basin. Arizona State University, Tempe, AZ. M.S. Thesis.
- Deacon, J. E. and W. L. Minckley. 1974. Desert Fishes. Pp. 385-488 in G. W. Brown (ed.). Desert Biology, Vol. II. Academic Press, Inc., New York, NY.
- Douglas, M. E., W. L. Minckley, and H. M. Tyus. 1989. Qualitative characters, identification of Colorado River chubs (Cyprinidae: genus *Gila*) and the "art of seeing well." Copeia 1989:653-662.
- Griffith, J. S. and T. R. Tiersch. 1989. Ecology of fishes in Redfield Canyon, Arizona, with emphasis on *Gila robusta intermedia*. Southwestern Naturalist 34:131-4.
- Hamman, R. L. 1981. Hybridization of three species of chub in a hatchery. Prog. Fish-Cult. 43:140-1.
- Hamman, R. L. 1982. Induced spawning and culture of bonytail chub. Prog. Fish-Cult. 44:201-3.
- Hamman, R. L. 1982. Spawning and culture of humpback chub. Prog. Fish-Cult. 44:213-6.
- Hastings, J. R. 1959. Vegetation change and arroyo cutting in southeastern Arizona. J. Arizona Academy Science 1:60:67.
- Hecht, B. 1984. Sequential changes in bed habitat conditions in the upper Carmel River following the Marble-Cone fire of August, 1977. Pp. 134-41 in R. E. Warner and K. M. Hendrix (eds.). California riparian systems. University of California Press, Berkeley, CA.

- 1 • Hendrickson, D. A. and W. L. Minckley. 1984. Cienegas--Vanishing climax communities
2 of the American Southwest. *Desert Plants*. 6:141-75.
- 3 • Hubbs, C. L. 1954. Establishment of a forage fish, the red shiner (*Notropis lutrensis*), in
4 the lower Colorado River system. *California Fish and Game* 40:287-94.
- 5 • Meffe, G. K. 1985. Predation and species replacement in American southwestern fishes:
6 a case study. *Southwestern Naturalist* 30:173-187.
- 7 • Meffe, G. K., and W. L. Minckley. 1987. Persistence and stability of fish and invertebrate
8 assemblages in a repeatedly disturbed Sonoran desert stream. *American Midland*
9 *Naturalist* 117:177-91.
- 10 • Miller, R. R. 1961. Man and the changing fish fauna of the American Southwest. Paper
11 Michigan Academy Science, Arts, and Letters 46:365-404.
- 12 • Miller, R. R. and C. H. Lowe. 1964. Part 2: An annotated check-list of the fishes of
13 Arizona. Pp. 133-51 in C. H. Lowe (ed.). *The vertebrates of Arizona*. University of Arizona
14 Press, Tucson, AZ.
- 15 • Minckley, W. L. 1969. Aquatic biota of the Sonoita Creek Basin, Santa Cruz County,
16 Arizona. *Ecological Studies Leaflet*, No. 15. 8 pp.
- 17 • Minckley, W. L. 1973. *Fishes of Arizona*. Arizona Game and Fish Department, Phoenix,
18 AZ.
- 19 • Minckley, W. L. 1985. Native fishes and natural aquatic habitats in U.S. Fish and Wildlife
20 Service Region II west of the Continental Divide. Report to U.S. Fish and Wildlife Service,
21 Albuquerque, NM. 158 pp.
- 22 • Minckley, W. L., and G. K. Meffe. 1987. Differential selection by flooding in stream-fish
23 communities of the arid American southwest. Pp. 93-104 in W. J. Matthews and D. E.
24 Heins (eds.). *The zoogeography of North American freshwater fishes*. John Wiley and
25 Sons, Incorporated, NY.
- 26 • Minckley, W. L., and J. E. Deacon. 1968. Southwestern fishes and the enigma of
27 "endangered species." *Science* 159:1424-32.
- 28 • Muth, R. T., C. M. Haynes and C. A. Carlson, 1985. Culture of roundtail chubs, *Gila*
29 *robusta robusta* (Cyprinidae), through the larval period. *Southwestern Naturalist* 30:152-
30 4.
- 31 • Nelson, J. S., E. J. Crossman, H. Espinosa-Perez, L. T. Findley, C. R. Gilbert, R. N. Lea,
32 and J. D. Williams. 2004. Common and scientific names of fishes from the United States,
33 Canada, and Mexico. American Fisheries Society, Special Publication 29, Bethesda, MD.
34 386 pp.
- 35 • Page, L. M., and B. M. Burr. 1991. A field guide to freshwater fishes: North America north
36 of Mexico. Houghton Mifflin Company, Boston, MA. 432 pp.
- 37 • Propst, D. L., P. C. Marsh and W. L. Minckley. 1985. Arizona survey for spikedace (*Meda*
38 *fulgida*) and loach minnow (*Tiaroga cobitis*): Fort Apache and San Carlos Apache Indian
39 Reservations and Eagle Creek, 1985. Report to Office of Endangered Species, U.S. Fish
40 and Wildlife Service, Albuquerque, NM. 8 pp.
- 41 • Rinne, J. N. 1976. Cyprinid fishes of the genus *Gila* from the lower Colorado River Basin.
42 *Wassman J. Biology* 34:65-107.
- 43 • Robins, C. Richard, et. al. 1980. A List of Common and Scientific Names of Fishes from
44 the United States and Canada. 4th edition. American Fisheries Society, Special
45 Publication No. 12, Bethesda, MD.

- 1 • Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B.
2 Scott. 1991. Common and scientific names of fishes from the United States and Canada.
3 American Fisheries Society, Special Publishing 20. 183 pp.
- 4 • Rucks, M. G. 1984. Composition and trend of riparian vegetation on five perennial
5 streams in southeastern Arizona. Pp. 97-107 in R. E. Warner and K. M. Hendrix (eds.).
6 California riparian systems. University of California Press, Berkeley, CA.
- 7 • Sublette, J. E., M. D Hatch, and M. Sublette. 1990. The fishes of New Mexico. University
8 New Mexico Press, Albuquerque, NM. 393 pp.
- 9 • U.S. Fish and Wildlife Service. 9 August 2002. Listing the Gila chub as endangered with
10 critical habitat. Federal Register 67(154):51948-51985.
- 11 • Vanicek, C. D., and R. H. Kramer. 1969. Life history of the Colorado squawfish,
12 *Ptychocheilus lucius*, and the Colorado chub, *Gila robusta*, in the Green River in
13 Dinosaur National Monument, 1964-1966. Trans. American Fish. Soc. 98:193-195.
- 14 • Varela-Romero, A., et al. [9 other authors]. 1990. Re-discovery of *Gila intermedia* and
15 *Gila purpurea* in northern Sonora, Mexico. Proceedings of the Desert Fishes Council
16 22:33.
- 17 • Weedman, D. A., A. L. Girmendonk, and K. L. Young. 1996. Status review of Gila chub,
18 *Gila intermedia*, in the United States and Mexico. Technical Report 91, Nongame and
19 Endangered Wildlife Program, Arizona Game and Fish Department, Phoenix. 120 pp.
- 20 • Williams, J. E. and D. W. Sada. 1985. America's desert fishes: increasing their protection
21 under the Endangered Species Act. Endangered Species Tech. Bull. 10:8-14.
- 22 • Williams, J. E., D. W. Sada, C. D. Williams, et. al. 1988. American Fisheries Society
23 guidelines for introductions of threatened and endangered fishes. Fisheries. 13:5-11.
- 24 • Winn, H. E., and R. R. Miller. 1954. Native postlarval fishes of the lower Colorado River
25 basin, with a key to their identification. California Fish and Game 40:273-85. 4 plates.
26

Yaqui Chub (*Gila purpurea*)

The Yaqui chub was designated as a federally Enangered species on August 31, 1984.

Historic range: Historically the Yaqui chub occurred in the Rio Yaqui drainage in (b) (7)(E) County, extreme southeastern Arizona, USA, and in a short perennial reach of the (b) (7)(E) (b) (7)(E) just south of the U.S./Mexico border in Sonora, Mexico. Current distribution in Mexico is unknown. The species was nearly extirpated in the United States, persisting only in one artesian well in (b) (7)(E) Creek drainage (McNatt 1974). It was introduced and established in (b) (7)(E) Creek, Swisshelm Mountains, Arizona, in 1969 (Minckley 1973). Records from Morse Canyon, northern Chiracahua Mountains, Arizona, are not supported by specimens (Willcox Playa basin; McNatt 1974). In the United States, populations are limited primarily to several sites in the (b) (7)(E) and (b) (7)(E) (b) (7)(E), (b) (7)(E) County, Arizona. Populations from the drainages of the Rio Sonora, Rio Matape, and portions of the Rio Yaqui in Sonora, Mexico, formerly were included in *G. purpurea*; they were described as a new species (*Gila eremica*) by DeMarais (1991).

Basic Description: A fish less than six inches long.

Reproduction Comments: Spawning occurs throughout the warmer months, with greater activity in spring; matures often within the first summer; high reproductive potential (USFWS 1994).

Ecology Comments: Large populations develop quickly from a few adults.

Habitat Type: Freshwater

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): CREEK, Moderate gradient, Pool

Habitat Comments: Habitat includes deep pools in creeks, springheads, scoured areas of cienegas, and other stream-associated quiet waters (USFWS 1994); this fish seeks shade, often near undercut banks or debris; it is often associated with higher aquatic plants (Lee et al. 1980). Similarly, in artificial ponds, adults tend to occupy the lower part of the water column and seek shade (USFWS 1994). Young occupy near-shore zones, often near the lower ends of riffles (USFWS 1994). Spawning occurs probably in deep pools where there is aquatic vegetation (Matthews and Moseley 1990).

Adult Food Habits: Herbivore, Invertivore, Piscivore

Immature Food Habits: Herbivore, Invertivore, Piscivore

Food Comments: Eats algae, terrestrial insects, and arachnids. Aquatic insects and small fishes (Poeciliopsis) are eaten when available; also detritus (Matthews and Moseley 1990).

Stewardship Overview: Actions needed (USFWS 1994): 1) Develop co-operative effort with Mexico for the recovery of Yaqui fishes; 2) Secure habitat and water sources for the Yaqui fishes in the USA and Mexico; 3) Conduct research on the biology and habitat requirements of Yaqui fishes; 4) Manage the fish and their essential habitats; 5) Introduce and maintain self-sustaining populations within their historic range; and 6) Monitor existing and established populations and habitats. Management needs: protect (b) (7)(E) aquifers, and (b) (7)(E)

watersheds, to ensure adequate perennial flow; ameliorate effects of non-native fishes within chub management streams; establish and maintain self-sustaining populations on [REDACTED], and West Turkey Creek (Arizona Game and Fish Department 2001).

Management Requirements: Securing habitat and water sources are important management needs. See recovery plan (USFWS 1994).

References:

- Arizona Game and Fish Department. 2001. *Gila purpurea*. Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- DeMarais, B. D. 1991. *Gila eremica*, a new cyprinid fish from northwestern Sonora, Mexico. *Copeia* 1991:178-189.
- Hendrickson, D. A., and J. E. Brooks. 1991. Transplanting short-lived fishes in North American deserts: review, assessment, and recommendations. Pages 283-98 in W. L. Minckley and J. E. Deacon (editors). *Battle Against Extinction: Native Fish Management in the American West*. University of Arizona Press, Tucson, AZ.
- Hendrickson, D., W. L. Minckley, R. R. Miller, D. J. Siebert, and P. H. Minckley. 1980. Fishes of the Rio Yaqui Basin, Mexico and the United States. *J. Ariz. Acad. Sci.* 15(3):65-106.
- Kline, J., T. Archdeacon, A. C. Iles, and S. A. Bonar. 2007. Factors influencing distribution of introduced Asian tapeworm and effects on selected southwestern fishes (Yaqui topminnow and Yaqui chub). Arizona Cooperative Fish and Wildlife Research Unit Fisheries Research Report 01-07.
- Lee, D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer, Jr. 1980. *Atlas of North American Freshwater Fishes*. North Carolina State Museum of Natural History. 867 pp.
- Matthews, J.R. and C.J. Moseley (eds.). 1990. *The Official World Wildlife Fund Guide to Endangered Species of North America*. Volume 1. Plants, Mammals. xxiii + pp 1-560 + 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians, Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180. Beacham Publications, Inc., Washington, D.C.
- McNatt. R.M. 1974. Re-evaluation of the native fishes of the Rio Yaqui in the United States. *Proc. Ann. Conf. West. Assoc. Game Fish Comm.* 54:273-279.
- Minckley, W. L. 1973. *Fishes of Arizona*. Arizona Game and Fish Department, Phoenix, AZ.
- Minckley, W. L., and J. E. Deacon. 1991. *Battle Against Extinction: Native Fish Management in the American West*. University of Arizona Press, Tucson. xviii + 517 pp.
- Nelson, J. S., E. J. Crossman, H. Espinosa-Perez, L. T. Findley, C. R. Gilbert, R. N. Lea, and J. D. Williams. 2004. *Common and scientific names of fishes from the United States, Canada, and Mexico*. American Fisheries Society, Special Publication 29, Bethesda, MD. 386 pp.
- Page, L. M., and B. M. Burr. 1991. *A field guide to freshwater fishes: North America north of Mexico*. Houghton Mifflin Company, Boston, MA. 432 pp.
- Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B. Scott. 1991. *Common and scientific names of fishes from the United States and Canada*. American Fisheries Society, Special Publishing 20. 183 pp.

- 1 • U.S. Fish and Wildlife Service. 1990. Endangered and threatened species recovery
2 program: report to Congress. 406 pp.
- 3 • U.S. Fish and Wildlife Service. 1984. Final rule to determine the Yaqui chub to be an
4 endangered species with critical habitat, and to determine the beautiful shiner and Yaqui
5 catfish to be threatened species with critical habitat. Federal Register 49(171):34490-
6 34497.
- 7 • U.S. Fish and Wildlife Service. 1994. Yaqui fishes recovery plan. USDI Fish and Wildlife
8 Service, Albuquerque, NM. iv + 48 pp.
- 9 • U.S. Fish and Wildlife Service. 1995. San Bernardino and Leslie Canyon National Wildlife
10 Refuges. Comprehensive management plan, 1995-2015. Includes the environmental
11 assessment under seprate cover. Albuquerque, NM.
- 12 • Williams, J. E., J. E. Johnson, D. A. Hendrickson, S. Contreras-Balderas, J. D. Williams,
13 M. Navarro-Mendoza, D. E. McAllister, and J. E. Deacon. 1989. Fishes of North America
14 endangered, threatened, or of special concern: 1989. Fisheries 14:2-20.

15

Yaqui Catfish (*Ictalurus pricei*)

The Yaqui catfish was designated as a federally Threatened species on August 31, 1984.

Historic range: Originally described from the (b) (7)(E), Sonora. Historical range most likely included the uppermost Rio Yaqui system, Arizona, and the basins of the Rio Yaqui and Rio Casas Grandes, Sonora and Chihuahua, Mexico (USFWS 1994). Now definitely known only from the Rio Yaqui basin, Mexico, though catfishes in other basins to the south may be this species. An introduced population existed in Arizona in the Santa Cruz River system (in a reservoir fed by Monkey Spring) from 1899 to the 1950s (Minkley 1973, Lee et al. 1980). As of the mid-1990s, stock was being held at Dexter National Fish Hatchery for future reintroduction onto the (b) (7)(E) in Arizona.

Basic Description: A small catfish.

Ecology Comments: Little information on life history available but probably similar to channel catfish (Minckley 1973). Grows rapidly and attains large sizes in ponds at Dexter NFHTC.

Habitat Type: Freshwater

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): CREEK, MEDIUM RIVER, Moderate gradient

Special Habitat Factors: Benthic

Habitat Comments: Small to medium rivers; most abundant in larger rivers in medium to slow currents over gravel/sand substrate.

Length: 50 cm

Management Requirements: Securing habitat and water sources is a major management need (USFWS 1994). Could be reintroduced in the (b) (7)(E) if sufficient habitat there can be secured and maintained. However, leases on geothermal resources granted by BLM on lands adjacent to the NWR possibly could result in a decrease in the already diminished water tables in the region, and the threat of pollution of groundwater could be increased. These threats are to be evaluated by BLM in consultation with the USFWS.

References:

- Hendrickson, D. A., et al. 1981. Fishes of the Rio Yaqui, Mexico and United States. J. Arizona-Nevada Academy Science 15(3; 1980):65-106.
- Lee, D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer, Jr. 1980. Atlas of North American Freshwater Fishes. North Carolina State Museum of Natural History. 867 pp.
- Lundberg, J. G. 1992. The phylogeny of ictalurid catfishes: a synthesis of recent work. Pages 392-420 in R. L. Mayden, editor. Systematics, historical ecology, and North American freshwater fishes. Stanford Univ. Press, Stanford, CA. xxvi + 969 pp.
- Matthews, J.R. and C.J. Moseley (eds.). 1990. The Official World Wildlife Fund Guide to Endangered Species of North America. Volume 1. Plants, Mammals. xxiii + pp 1-560 +

33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians, Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180. Beacham Publications, Inc., Washington, D.C.

- Minckley, W. L. 1973. Fishes of Arizona. Arizona Game and Fish Department, Phoenix, AZ.
- Nelson, J. S., E. J. Crossman, H. Espinosa-Perez, L. T. Findley, C. R. Gilbert, R. N. Lea, and J. D. Williams. 2004. Common and scientific names of fishes from the United States, Canada, and Mexico. American Fisheries Society, Special Publication 29, Bethesda, MD. 386 pp.
- Page, L. M., and B. M. Burr. 1991. A field guide to freshwater fishes: North America north of Mexico. Houghton Mifflin Company, Boston, MA. 432 pp.
- Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B. Scott. 1991. Common and scientific names of fishes from the United States and Canada. American Fisheries Society, Special Publishing 20. 183 pp.
- U.S. Fish and Wildlife Service. 1990. Endangered and threatened species recovery program: report to Congress. 406 pp.
- U.S. Fish and Wildlife Service. 1994. Yaqui fishes recovery plan. USDI Fish and Wildlife Service, Albuquerque, NM. iv + 48 pp.
- U.S. Fish and Wildlife Service. 1995. San Bernardino and Leslie Canyon National Wildlife Refuges. Comprehensive management plan, 1995-2015. Includes the environmental assessment under separate cover. Albuquerque, NM.

et al. 1986), as also documented for *Rhinichthys osculus* (John 1963). Gonads generally increase in size in February. Spawning first occurs in March when water temperatures reach approximately 19 C and proceeds until June (Minckley 1973, Anderson 1978), but mature ovaries have been noted in September (Minckley 1981). Propst et al. (1986) identified April as the peak breeding period and stated that spawning was completed by mid-May. Older females spawn earlier than younger females (Anderson 1978). Number of eggs per female ranges from 80 to 300 or more depending on female size. Anderson (1978) examined a sample of 29 females from the Gila River, 10 km south of Cliff, New Mexico; these ranged in size from 38 to 70 mm TL, with 88 to 246 mature ova per female. Anderson (1978) computed the relationship between female body size and fecundity as follows: number of ova = $-152.85 + 5.61 \text{ TL}$ ($r = 0.844$). Ovum diameter at spawning is near 1.5 mm. Age II females spawn at least twice per season, but most reproductive effort is by age I females (Barber et al. 1970, Anderson 1978, Sublette et al. 1990). Young first appear in April and May and reach 41 to 47 mm TL by November. Standard length (tip of snout to end of hypural plate) is related to total length by the following equation: $\text{SL} = 0.85 \text{ TL} - 0.12$, $r^2 = 0.99$, $n = 100$ (Marsh 1988). Total length averages 47 mm TL at the end of the first year, and 59 to 74 mm at the end of the second year. Sexual maturity occurs at about 40 mm in both sexes (Barber et al. 1970), and most become sexually mature in their second summer of life. Longevity typically is one to two years. Few live through their fourth summer and the largest individuals rarely exceed 70 mm (TL?) (Minckley 1973). Anderson (1978) reported an 81 mm TL female. Growth continues in the winter in Aravaipa Creek (Barber et al. 1970) but not in the cooler Gila River in New Mexico (Anderson 1978).

Ecology Comments: The spinedace is associated with a native fish fauna that includes roundtail chub (*Gila robusta*), loach minnow (*Tiaroga cobitis*), speckled dace (*Rhinichthys osculus*), longfin dace (*Agosia chrysogaster*), Sonora sucker (*Catostomus insignis*), and desert sucker (*Pantosteus clarki*). Historically, it was also associated with the woundfin (*Plagopterus argentissimus*), bonytail (*Gila elegans*), squawfish (*Ptychocheilus lucius*), and razorback sucker (*Xyrauchen texanus*), all now extirpated from the Gila River basin. Due to difficulties in tagging small fishes, movement patterns of spinedace adults are unquantified. Minckley (1981) showed that populations of spinedace in Aravaipa Creek increased following years of relatively high flow.

Habitat Type: Freshwater

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): CREEK, Low gradient, MEDIUM RIVER, Moderate gradient, Pool, Riffle

Special Habitat Factors: Benthic

Habitat Comments: Favors permanent, flowing, unpolluted water of low gradient streams having pool, riffle, run, and backwater areas; sand, gravel, and cobble substrates with low to moderate amounts of fine sediment and substrate embeddedness; abundant aquatic insects; natural hydrologic conditions, including recurrent flooding; few or no predatory or competitive non-native species present; a healthy riparian community; and moderate to high bank stability (USFWS, Federal Register, 8 March 1994; USFWS 1999). In larger rivers, spinedace often are found in the vicinity of tributary mouths. Adults favor slow to swift velocities (0-100 cm/sec) in shallow water (3-38 cm) with shear zones where rapid flow borders slower flow, areas of sheet flow at the upper ends of mid-channel sand/gravel bars, and eddies at downstream riffle edges. Juveniles favor slow to moderate flow (0-60 cm/sec) in shallow water (3-70 cm) with moderate amounts of instream cover; shallow stream margins and backwater areas, over silt, sand, or gravel bottoms, adjacent to pools. Periodic spates that scour and clean sands and gravels are essential to feeding and reproduction (Sublette et al. 1990). See Barber and Minckley (1966),

Anderson (1978), Propst and Bestgen (1986), Propst et al. (1986), and Sublette et al. (1990) for further details. May partition habitat with red shiner in areas where the two species co-occur (Rinne 1991). Spawns over shallow (less than 15 cm deep), sand-gravel-bottomed riffles where water flow is moderate (Minckley 1973, Sublette et al. 1990). Eggs develop in sand or gravel at spawning site (Sublette et al. 1990). Stability of the substrate is likely important during times of egg deposition and hatching (Minckley 1981). Larvae occur in areas of slow to moderate flow (0-30 cm/sec) in shallow water (3-30 cm) with abundant instream cover. Habitat utilization in the Cliff-Valley reach of the Gila River was studied by Propst et al. (1986). Juveniles (26-35 mm TL) were found to occupy an average depth of 16.1 cm and average current speed of 16.8 cm/s. Adults (>36 mm TL) in the same reach occupied an average depth of 19.3 cm and current of 49.1 cm/s. Spikedace occupied swifter waters in the warmer months of June to November than in the cooler months of December to May. Although habitat availability was not recorded, Propst et al. (1986) believed this to be a real shift. Sixty-percent of larval spikedace were captured over sand-dominated substrate, 18% over gravel, and 18% over cobble substrates. Juveniles were found over gravel substrates (46%), sand-dominated substrates (45%), and cobble substrates (9%). Adults were captured over gravel substrates (47%), cobble substrates (32%), and sand-dominated substrates (19%). Rinne and Kroeger (1988) observed spikedace in Aravaipa Creek at an average depth of 20 cm and current speed of 35 cm/s over gravel and pebble substrates (3-64 mm diameter). Schools of 10 or more fish were found in deeper and slower water than solitary fish. Seasonal differences were documented in use of depths but not currents. Spikedace collected in December, February, and August occupied shallower depths than those collected in April, May, and September. Rinne and Kroeger stated these differences showed no discernible pattern and were probably related to availability.

Adult Food Habits: Invertivore, Piscivore

Immature Food Habits: Invertivore, Piscivore

Food Comments: Diet is mainly aquatic and terrestrial insects, such as larval baetid ephemeropterans, and secondarily other larval ephemeropterans, hydropsychid trichopterans, and chironomid and simuliid dipterans (Anderson 1978, Schreiber and Minckley 1981, Barber and Minckley 1983, Abarca 1989). Schreiber and Minckley (1981) reported that up to approximately 30% of the diet was made up of emerging or adult insects. Also eats (seasonally) some fry of other fish species. In pools, eats mayflies; diet is more diverse in riffles and runs. Dipteran larvae are most important for small individuals, mayfly adults and nymphs for adults.

Phenology Comments: Feeding activity peaks in late afternoon and early evening (Barber and Minckley 1983). Larval cyprinids in the Gila River of New Mexico were found to be primarily diurnal drifters; 87% of cyprinid larvae collected were in noon or dusk drift samples (Bestgen et al. 1987). Additionally, a ratio of 6.5:1, nearshore vs. midstream, in captured larvae was found in noon samples, but a 1:1 ratio was found in dawn samples.

Stewardship Overview: Existing populations must be carefully monitored and protected by eliminating detrimental water and land use and exposure to non-native fishes. Research is needed to identify specific aspects of these practices that result in the demise of spikedace. Spikedace are not the only native fish threatened, endangered, or extirpated from the Gila River Basin. An ecological approach that addresses the habitat needs of all native fish species is necessary to protect remaining populations of native fishes. Degraded habitat should be reclaimed and enhanced, and spikedace should be reintroduced where chances for success are judged good.

Restoration Potential: Recovery potential is good only if adequate suitable habitat within the present or historical range is vigorously protected. Remaining populations continue to be threatened by habitat modification, predation by and competition with non-native fishes, and continued introduction and dispersal of non-native fishes. Reestablishment of the spikedace into its former range is problematic until the causes of its demise are identified and corrected.

Preserve Selection & Design Considerations: Preserves should be in areas of designated Critical Habitat (see Federal Register, 8 March 1994, p. 10906).

Management Requirements: The following management needs were identified by Marsh (1988): protect existing populations not infected by non-native fishes by building fish barriers or enhancing natural barriers (barrier design should not significantly alter stream flow and the potential impact on natural upstream and downstream movements of native fishes should be assessed; barrier design must be approved by appropriate agencies and the Desert Fishes Recovery Team); identify target areas amenable to management; determine and implement necessary habitat and landscape improvements (including removal of non-native fishes); reintroduce populations to selected streams within historic range, ensuring that genetic considerations are addressed (local stocks with affinities to those formerly occupying target streams should be utilized for reintroduction; e.g., Aravaipa Creek for the San Pedro, Gila River for the San Francisco; stockings should be done according to guidelines set up by the American Fisheries Society, Desert Fishes Recovery Team, and the U.S. Fish and Wildlife Service); assure closure of potential immigration routes to preclude reinvasion of non-native fishes. Proper management and maintenance of riparian zones are essential to native fish populations. Changes in the riparian zone can affect leaf fall and energy flow, flow, natural cover, temperature, and deposition of eroded materials (Baltz and Moyle 1984). Of five riparian systems studied in Arizona, only Aravaipa Creek, where cattle have been excluded since 1973, showed successful reproduction and dominance of the broadleaf riparian community (Rucks 1984). Cattle-browsing is a major factor in the replacement of a broadleaf riparian community by a riparian scrub community (Rucks 1984). A change from a broadleaf to scrub riparian community can change energy flow, tree-fall cover, amount of shade, and stream temperature. Fire would be a preferred method of watershed management when necessary. However, the choice of fire as a management tool must take into account the fuel levels present. A crown fire ("hot fire") can lead to increased runoff and result in the filling of riffles or other spawning areas. The effects of a crown fire and subsequent runoff were reversed in three years in the upper Carmel River, California (Hecht 1984). Removal of spawning areas for a period of two to three years would cause local extinctions of spinedace due to their short lifespan. If watershed management is necessary, controlled burns, frequent enough to prevent build-up of high fuel levels, set during non-spawning periods or periods of decreased spawning activity (autumn), should be employed.

Monitoring Requirements: Known populations should be monitored biannually in the spring during the breeding season and in late autumn to check recruitment. Data are needed to distinguish natural fluctuations in abundance from population declines due to human-caused perturbation. Reintroduced populations should be monitored for success or failure. An immediate monitoring program is needed for Aravaipa Creek, Arizona, due to the recent discovery of red shiner in September 1990 (Minckley 1990). Additionally, the black bullhead has increased in abundance in Aravaipa Creek and may prey on spinedace (Marsh 1990); this situation should be monitored. Standardized monitoring techniques should be adopted so that data will be comparable over locations and time. Techniques could be those recommended by the Desert Fishes Recovery Team. Techniques available for determination of absolute abundance for fishes include depletion sampling, mark-and-recapture, passive capture devices, and underwater censusing. These may be modified or others developed specifically for application to spinedace. Such techniques should be adjusted as dictated by experience, and uniformly applied. Minckley (1981) found that 6 to 10 passes with an electroshocking device, in an area blocked off with nets, were required to capture 99% of the spinedace. Natural units of a stream should be sampled, i.e. riffles, pools, runs and channels, rather than predetermined distances. Then the natural units could be measured and the results reported as densities per habitat type.

References:

- Abarca, F. J. 1989. Potential food habit overlap between spinedace (*Meda fulgida*) and red shiner (*Cyprinella lutrensis*). Arizona State University, Tempe. M.S. thesis.

- 1 • Anderson, R. M. 1978. The distribution and aspects of the life history of *Meda fulgida* in
2 New Mexico. New Mexico State University, Las Cruces. M.S. thesis. 62 pp.
- 3 • Arizona Game and Fish Commission. 1988. Threatened native wildlife in Arizona.
4 Arizona Game and Fish Department Publication, Phoenix, AZ. 32 pp.
- 5 • Baltz, D. M., and P. B. Moyle. 1984. The influence of riparian vegetation on steam fish
6 communities of California. Pp. 183-7 in R. E. Warner and K. M. Hendrix (eds.). California
7 riparian systems. University of California Press, Berkeley, CA.
- 8 • Barber, W. E., D. C. Williams, and W. L. Minckley. 1970. Biology of the Gila spikedace,
9 *Meda fulgida*, in Arizona. Copeia 1970:9-18.
- 10 • Barber, W. E., and W. L. Minckley. 1966. Fishes of Aravaipa Creek, Graham and Pinal
11 Counties, Arizona. Southwestern Nat. 11:313-24.
- 12 • Barber, W. E., and W. L. Minckley. 1983. Feeding ecology of a southwestern cyprinid
13 fish, the spikedace, *Meda fulgida* Girard. Southwestern Naturalist 28:33-40.
- 14 • Bestgen, K. R. 1985. Results of identification of collections of larval fish made in the
15 upper Salt and Gila Rivers, AZ. Report to U.S. Fish and Wildlife Service, September 10,
16 1985.
- 17 • Bestgen, K. R., D. L. Propst, and C. W. Painter. 1987. Transport Ecology of larval fishes
18 in the Gila River, New Mexico. Proc. Desert Fishes Council 16-18:174.
- 19 • Bestgen, K. R., and D. L. Propst. 1987. Red shiner vs. native fishes: replacement or
20 displacement? Proc. Desert Fishes Council 18:209.
- 21 • Bureau of Land Management. 1989. Fisheries habitat management on public land.
22 Washington, D.C.
- 23 • Douglas, M. E., P. C. Marsh, and W. L. Minckley. 1994. Indigenous fishes of western
24 North America and the hypothesis of competitive displacement: *Meda fulgida*
25 (Cyprinidae) as a case study. Copeia 1994:9-19.
- 26 • Gilbert, C. H., and N. B. Scofield. 1898. Notes on a collection of fishes from the Colorado
27 Basin in Arizona. Proc. U.S. Nat. Mus. 20:487-99.
- 28 • Girard, C. 1857. Researches upon the cyprinoid fishes inhabiting the fresh waters of the
29 United States of America, west of the Mississippi Valley, from specimens in the museum
30 of the Smithsonian Institution. Proc. Acad. Nat. Sci. Philadelphia. 8:165-213.
- 31 • Hardy, T. B., B. Bartz, and W. Carter. 1990. Instream flow recommendations for the
32 fishes of Aravaipa Creek, Arizona. Submitted to the Nature Conservancy. Twelve-Nine,
33 Inc., Logan, UT.
- 34 • Hecht, B. 1984. Sequential changes in bed habitat conditions in the upper Carmel River
35 following the Marble-Cone fire of August, 1977. Pp. 134-41 in R. E. Warner and K. M.
36 Hendrix (eds.). California riparian systems. University of California Press, Berkeley, CA.
- 37 • Hubbs, C. L. 1954. Establishment of a forage fish, the red shiner (*Notropis lutrensis*), in
38 the lower Colorado River system. California Fish and Game 40:287-94.
- 39 • John, K. R. 1963. The effect of torrential rains on the reproductive cycle of *Rhinichthys*
40 *osculus* in the Chiricahua Mountains, Arizona. Copeia 1963:286-91.
- 41 • Lee, D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer,
42 Jr. 1980. Atlas of North American Freshwater Fishes. North Carolina State Museum of
43 Natural History. 867 pp.
- 44 • Marsh, P. C. 1988. Spikedace (*Meda fulgida*) recovery plan draft. Center for
45 Environmental Studies, Arizona State University, prepared for Region II, U.S. Fish and
46 Wildlife Service, Albuquerque, NM. 55 pp.

- 1 • Marsh, P. C., F. J. Abarca, M. E. Douglas, and W. L. Minckley. 1989. Spikedace (*Meda*
2 *fulgida*) and loach minnow (*Tiaroga cobitis*) relative to introduced red shiner (*Cyprinella*
3 *lutrensis*). Final report to Arizona Game and Fish Dept., Phoenix, AZ.
- 4 • Matthews, J.R. and C.J. Moseley (eds.). 1990. The Official World Wildlife Fund Guide to
5 Endangered Species of North America. Volume 1. Plants, Mammals. xxiii + pp 1-560 +
6 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians,
7 Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180.
8 Beacham Publications, Inc., Washington, D.C.
- 9 • Meffe, G. K., and W. L. Minckley. 1987. Persistence and stability of fish and invertebrate
10 assemblages in a repeatedly disturbed Sonoran desert stream. American Midland
11 Naturalist 117:177-91.
- 12 • Miller, R. R. 1961. Man and the changing fish fauna of the American Southwest. Paper
13 Michigan Academy Science, Arts, and Letters 46:365-404.
- 14 • Miller, R. R., and C. L. Hubbs. 1960. The spiny-rayed cyprinid fishes (Plagopterini) of the
15 Colorado River system. University of Michigan Museum of Zoology Miscellaneous
16 Publication 115:1-39.
- 17 • Minckley, W. L. 1973. Fishes of Arizona. Arizona Game and Fish Department, Phoenix,
18 AZ.
- 19 • Minckley, W. L. 1981. Ecological studies of Aravaipa Creek, central Arizona, relative to
20 past, present, and future uses. Final Report. U.S. Bureau of Land Management, Safford,
21 Arizona. Contract Number YA-512-CT6-98. 362 pp.
- 22 • Minckley, W. L. 1985. Native fishes and natural aquatic habitats in U.S. Fish and Wildlife
23 Service Region II west of the Continental Divide. Report to U.S. Fish and Wildlife Service,
24 Albuquerque, NM. 158 pp.
- 25 • Minckley, W. L., and G. K. Meffe. 1987. Differential selection by flooding in stream-fish
26 communities of the arid American southwest. Pp. 93-104 in W. J. Matthews and D. E.
27 Heins (eds.). The zoogeography of North American freshwater fishes. John Wiley and
28 Sons, Incorporated, NY.
- 29 • Minckley, W. L., and J. E. Deacon. 1968. Southwestern fishes and the enigma of
30 "endangered species." Science 159:1424-32.
- 31 • Nelson, J. S., E. J. Crossman, H. Espinosa-Perez, L. T. Findley, C. R. Gilbert, R. N. Lea,
32 and J. D. Williams. 2004. Common and scientific names of fishes from the United States,
33 Canada, and Mexico. American Fisheries Society, Special Publication 29, Bethesda, MD.
34 386 pp.
- 35 • New Mexico State Game Commission. 1985. Regulation Number 624, as amended 28
36 March 1985. New Mexico Department of Game and Fish, Santa Fe, NM.
- 37 • Page, L. M., and B. M. Burr. 1991. A field guide to freshwater fishes: North America north
38 of Mexico. Houghton Mifflin Company, Boston, MA. 432 pp.
- 39 • Propst, D. 1986. Distribution, status and biology of the spikedace in the Gila River Basin,
40 New Mexico. New Mexico Department of Game and Fish, Santa Fe.
- 41 • Propst, D. L., K. R. Bestgen, and C. W. Painter. 1986. Distribution, status, and biology of
42 the spikedace (*Meda fulgida*) in New Mexico. Endangered Species Report, No. 15, U.S.
43 Fish and Wildlife Service, Albuquerque, NM. 93 pp.
- 44 • Propst, D. L., and K. R. Bestgen. 1986. Microhabitat utilization patterns of spikedace in
45 New Mexico. Proc. Desert Fishes Council 18 (1986).

- 1 • Rinne, J. N. 1991. Habitat use by spikedace, *Meda fulgida* (Pisces: Cyprinidae) in
2 southwestern streams with reference to probable habitat competition by red shiner,
3 *Notropis lutrensis* (Pisces: Cyprinidae). Southwestern Naturalist 36:7-13.
- 4 • Rinne, J. N., and E. Kroeger. 1988. Physical habitat use by spikedace, *Meda fulgida* in
5 Aravaipa Creek, Arizona. Proc. Western Association of Fish and Wildlife Agencies
6 68:282-90.
- 7 • Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B.
8 Scott. 1991. Common and scientific names of fishes from the United States and Canada.
9 American Fisheries Society, Special Publishing 20. 183 pp.
- 10 • Rucks, M. G. 1984. Composition and trend of riparian vegetation on five perennial
11 streams in southeastern Arizona. Pp. 97-107 in R. E. Warner and K. M. Hendrix (eds.).
12 California riparian systems. University of California Press, Berkeley, CA.
- 13 • Schreiber, D. C., and W. L. Minckley. 1981. Feeding interrelationships of native fishes in
14 a Sonoran Desert stream. Great Basin Naturalist 41:409-26.
- 15 • Sublette, J. E., M. D Hatch, and M. Sublette. 1990. The fishes of New Mexico. University
16 New Mexico Press, Albuquerque, NM. 393 pp.
- 17 • U.S. Fish and Wildlife Service (USFWS). 10 December 1999. Proposed designation of
18 critical habitat for the spikedace and the loach minnow. Federal Register 64(237):69323-
19 69355.
- 20 • U.S. Fish and Wildlife Service (USFWS). 1986. Determination of threatened status for the
21 spikedace. Federal Register 51(126):23769-80.
- 22 • U.S. Fish and Wildlife Service (USFWS). 1990. Endangered and threatened species
23 recovery program: report to Congress. 406 pp.
- 24 • U.S. Fish and Wildlife Service (USFWS). 1985. Proposal to determine the spikedace to
25 be a threatened species and to determine its critical habitat. Federal Register 50:25390-
26 25398.
- 27 • U.S. Fish and Wildlife Service (USFWS). 1986. Determination of threatened status for the
28 spikedace. Federal Register 51:23769-81.
- 29 • Winn, H. E., and R. R. Miller. 1954. Native postlarval fishes of the lower Colorado River
30 basin, with a key to their identification. California Fish and Game 40:273-85. 4 plates.
- 31

Gila Topminnow (*Poeciliopsis occidentalis occidentalis*)

The Gila topminnow is designated as a federally Endangered species.

Historic Range: Native range: Gila River system in Arizona and extreme western New Mexico; Rios de la Concepcion and Sonora, Sonora, Mexico. Currently occurs in the Gila river drainage, Arizona, particularly in the upper Santa Cruz River, Sonoita and Cienega creeks, and the middle Gila River; and in the Rio Sonora, Rio de la Concepcion, and Santa Cruz River (Weedman 1998). Extirpated in New Mexico; later reintroduced in New Mexico into a small pond on the Red Rock Wildlife Area, north of Lordsburg, in 1989; there is some question as to whether the fishes will be able to survive the cold winters of that area (Sublette et al. 1990).

Basic Description: A small fish (topminnow).

Reproduction Comments: In some areas reproduces throughout the year; in other areas breeding prolonged throughout spring and summer. Interval between broods apparently about 24 to 28 days. Depending on their size adults produce 1-15 young/brood (Minckley 1973). Life span apparently is about one year.

Habitat Type: Freshwater

Non-Migrant: Yes

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): CREEK, Low gradient, MEDIUM RIVER, Moderate gradient, Pool, SPRING/SPRING BROOK

Palustrine Habitat(s): HERBACEOUS WETLAND

Habitat Comments: Lowland and some upland streams of desert and grasslands, and margins of large, lowland rivers. Typical inhabitant of vegetated springs, brooks, and margins and backwaters of larger bodies of water (Lee et al. 1980). Prefers shallow, warm, fairly quiet waters but also can be found in moderate currents and depths up to 1 m; permanent and intermittent streams, marshes; preferred habitat has dense mats of algae and debris (usually along stream margins or below riffles) and sandy substrate sometimes covered with mud and debris (Matthews and Moseley 1990).

Adult Food Habits: Herbivore, Invertivore

Immature Food Habits: Herbivore, Invertivore

Food Comments: Eats detritus and algae; also feeds opportunistically on aquatic invertebrates (Lee et al. 1980).

Length: 3 cm

Restoration Potential: In Arizona, attempts to eradicate *Gambusia* from sites with natural topminnow populations have been unsuccessful (*Gambusia* reinvaded); fencing to protect habitat from livestock resulted in vegetation encroachment and extirpation of the topminnow at another site.

Management Requirements: Minckley (1999) emphasized the need for protection of existing populations, establishment of populations in artificial refugia, and elimination, exclusion, or

management against introduced piscivores. See Marsh and Minckley (1990) for recommendations on methods for eradicating *Gambusia* (poison fish, reintroduce topminnow, frequently monitor system) and removing vegetation (cattle grazing may be best method). See Minckley et al. (1991) for detailed information on management and reintroduction efforts. See also Hendrickson and Brooks (1991) for information on transplantation efforts. High levels of heterozygosity, which correlate with enhanced survivorship and fecundity, make the Sharp Spring population (Arizona) the best choice for source of fishes for the restocking effort in the Gila River system (Quattro and Vrijenhoek 1989). The captive stock at Dexter National Fish Hatchery was replaced by fishes from Sharp Spring in the mid-1980s (Minckley and Deacon 1991). Based on patterns of molecular variation, Parker et al. (1999) recommended that each of the four watersheds in which subspecies *occidentalis* is still naturally extant be managed and conserved separately (see also Sheffer et al. 1997). Weedman (1998) cited the following needed actions: protect remaining natural and long-lived established populations; reestablish and protect populations throughout historical range; monitor populations and their habitats; develop and implement genetic protocol for managing populations; study life history, genetics, ecology, habitat, and interactions with non-native aquatic species; inform and educate the public and resource managers.

References:

- Constantz, G. D. 1979. Life history patterns of a livebearing fish in contrasting environments. *Oecologia* (Berl.) 40:189-201.
- Echelle, A. A. 1991. Conservation genetics and genic diversity in freshwater fishes of western North America. Pages 141-153 in W. L. Minckley and J. E. Deacon, eds. *Battle against extinction: native fish management in the American West*. Univ. Arizona Press, Tucson, AZ.
- Hedrick, P. W., K. M. Parker, and R. N. Lee. 2001. Using microsatellite and MHC variation to identify species, ESUs, and MUs in the endangered Sonoran topminnow. *Molecular Ecology* 10:1399-1412.
- Hendrickson, D. A., and J. E. Brooks. 1991. Transplanting short-lived fishes in North American deserts: review, assessment, and recommendations. Pages 283-98 in W. L. Minckley and J. E. Deacon (editors). *Battle Against Extinction: Native Fish Management in the American West*. University of Arizona Press, Tucson, AZ.
- Lee, D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer, Jr. 1980. *Atlas of North American Freshwater Fishes*. North Carolina State Museum of Natural History. 867 pp.
- Marsh, P. C., and W. L. Minckley. 1990. Management of endangered Sonoran topminnow at Bylas Springs, Arizona: description, critique, and recommendations. *Great Basin Naturalist* 50:265-272.
- Matthews, J.R. and C.J. Moseley (eds.). 1990. *The Official World Wildlife Fund Guide to Endangered Species of North America*. Volume 1. Plants, Mammals. xxiii + pp 1-560 + 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians, Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180. Beacham Publications, Inc., Washington, D.C.
- Meffe, G. K. 1985. Predation and species replacement in American southwestern fishes: a case study. *Southwestern Naturalist* 30:173-187.
- Meffe, G. K., and F. F. Snelson, Jr., editors. 1989. *Ecology and evolution of livebearing fishes (Poeciliidae)*. Prentice Hall, Englewood Cliffs, NJ. 453 pp.
- Meffe, G. K., and R. C. Vrijenhoek. 1988. Conservation genetics in the management of desert fishes. *Conservation Biology* 2:157-169.

- 1 • Minckley, W. L. 1973. Fishes of Arizona. Arizona Game and Fish Department, Phoenix,
2 AZ.
- 3 • Minckley, W. L. 1999. Ecological review and management recommendations for recovery
4 of the endangered Gila topminnow. Great Basin Naturalist 59:230-244.
- 5 • Minckley, W. L., G. K. Meffe, and D. L. Soltz. 1991. Conservation and management of
6 short-lived fishes: the cyprinodontoids. Pages 247-82 in W. L. Minckley and J. E. Deacon
7 (editors). Battle Against Extinction: Native Fish Management in the American West.
8 University of Arizona Press, Tucson, AZ.
- 9 • Minckley, W. L., J. N. Rinne, and J. E. Johnson. 1977. Status of the Gila topminnow and
10 its co-occurrence with mosquitofish. USDA Forest Service Res. Pap. RM-198, Rocky
11 Mountain Forest and Range Expt. Sta., Fort Collins. 8 pp.
- 12 • Minckley, W. L., and J. E. Deacon. 1991. Battle Against Extinction: Native Fish
13 Management in the American West. University of Arizona Press, Tucson. xviii + 517 pp.
- 14 • Nelson, J. S., E. J. Crossman, H. Espinosa-Perez, L. T. Findley, C. R. Gilbert, R. N. Lea,
15 and J. D. Williams. 2004. Common and scientific names of fishes from the United States,
16 Canada, and Mexico. American Fisheries Society, Special Publication 29, Bethesda, MD.
17 386 pp.
- 18 • Page, L. M., and B. M. Burr. 1991. A field guide to freshwater fishes: North America north
19 of Mexico. Houghton Mifflin Company, Boston, MA. 432 pp.
- 20 • Parker, K. M., R. J. Sheffer, and P. W. Hedrick. 1999. Molecular variation and
21 evolutionarily significant units in the endangered Gila topminnow. Conservation Biology
22 13:108-116.
- 23 • Quattro, J. M., and R. C. Vrijenhoek. 1989. Fitness differences among remnant
24 populations of the endangered Sonoran topminnow. Science 245:976-978.
- 25 • Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B.
26 Scott. 1991. Common and scientific names of fishes from the United States and Canada.
27 American Fisheries Society, Special Publishing 20. 183 pp.
- 28 • Sheffer, R. J., P. W. Hedrick, W. L. Minckley, and A. L. Velasco. 1997. Fitness of the
29 endangered Gila topminnow. Conservation Biology 11:162-171.
- 30 • Simons, L. H., D. A. Hendrickson, and D. Papoulias. 1989. Recovery of the Gila
31 topminnow: a success story? Conservation Biology 3:11-15.
- 32 • Sublette, J. E., M. D Hatch, and M. Sublette. 1990. The fishes of New Mexico. University
33 New Mexico Press, Albuquerque, NM. 393 pp.
- 34 • U.S. Fish and Wildlife Service (USFWS). 1967. Native fish and wildlife: endangered
35 species. Federal Register 32(48):4001.
- 36 • U.S. Fish and Wildlife Service (USFWS). 1990. Endangered and threatened species
37 recovery program: report to Congress. 406 pp.
- 38 • Weedman, D. A. 1998. Gila topminnow, *Poeciliopsis occidentalis occidentalis*, revised
39 recovery plan. U.S. Fish and Wildlife Service, Albuquerque, NM.
- 40 • Williams, J. E., J. E. Johnson, D. A. Hendrickson, S. Contreras-Balderas, J. D. Williams,
41 M. Navarro-Mendoza, D. E. McAllister, and J. E. Deacon. 1989. Fishes of North America
42 endangered, threatened, or of special concern: 1989. Fisheries 14:2-20.

Yaqui Topminnow (*Poeciliopsis occidentalis sonoriensis*)

The Yaqui topminnow has an implied federally Endangered species status because it is a subspecies of the federally endangered Gila topminnow (*Poeciliopsis occidentalis*).

Historic Range: Native range: Rio Yaqui basin in Arizona (b) (7)(E) and several tributaries of the Rio Yaqui in Sonora, Mexico (Minckley et al. 1991, draft recovery plan). Hendrickson et al. (1980) found this fish to be widely distributed below elevations of 1300 m in the Rio Yaqui basin. Presently occurs in the U.S. at several locations within the (b) (7)(E) (draft recovery plan).

Basic Description: A fish (topminnow) that reaches a maximum length of about 6 cm.

Reproduction Comments: Breeds year-round where winter temperatures are ameliorated by spring flows, breeds mainly April-October otherwise; adult females produce broods of up to 20+ young at intervals of about 20 days; few live more than 1 year (USFWS 1994).

Habitat Type: Freshwater

Non-Migrant: Yes

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): CREEK, Low gradient, MEDIUM RIVER, Moderate gradient, Pool, SPRING/SPRING BROOK

Palustrine Habitat(s): HERBACEOUS WETLAND

Habitat Comments: Lowland and some upland streams of desert and grasslands, and margins of large, lowland rivers. Typical inhabitant of vegetated springs, brooks, and margins and backwaters of larger bodies of water (Lee et al. 1980). Prefers shallow, warm, fairly quiet waters but also can be found in moderate currents and depths up to 1 m; permanent and intermittent streams, marshes; preferred habitat has dense mats of algae and debris (usually along stream margins or below riffles) and sandy substrate sometimes covered with mud and debris (Matthews and Moseley 1990). On the (b) (7)(E), occurs in shallows of artesian well outflows, ponds, and pool margins (draft recovery plan).

Adult Food Habits: Herbivore, Invertivore

Immature Food Habits: Herbivore, Invertivore

Food Comments: Detritus and algae; also feeds opportunistically on aquatic invertebrates such as amphipods and insect larvae (Minckley 1973, Lee et al. 1980).

Length: 3 cm

Management Requirements: Securing habitat and water sources are major management needs (USFWS 1994).

References:

- 1 • Hendrick, P. W., K. M. Parker, and R. N. Lee. 2001. Using microsatellite and MHC
2 variation to identify species, ESUs, and MUs in the endangered Sonoran topminnow.
3 *Molecular Ecology* 10:1399-1412.
- 4 • Lee, D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer,
5 Jr. 1980. *Atlas of North American Freshwater Fishes*. North Carolina State Museum of
6 Natural History. 867 pp.
- 7 • Matthews, J.R. and C.J. Moseley (eds.). 1990. *The Official World Wildlife Fund Guide to*
8 *Endangered Species of North America*. Volume 1. Plants, Mammals. xxiii + pp 1-560 +
9 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians,
10 Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180.
11 Beacham Publications, Inc., Washington, D.C.
- 12 • Minckley, W. L. 1973. *Fishes of Arizona*. Arizona Game and Fish Department, Phoenix,
13 AZ.
- 14 • Minckley, W. L. 1999. Ecological review and management recommendations for recovery
15 of the endangered Gila topminnow. *Great Basin Naturalist* 59:230-244.
- 16 • Minckley, W. L., G. K. Meffe, and D. L. Soltz. 1991. Conservation and management of
17 short-lived fishes: the cyprinodontoids. Pages 247-82 in W. L. Minckley and J. E. Deacon
18 (editors). *Battle Against Extinction: Native Fish Management in the American West*.
19 University of Arizona Press, Tucson, AZ.
- 20 • Nelson, J. S., E. J. Crossman, H. Espinosa-Perez, L. T. Findley, C. R. Gilbert, R. N. Lea,
21 and J. D. Williams. 2004. *Common and scientific names of fishes from the United States,*
22 *Canada, and Mexico*. American Fisheries Society, Special Publication 29, Bethesda, MD.
23 386 pp.
- 24 • Page, L. M., and B. M. Burr. 1991. *A field guide to freshwater fishes: North America north*
25 *of Mexico*. Houghton Mifflin Company, Boston, MA. 432 pp.
- 26 • Quattro, J. M., P. L. Leberg, M. E. Douglas, and R. C. Vrijenhoek. 1996. Molecular
27 evidence for a unique evolutionary lineage of endangered Sonoran Desert fish (genus
28 *Poeciliopsis*). *Conservation Biology* 10:128-135.
- 29 • U.S. Fish and Wildlife Service (USFWS). 1994. Yaqui fishes recovery plan. USDI Fish
30 and Wildlife Service, Albuquerque, NM. iv + 48 pp.
- 31 • U.S. Fish and Wildlife Service (USFWS). 1995. San Bernardino and Leslie Canyon
32 National Wildlife Refuges. Comprehensive management plan, 1995-2015. Includes the
33 environmental assessment under separate cover. Albuquerque, NM.

34

Loach Minnow (*Tiaroga cobitis*)

The loach minnow was designated as a federally Threatened species on October 28, 1986.

Historic Range: Once locally common throughout much of the Verde, Salt, San Pedro, San Francisco, and Gila (upstream from Phoenix) river systems, Arizona, New Mexico, and Sonora, occupying suitable habitat in both the mainstreams and perennial tributaries, at elevations up to about 2200 m. Extirpated throughout much of its former range in Arizona. Occurred historically in the San Pedro River, Sonora, Mexico, but habitat there has been largely destroyed by diversion of water for agriculture. Now restricted to about 645 km of stream in portions of the upper Gila River (Grant, Catron, and Hidalgo counties, New Mexico), the San Francisco and Tularosa rivers and their tributaries Negrito and Whitewater creeks (Catron County, New Mexico), the Blue River and its tributaries Dry Blue, Campbell Blue, Little Blue, Pace, and Frieborn creeks (Greenlee County, Arizona, and Catron County, New Mexico), Aravaipa Creek and its tributaries Turkey and Deer creeks (Graham and Pinal counties, Arizona), Eagle Creek (Graham and Greenlee counties, Arizona), the White River (Apache, Gila, and Navajo counties, Arizona), and the Black River (Apache and Greenlee counties, Arizona) (USFWS 1999). Common only in Aravaipa Creek, the Blue River, and limited portions of the San Francisco, upper Gila, and Tularosa rivers in New Mexico (USFWS 1999). Marsh et al. (2003) reported a new record from North Fork of East Fork Black River, Arizona, and a rediscovered population in Eagle Creek, Arizona; the species recorded in the latter location in 1950 and the mid-1990s but has not been seen there since 1997.

Basic Description: A small fish (minnow), up to 6 cm long.

Reproduction Comments: In New Mexico, most spawners were in their second summer (Propst and Bestgen 1991). Spawning occurs in Arizona mainly March-June, with some breeding December-February; nests with eggs found also in September (Vives and Minckley 1990). Spring (e.g., April) spawning recorded in New Mexico. Female produces between 250 to 1,200 ova (Minckley 1973). Eggs hatch in about 6 days at 21 C. Male may provide some care to developing eggs (female also?) (Vives and Minckley 1990).

Habitat Type: Freshwater

Non-Migrant: Yes

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): CREEK, High gradient, MEDIUM RIVER, Moderate gradient, Riffle

Special Habitat Factors: Benthic

Habitat Comments: Lives on bottom in permanent, flowing, unpolluted creeks and small to medium rivers of low to moderate gradient, low amounts of fine sediment and substrate embeddedness, abundant aquatic insects, and a healthy, intact riparian community with moderate to high bank stability; typically on turbulent riffles, sometimes in association with filamentous algae; habitat resembles that of many eastern darters (Percidae) (Lee et al. 1980). Obligate riffle-dweller, occurs in shallow (<20 cm) water over gravel/ cobble substrate (Rinne 1989, Propst and Bestgen 1991) or in interstices between rocks, often in association with eddying currents (Sublette et al. 1990). Adults inhabit moderate to swift (15-100 cm/sec), shallow (3-40 cm) water with gravel, cobble, and rubble substrates; juvenile habitat is similar but includes also sand substrates (Federal Register, 8 March 1994). Persists mainly in streams having relatively natural flow regimes and a predominance of native species (Propst and Bestgen 1991). Recurrent flooding is important in keeping substrate free of sediments and in helping this species

maintain a competitive edge over invading non-native fishes. Eggs are laid in cavities under flattened cobble (or uncemented cobble and rubble) in slow to swift (3-85 cm/sec), shallow (3-30 cm) water; eggs adhere to under surface (Sublette et al. 1990, Vives and Minckley 1990); males guard cavities and eggs. Larvae apparently use low velocity nursery areas: 0-30 cm/sec, 3-30 cm deep, with sand, gravel, and cobble substrates and abundant instream cover (Sublette et al. 1990; Propst and Bestgen 1991; Federal Register, 8 March 1994).

Adult Food Habits: Invertivore

Immature Food Habits: Invertivore

Food Comments: Restricted diet; feeds opportunistically on riffle-inhabiting insect larvae (e.g., simuliid dipterans and mayflies). Immatures feed principally on chironomids, adults eat various benthic insects (dipterans, mayflies, stoneflies, caddisflies) (Sublette et al. 1990, Propst and Bestgen 1991).

Length: 6 cm

References:

- Coburn, M. M., and T. M. Cavender. 1992. Interrelationships of North American cyprinid fishes. Pages 328-373 in R. L. Mayden, editor. Systematics, historical ecology, and North American freshwater fishes. Stanford Univ. Press, Stanford, CA. xxvi + 969 pp.
- Johnston, C. E., and L. M. Page. 1992. The evolution of complex reproductive strategies in North American minnows (Cyprinidae). Pages 600-621 in R. L. Mayden, editor. Systematics, historical ecology, and North American freshwater fishes. Stanford Univ. Press, Stanford, CA. xxvi + 969 pp.
- Lee, D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer, Jr. 1980. Atlas of North American Freshwater Fishes. North Carolina State Museum of Natural History. 867 pp.
- Marsh, P. C., B. E. Bagley, G. W. Knowles, G. Schiffmiller, and P. A. Sowka. 2003. New and rediscovered populations of loach minnow, *Tiaroga cobitis* (Cyprinidae), in Arizona. Southwestern Naturalist 48:666-669.
- Matthews, J.R. and C.J. Moseley (eds.). 1990. The Official World Wildlife Fund Guide to Endangered Species of North America. Volume 1. Plants, Mammals. xxiii + pp 1-560 + 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians, Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180. Beacham Publications, Inc., Washington, D.C.
- Minckley, W. L. 1973. Fishes of Arizona. Arizona Game and Fish Department, Phoenix, AZ.
- Nelson, J. S., E. J. Crossman, H. Espinosa-Perez, L. T. Findley, C. R. Gilbert, R. N. Lea, and J. D. Williams. 2004. Common and scientific names of fishes from the United States, Canada, and Mexico. American Fisheries Society, Special Publication 29, Bethesda, MD. 386 pp.
- Page, L. M., and B. M. Burr. 1991. A field guide to freshwater fishes: North America north of Mexico. Houghton Mifflin Company, Boston, MA. 432 pp.
- Propst, D. L. 1986. Distribution, status, and biology of the loach minnow in the Gila River basin. New Mexico Department of Game and Fish, Santa Fe.
- Propst, D. L., and K. R. Bestgen. 1991. Habitat and biology of the loach minnow, *Tiaroga cobitis*, in New Mexico. Copeia 1991:29-38.

- 1 • Rinne, J. N. 1989. Physical habitat use by loach minnow, *Tiaroga cobitus* (Pisces:
2 Cyprinidae), in southwestern desert streams. Southwestern Naturalist 34:109-17.
- 3 • Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B.
4 Scott. 1991. Common and scientific names of fishes from the United States and Canada.
5 American Fisheries Society, Special Publishing 20. 183 pp.
- 6 • Sublette, J. E., M. D Hatch, and M. Sublette. 1990. The fishes of New Mexico. University
7 New Mexico Press, Albuquerque, NM. 393 pp.
- 8 • U.S. Fish and Wildlife Service (USFWS). 10 December 1999. Proposed designation of
9 critical habitat for the spikedace and the loach minnow. Federal Register 64(237):69323-
10 69355.
- 11 • U.S. Fish and Wildlife Service (USFWS). 1990. Endangered and threatened species
12 recovery program: report to Congress. 406 pp. Vives, S. P., and W. L. Minckley. 1990.
13 Autumn spawning and other reproductive notes on loach minnow, a threatened cyprinid
14 fish of the American southwest. Southwestern Naturalist 35:451-454.
- 15 • Williams, J. E., J. E. Johnson, D. A. Hendrickson, S. Contreras-Balderas, J. D. Williams,
16 M. Navarro-Mendoza, D. E. McAllister, and J. E. Deacon. 1989. Fishes of North America
17 endangered, threatened, or of special concern: 1989. Fisheries 14:2-20.
- 18 • Woodman, D. A. 1992. Systematic relationships within the cyprinid genus *Rhinichthys*.
19 Pages 374-391 in R. L. Mayden, editor. Systematics, historical ecology, and North
20 American freshwater fishes. Stanford Univ. Press, Stanford, CA. xxvi + 969 pp.
- 21

Huachuca Springsnail (*Pyrgulopsis thompsoni*)

The Huachuca springsnail was designated as a Federal Candidate species on September 12, 2006.

Historic Range: Range is the upper portion of the Santa Cruz and San Pedro River basins in Arizona and Sonora, Mexico. Originally it covered only six sites in Santa Cruz County, Arizona and Sonora, Mexico. These sites were: Cottonwood Springs, Monkey Spring, Canelo Hills Cienega, Sheehy Spring, Peterson Ranch Springs, and Ojo Caliente, (Hershler and Landye, 1988). Since that time, Landye in 1992 examined sixteen springs on (b) (7)(E) and found occurrences at nine springs. The nine additional sites are Upper Garden Canyon Spring, Lower Garden Canyon Spring, McClure Spring, Broken Pipe Spring, Cave Spring, Sawmill Canyon Spring, Upper Water Supply Spring, Lower Water Supply Spring, Blacktail Spring (Landye, pers. comm.). An additional site in Mexico was reported at Cienega Los Fresnos (Stefferd, pers. comm.).

Basic Description: a snail

General Description: This is considered a medium to large species relative to other Hydrobiidae snails, with a shell 1.7 to 3.2 mm long. The shell is ovate-conic with "3.25 to 5 moderately convex, slightly shouldered whorls". The aperture may be fused or separate from the body whorl. The pigmentation of the snout and anterior part of the foot tends from light to dark with the remaining portion and the head generally unpigmented. There appears to be some sexual dimorphism in two of four populations studied, in one case the males being larger than the females and vice versa in the other population. The identification is based upon characteristics of the reproductive organs. The penis which is considered moderate in size may be "squat to elongate". The ventral penial lobe surface has a glandular ridge, this is generally located at the tip of the lobe. The penial filament may be 35 to 103 per cent of the penis length and centered at 80 to 93% of the penis length. The whole of the penis exhibits a dark pigmentation. The testis and prostrate make up 37 to 54% and 7 to 8% of the body length, respectively. Between 55 and 85% of the bursa length is posterior to the albumen gland (Hershler and Landye 1988).

Ecology Comments: Little is known about the life history, biology or ecology of this small snail, but Landye (1993) suggested that it may be similar to another Hydrobiidae species, *P. morrisoni*, the Page Spring snail. The Page Spring snail experiences what appears to be a population crash in December and young appear in January.

Habitat Type: Freshwater

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): SPRING/SPRING BROOK

Palustrine Habitat(s): HERBACEOUS WETLAND

Habitat Comments: Habitat is restricted to springs and cienega wetland habitats. Within these habitats it is commonly found in shallow water on rocks around the spring sources.

Stewardship Overview: This snail occurs in cienegas and isolated springs in the upper Santa Cruz and San Pedro River drainages; a range-wide survey to determine the distribution is critically needed as is basic information on ecology, life-cycle, and population dynamics. Currently, the only management strategy is to maintain inhabited cienega and spring-fed wetland

habitat by (i) reducing the impacts of livestock on wetland vegetation, and ensuring bank stability and water quality; (ii) protecting the aquifer sources of these wetlands from groundwater pumping, water diversion, and pollution; and (iii) preventing erosion and incision of the stream channel through good land-use practices or construction of erosion-control structures.

Restoration Potential: Given the lack of knowledge about biology and ecology, including response to disturbance, recovery potential is unknown.

Preserve Selection & Design Considerations: Protection requires protection of wetland habitats, protection of the aquifer sources of these wetlands from groundwater pumping, maintaining channel stability upstream and downstream in the watershed (i.e., discouraging channel incision and erosion) and assuring high standards of water quality upstream. Within the site, protection requires maintenance of suitable firm (rocky) substrates, which seems to be a component of preferred habitat.

Management Requirements: With so little information, it is difficult to prescribe management directives. As a default, management should be targeted at maintaining the inhabited cienega and spring fed wetland habitat by (i) reducing the impacts of livestock on wetland vegetation, and ensuring bank stability and water quality; (ii) protecting the aquifer sources of these wetlands from groundwater pumping, water diversion, and pollution; and (iii) preventing erosion and incision of the stream channel through good land-use practices or construction of erosion-control structures.

Monitoring Requirements: It would be useful to assess the numbers of sites within the San Pedro and Santa Cruz River drainages. Based on the Landye 1992 survey, additional populations are likely to be found. Once a range-wide survey is completed, then one can begin to assess the nature and severity of threats. An important need is the development of a monitoring protocol to assess population size.

References:

- Arizona Game and Fish Department. 1992. Status Summary. Huachuca Springsnail (*Pyrgulopsis thomsoni*).
- Hendrickson, D. A. and W. L. Minckley. 1984. Cienegas--Vanishing climax communities of the American Southwest. *Desert Plants*. 6:141-75.
- Hershler, R. and J.J. Landye. 1988. Arizona Hydrobiidae (Prosobranchia: Rissoacea). *Smithsonian Contributions to Zoology*, 459: 1-58.
- Turgeon, D.D., J.F. Quinn, Jr., A.E. Bogan, E.V. Coan, F.G. Hochberg, W.G. Lyons, P.M. Mikkelsen, R.J. Neves, C.F.E. Roper, G. Rosenberg, B. Roth, A. Scheltema, F.G. Thompson, M. Vecchione, and J.D. Williams. 1998. Common and scientific names of aquatic invertebrates from the United States and Canada: Mollusks. 2nd Edition. American Fisheries Society Special Publication 26, Bethesda, MD: 526 pp.
- U.S. Fish and Wildlife Service (USFWS). 1997. Category and Listing Priority Forms.

Sonoran Tiger Salamander (*Ambystoma tigrinum stebbinsi*)

The Sonoran tiger salamander was designated as a federally Endangered species on January 6, 1997.

Historic Range: Santa Cruz and San Pedro river drainages, Santa Cruz and (b) (7)(E) counties, Arizona, including sites in the San Rafael Valley (SRV) and adjacent foothills of the (b) (7)(E) and (b) (7)(E). The range of the subspecies and its occupied and potentially occupied habitat is thought to extend from the crest of the (b) (7)(E) west to the crest of the (b) (7)(E), including the SRV and adjacent foothills from its origins in Sonora north to the Canelo Hills (USFWS 2002). Salamanders suspected of being Sonora tiger salamanders have been collected from Los Fresnos cienega in the School Canyon drainage approximately 3 km south of the border (Varela-Romero *et al.* 1992). Genetic testing showed that some SRV ponds contain salamanders with genetic characteristics similar to barred tiger salamanders. Salamanders with these "mavortium-like" sequences are more common on the outskirts of the SRV and ponds close to (b) (7)(E) Lake, which, because of prior use of imported waterdogs as fish bait, is where introduced barred tiger salamanders are expected to be found (Ziemba *et al.* 1998). Tiger salamanders have also been found in areas just outside the SRV, such as (b) (7)(E), (b) (7)(E), Copper Canyon, and (b) (7)(E). Of these localities, genetic testing has only been performed on salamanders from the Fort, and with the exception of one pond within a kilometer of the SRV, salamanders on the Fort appear to be barred tiger salamanders (Andrew Storfer, University of Florida, pers. comm.) (USFWS 2002).

Basic Description: A robust salamander.

General Description: Metamorphosed terrestrial Sonora tiger salamanders have a color pattern ranging from "a reticulate pattern with an irregular network of light coloration, often coupled with light spots, on a dark background color", to a pattern of large, well-defined light or yellow spots or transverse bars, some of which encroach on the dark venter (Jones *et al.* 1988). Metamorphosed Sonora tiger salamanders measure from about 45 to 150 mm snout to vent length (SVL). Branchiate adults are gray to olive on the dorsum, head, and tail, and off-white to yellow on the ventral side. They have three external gills on each side of their head, and measure between 65 and 165 mm SVL. Male and female adult salamanders can be distinguished by the presence of two black folds of tissue (cloacal folds) on the caudal side of a male's vent. Larvae are gray on the dorsum, head, and tail, with little pigment on the ventral surface. They have external gills and hatch without legs, but grow hind and fore-limbs early in development (USFWS 2002).

Reproduction Comments: Breeds as early as January or as late as early May; breeding after monsoon rains in July and August is rare (Synder, cited by USFWS 2002). Some larvae hatched in spring metamorphose into terrestrial form from late July to early September; other individuals become sexually mature in the larval form or overwinter as immature larvae (USFWS 2002).

Non-Migrant: Yes

Locally Migrant: Yes

Long Distance Migrant: No

Mobility and Migration Comments: Movement patterns not thoroughly documented; most likely stay within a few hundred meters of their natal pond, but some may move 1.5-2.0 km or more between breeding and nonbreeding habitats or between ponds (see USFWS 2002).

Riverine Habitat(s): SPRING/SPRING BROOK

Palustrine Habitat(s): HERBACEOUS WETLAND, TEMPORARY POOL

1 **Special Habitat Factors:** Benthic, Burrowing in or using soil, Fallen log/debris

2 **Habitat Comments:** Cienegas, impounded cienegas, springs, livestock tanks; breeds mainly in
3 cattle ponds or tanks (USFWS 2002). Adult, metamorphosed salamanders inhabit adjacent
4 grassland and oak woodland terrestrial habitat when not in ponds (USFWS 2002). Mammal
5 burrows or loosened soils outside the pond likely provide refugia for metamorphosed
6 salamanders in the terrestrial environment, enabling them to burrow underground to avoid
7 extreme environmental conditions (USFWS 2002).

8 **Adult Food Habits:** Carnivore, Invertivore

9 **Immature Food Habits:** Carnivore, Invertivore

10 **Stewardship Overview:** Recovery Criteria: The Sonora tiger salamander may be reclassified to
11 threatened status when approximately 90 percent of salamander's currently-occupied range and
12 approximately 90 percent of current breeding ponds are protected and maintained to prevent
13 habitat loss and degradation, predator introductions, barred tiger salamander introductions, and
14 collection of salamanders for bait. Scientifically credible monitoring over a five year period must
15 indicate that the number of Sonora tiger salamander populations is not in decline and that there
16 are no new factors that threaten the persistence of Sonora tiger salamanders (USFWS 2002).
17 The Sonora tiger salamander will be considered for delisting when quantitative criteria in terms of
18 number of breeding populations and amount, distribution, and type of available habitat are
19 defined and met. Criteria will be based on research, continued monitoring, and population
20 viability analysis. In addition, regulatory mechanisms and land management commitments must
21 be implemented that provide for adequate long-term protection of the Sonora tiger salamander
22 and its habitat. These commitments and mechanisms should address habitat maintenance and
23 protection, management of non-native predators, disease transmission, introduction and
24 collection of salamanders, interbreeding with non-native salamanders, and public education.
25 Finally, the Sonora tiger salamander must be unlikely to need protection under the Endangered
26 Species Act in the foreseeable future (USFWS 2002).

27 **Actions Needed (USFWS 2002):**

- 28 1. Maintain and enhance habitat where salamanders have been found, and create new
29 habitat, if deemed necessary.
- 30 2. Control non-native predators (fish, bullfrogs, and crayfish) by enforcing and enhancing
31 existing policies prohibiting the introduction and pond to pond transport of these taxa and
32 by removing populations of non-native fish, bullfrogs, and crayfish.
- 33 3. Control introduction, transport, and collection of tiger salamanders in the San Rafael
34 Valley by enforcing existing policies prohibiting these acts and by removing populations
35 of barred tiger salamanders.
- 36 4. Create and enforce policies to minimize frequency of die-offs.
- 37 5. Monitor salamander populations and their habitat on public and, if permitted, private land,
38 to observe threats as they arise and fulfill research objectives.
- 39 6. Conduct research to acquire demographic and dispersal information and develop a
40 population viability analysis, better understand salamander disease, conduct genetic
41 analyses, investigate reports of low pH, and determine distribution of crayfish and
42 methods of crayfish removal.
- 43 7. Develop public education and information programs.
- 44 8. Practice adaptive management.

1 **References:**

- 2 • Collins, J. T. 1990. Standard common and current scientific names for North American
3 amphibians and reptiles. 3rd ed. Society for the Study of Amphibians and Reptiles.
4 Herpetological Circular No. 19. 41 pp.
- 5 • Jones, T. R., E. J. Routman, D. J. Begun, and J. P. Collins. 1995. Ancestry of an isolated
6 subspecies of salamander, *Ambystoma tigrinum stebbinsi* Lowe: the evolutionary
7 significance of hybridization. *Molecular Phylogenetics and Evolution* 4:194-202.
- 8 • Jones, T.R., Collins, J.P., Kocher, T.D. and Mitton, J.B. 1988. Systematic status and
9 distribution of *Ambystoma tigrinum stebbinsi* Lowe (Amphibia: Caudata). *Copeia*.
10 1988:621-635.
- 11 • Snyder, J. D. 1998. Ecology, management, and intellectual history of native and
12 introduced species. Masters Thesis, Arizona State University.
- 13 • Storfer, A., S. G. Mech, M. W. Reudink, R. E. Ziemba, J. Warren, and J. P. Collins. 2004.
14 Evidence for introgression in the endangered Sonora tiger salamander, *Ambystoma*
15 *tigrinum stebbinsi* (Lowe). *Copeia* 2004:783-796.
- 16 • U.S. Fish and Wildlife Service. 2002. Sonora tiger salamander (*Ambystoma tigrinum*
17 *stebbinsi*) recovery plan. U.S. Fish and Wildlife Service, Phoenix, AZ. iv + 67 pp.
- 18 • U.S. Fish and Wildlife Service. 6 January 1997. Determination of endangered status for
19 three wetland species found in southern Arizona and northern Sonora, Mexico. *Federal*
20 *Register* 62(3):665-689.
- 21 • Varela-Romero, A., C. Galindo-Duarte, E. Saucedo-Monarque, L. S. Anderson, P.
22 Warren, S. Stefferud, J. Stefferud, S. Rutman, T. Tibbits, and J. Malusa. 1992. Re-
23 discovery of *Gila intermedia* and *G. purpurea* in northern Sonora, Mexico. In D. A.
24 Hendrickson, editor. "Proceedings of the Desert Fishes Council. Volumes XXII and XXIII,
25 1990 and 1991 Annual Symposia, and Index for Volumes XVI Through XXIII" p. 33,
26 Desert Fishes Council, Bishop, CA.
- 27 • Ziemba, R. E., A. T. Storfer, J. Warren, and J. P. Collins. 1998. A survey of genetic
28 variation among populations of the Sonora tiger salamander (*Ambystoma tigrinum*
29 *stebbinsi*). Report to Arizona Game and Fish Department Heritage Program, Phoenix,
30 AZ.

31

Ramsey Canyon Leopard Frog (*Rana subaquavocalis*)

A conservation agreement among landowners and state and Federal agencies regarding the Ramsey Canyon leopard frog was implemented in 1997. It provides for captive breeding and reintroduction, acquisition of habitat, and habitat and population surveys (Federal Register, 19 September 1997).

Historic Range: Known from areas within a 10-km radius in the (b) (7)(E); current known range is limited to aquatic habitats in Tinker, Brown, Ramsey, and Miller canyons and several residential ponds in the area, (b) (7)(E) County, Arizona (Platz 1993, Platz and Grudzien 1993, Platz et al. 1997, Arizona Game and Fish Department 2001, Platz and Grudzien 2003). Currently exists in several canyons on the east side of the (b) (7)(E) (Goldberg et al. 2004). Ranges from 4,925 to 6,001 ft. (1502 - 1830 m) (Sredl et al. 1997).

Basic Description: A 3-4-inch frog.

General Description: A frog of the *Rana pipiens* complex, with prominent dorsal spots, dorsolateral folds, and extensive webbing on the hind feet. Snout-vent length 81-85 mm in adult males, 86-116 mm in adult females, and 60-62 mm in juveniles (type series, Platz 1993).

Diagnostic Characteristics: Differs from other members of the *Rana pipiens* complex by the following combination of characters: "dorsolateral folds interrupted posteriorly and deflected medially; incomplete supralabial stripe (diffuse anterior to eye); enhanced melanism on venter; yellowish pigmentation on groin, which may extend onto posterior venter; numerous white papillae around cloacal aperture and adjacent dorsum and thighs; stocky body proportions; knob-like terminal swellings on toes in large adults; a long (average length 2.1 sec at 17 C), snore-like mating call consisting of 28-54 pulses of moderate pulse rate (averaging 19.6 pulses/sec at 17 C). The call is given entirely underwater (at a depth of 1.0-1.3 m) and is therefore inaudible in air" (Platz 1993). Differs from *R. yavapaiensis*, *R. pipiens*, and *R. blairi* by the presence of extensive mottling in the chin region. Differs from *R. pipiens* and *R. blairi* by lacking a well-defined, light-colored, complete supralabial stripe. Differs from *R. pipiens* also by lacking continuous dorsolateral folds and green axillary pigmentation, and by having external vocal sacs. Differs from *Rana berlandieri* by the stockier build of adults and by the yellow pigmentation in the groin region (occasionally present to a limited extent in *berlandieri*). Differs from *R. chiricahuensis* in larger adult size and expanded, knob-like toe tips in large adults.

Reproduction Comments: Males vocalize from at least mid-March through mid-July (Platz 1993). Egg masses have been recorded from mid-March through early October (AGFD, unpublished data). Mating seems to begin once water temperatures have reached at least 10 C (50 F), and oviposition may be correlated with temperatures rather than rainfall. Eggs hatch in about 14 days in the wild (Platz 1997). In captivity, eggs hatch in about 10 days when held at 23-25 C (73-77 F) (M. Demlong, unpublished data). Larvae metamorphose in the year they were oviposited or may overwinter as tadpoles (Platz and Grudzien 1993, Platz et al. 1997). Larvae metamorphose in as few as 100 days in captivity, but frequently take 160 to 200 days (M. Demlong, unpublished data). Platz (1997) suggested that sexual maturity is reached rather late in life, at approximately 6 years postmetamorphosis, but captive-reared frogs at the Phoenix Zoo and released in Miller Canyon produced egg masses one year after metamorphosis. Some individuals live at least 10 years after metamorphosis (Platz and Grudzien 1993, Platz et al. 1997). May have a lek breeding system, but further study is needed (Platz and Grudzien 1993).

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: No

Mobility and Migration Comments: Although detailed study of movements has not been done, marked frogs have moved several hundred meters within Ramsey Canyon (M. Sredl, unpublished data) (Arizona Game and Fish Department 2001).

Riverine Habitat(s): CREEK, Low gradient, Moderate gradient, Pool

Palustrine Habitat(s): TEMPORARY POOL

Special Habitat Factors: Benthic

Habitat Comments: Habitats are found in pine-oak, oak woodland, and semi-desert grassland areas of the (b) (7)(E). Vegetation at sites is variable but includes horsetail (*Equisetum* spp.), spikerush (*Eleocharis* spp.), cattail (*Typha* spp.), watercress (*Rorippa*), monkey flower (*Mimulus*), and grasses. Emergent vegetation and root masses provide cover sites (M. Sredl unpublished data) (Arizona Game and Fish Department 2001). Most occupied habitats are modified or artificial aquatic systems (Sredl et al. 1997). Ponds, streams, plunge pools are occupied. Adults and several tadpoles in upper Brown Canyon were found in a plunge pool (elev. 1675 m). Most of the frogs in Ramsey Canyon occupy a ground-level concrete tank (14 m X 14 m) approximately 1.3 m deep, fed by the natural stream adjacent to the tank; frogs also occur at various plunge pools along a 1000 m length of the stream, starting with plunge pools adjacent to the visitors' center and continuing above the tank population. Adults and larvae were observed at a small excavation in rock (a water pocket 2 m in diameter) 2 km below the entrance to Ramsey Canyon (Platz 1993). Occurs also in an earthen stock tank (Platz and Grudzien 1993). Males call while submerged, as may males of certain other RANA species. Eggs are laid in spherical masses, attached to submerged vegetation, so that the egg mass is held near the surface of the water (Arizona Game and Fish Department 2001).

Length: 10 cm

Management Requirements: Management needs include habitat restoration and removal of non-native species; captive rearing of larvae and release of juveniles began in 1995. Arizona Game and Fish Department (AGFD) is attempting to mitigate threats and enhance populations of Ramsey Canyon leopard frogs through captive rearing programs and translocations in the (b) (7)(E) of southeastern Arizona (Sredl et al. 2002). Eggs and larvae have been collected and reared in captivity to increase initial survival rates. The captive-reared frogs and larvae have been released at several sites including Ramsey Canyon, the Barchas Ranch, and Miller Canyon (Arizona Game and Fish Department 2001). An attempt to eradicate bullfrogs from Lower Garden Canyon Pond was unsuccessful (Sredl et al. 2002).

Biological Research Needs: Studies focusing on factors that may play a role in population declines, including the disease caused by chytrid fungus, would be valuable (Arizona Game and Fish Department 2001).

References:

- Arizona Game and Fish Department. 2001. *Rana subaquavocalis*. Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- Berger, L., et al. 1998. Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proceedings of the National Academy of Science* 95:9031-9036.
- Blackburn, L., P. Nanjappa, and M. J. Lannoo. 2001. *An Atlas of the Distribution of U.S. Amphibians*. Copyright, Ball State University, Muncie, IN, USA.

- 1 • Field, K.J., Beatty, Sr, T.L. and Beatty, Jr, T.L. 2003. *Rana subaquavocalis* (Ramsey
2 Canyon leopard) frog diet. Herpetological Review. 34:235.
- 3 • Goldberg, C. S., K. J. Field, and M. J. Sredl. 2004. Mitochondrial DNA sesquences do not
4 support species status of the Ramsey Canyon leopard frog (*Rana subaquavocalis*).
5 Journal of Herpetology 38:313-319.
- 6 • Goldberg, C.S., Field, K.J. and Sredl, M.J. 2003. Ramsey Canyon leopard frog identity
7 crisis: mitochondrial DNA analyses support designation as Chiricahua leopard frog.
8 Arizona Game and Fish Department, Nongame and Endangered Wildlife Program
9 Technical Report 218. Phoenix, AZ.
- 10 • Platz, J. E. 1993. *Rana subaquavocalis*, a remarkable new species of leopard frog (*Rana*
11 *piapiens complex*) from southeastern Arizona that calls under water. Journal of
12 Herpetology 27:154-62.
- 13 • Platz, J. E. 1997. Status survey of the Ramsey Canyon leopard frog, *Rana*
14 *subaquavocalis*. Arizona Game and Fish contract report, Phoenix, AZ.
- 15 • Platz, J. E., A. Lathrop, L. Hofbauer, and M. Vradenburg. 1997. Age distribution and
16 longevity in the Ramsey Canyon leopard frog, *Rana subaquavocalis*. Journal of
17 Herpetology 31:552-557.
- 18 • Platz, J. E., and T. A. Grudzien. 2003. Limited genetic heterozygosity and status of two
19 populations of the Ramsey Canyon leopard frog: *Rana subaquavocalis*. Journal of
20 Herpetology 37:758-761.
- 21 • Platz, J. E., and T. Grudzien. 1993. Geographic distribution and genetic variability
22 assessment of the Ramsey Canyon leopard frog. Final report.
- 23 • Sredl, M. J., J. E. Wallace, and V. Miera. 2000. Aquatic herpetofauna inventory of Fort
24 Huachuca and vicinity. Nongame and Endangered Wildlife Program Technical Report
25 167. Arizona Game and Fish Department, Phoenix, AZ.
- 26 • Sredl, M. J., J. M. Howland, J. E. Wallace, and L. S. Saylor. 1997. Status and distribution
27 of Arizona's native ranid frogs. Pages 37-89 in M. J. Sredl, editor. Ranid frog
28 conservation and management. Arizona Game and Fish Department, Nongame and
29 Endangered Wildlife Program, Technical Report 121.
- 30 • Sredl, M. J., K. J. Field, and A. M. Peterson. 2002. Mitigating threats and managing the
31 Ramsey Canyon leopard frog in Arizona. Nongame and Endangered Wildlife Program
32 Technical Report 207. Arizona Game and Fish Department, Phoenix, AZ.
- 33 • Sredl, M.J. In press *Rana subaquavocalis* Platz 1993, Ramsey Canyon leopard frog.
34 Status and Conservation of U.S. Amphibians. Volume 2: Species Accounts. Lannoo,
35 M.J., editor. University of California Press. Berkeley, CA.
- 36 • Sredl, M.J. and Saylor, L.S. 1998. Conservation and Management Zones and the role of
37 earthen cattle tanks in conserving Arizona leopard frogs on large landscapes.
38 Environmental, Economic, and Legal Issues Related to Rangeland Water Developments.
39 Feller, J.M. and Strouse, D.S., editor. 211-225. The Center for the Study of Law, Science
40 and Technology, Arizona State University. Tempe, AZ.

41

Chiricahua Leopard Frog (*Rana chiricahuensis*)

The Chiricahua leopard frog was designated as a federally Threatened species on June 13, 2002.

Historic Range: This species occurs from southeastern Arizona (drainages of the Madrean Archipelago and surrounding desert grasslands, south of the Gila River in (b) (7)(E) Santa Cruz, Pima, and Graham counties) and extreme southwestern New Mexico (Hidalgo County) in the United States, south along the eastern slope of the Sierra Madre Occidental in Sonora and Chihuahua, Mexico. It occurs at elevations of 1,060-2,010m in Arizona (Arizona Game and Fish Department 1995, Degenhardt et al. 1996, Sredel et al. 1997). Its southern range limit is poorly defined due to taxonomic uncertainties. See RANA SP 1 for information on the distribution of northern montane populations that may represent a different species.

Basic Description: A leopard frog.

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: No

Riverine Habitat(s): CREEK, Pool, SPRING/SPRING BROOK

Lacustrine Habitat(s): Shallow water

Palustrine Habitat(s): Riparian

Special Habitat Factors: Benthic, Fallen log/debris

Habitat Comments: This species occurs in a wide variety of habitats at a wide range of altitudes in pine and pine-oak forests with permanent water ponds of moderate depth as well as montane streams. It is highly aquatic. It breeds in a wide variety of aquatic habitats, ranging from stock ponds, reservoirs, and lakes to spring-fed streams (Jennings and Scott 1993, USFWS 2000).

Adult Food Habits: Invertivore

Immature Food Habits: Herbivore

Food Comments: Adults mainly invertivorous. Larvae eat algae, organic debris, plant tissue, and minute organisms in water.

Phenology Comments: Inactive in cold temperatures.

Length: 14 cm

Management Requirements: See USFWS (2000) for information on management programs.

References:

- Anonymous. 1993. Postmetamorphic death syndrome. Froglog No. 7.
- Arizona Game and Fish Department 1996. Wildlife of special concern in Arizona (public review draft). Nongame and Endangered Wildlife Program. 40 pp.
- Arizona Game and Fish Department. 1992. *Bufo retiformis*. Unpublished abstract. 3 pp.

- 1 • Arizona Game and Fish Department. 1996. Wildlife of special concern in Arizona (public
2 review draft). Nongame and Endangered Wildlife Program, Phoenix, AZ. 40 pp.
- 3 • Blackburn, L., P. Nanjappa, and M. J. Lannoo. 2001. An Atlas of the Distribution of U.S.
4 Amphibians. Copyright, Ball State University, Muncie, IN, USA.
- 5 • Bradley, G.A., Rosen, P.C., Sredl, M.J., Jones, T.R. and Longcore, J.E. 2002.
6 Chytridiomycosis in native Arizona frogs. Journal of Wildlife Diseases. 38:206-212.
- 7 • Clarkson W.R. and Rorabaugh, J.C. 1989. Status of leopard frogs (*Rana pipiens*
8 *complex*: Ranidae) in Arizona and southeastern California. Southwestern Naturalist
9 34:531-538.
- 10 • Clarkson, R. W., and J. C. Rorabaugh. 1989. Status of leopard frogs (*Rana pipiens*
11 *complex*: Ranidae) in Arizona and southeastern California. Southwest. Nat. 34:531-538.
- 12 • Degenhardt, W. G., C. W. Painter, and A. H. Price. 1996. Amphibians and reptiles of New
13 Mexico. University of New Mexico Press, Albuquerque, NM. xix + 431 pp.
- 14 • Fernandez, P.J 1996. A facility for captive propagation of Chiricahua leopard frogs (*Rana*
15 *chiricahuensis*). Herpetoculture. 7-12. International Herpetological Symposium.
- 16 • Frost, D. R. 1985. Amphibian species of the world. A taxonomic and geographical
17 reference. Allen Press, Inc., and The Association of Systematics Collections, Lawrence,
18 KS. v + 732 pp.
- 19 • Frost, J. S., and J. E. Platz. 1983. Comparative assessment of modes of reproductive
20 isolation among four species of leopard frogs (*Rana pipiens complex*). Evolution 37:66-
21 78.
- 22 • Frost, J. S., and J. T. Bagnara. 1977. Sympatry between *Rana blairi* and the southern
23 form of leopard frog in southeastern Arizona (Anura: Ranidae). Southwestern Naturalist
24 22:443-453.
- 25 • Goldberg, C.S., Field, K.J. and Sredl, M.J. 2003. Ramsey Canyon leopard frog identity
26 crisis: mitochondrial DNA analyses support designation as Chiricahua leopard frog.
27 Arizona Game and Fish Department, Nongame and Endangered Wildlife Program
28 Technical Report 218. Phoenix, AZ.
- 29 • Green, D. M., and D. M. Delisle. 1985. Allotriploidy in natural hybrid frogs, *Rana*
30 *chiricahuensis* x *R. pipiens*, from Arizona: chromosomes and electrophoretic evidence.
31 Journal of Herpetology 19:385-390.
- 32 • Hillis, D.M. 1988. Systematics of the *Rana pipiens* complex: puzzle and paradigm.
33 Annual Review of Ecology and Systematics. 19:39-63.
- 34 • Hillis, D.M., Frost, J.S. and Wright, D.A. 1983. Phylogeny and biogeography of the *Rana*
35 *pipiens complex*: a biochemical evaluation. Systematic Zoology. 32:132-143.
- 36 • Jennings, R. D. 1995. Investigations of recently viable leopard frog populations in New
37 Mexico: *Rana chiricahuensis* and *Rana yavapaiensis*. Report submitted to New Mexico
38 Department of Game and Fish, Endangered Species Program, Santa Fe, NM. 36 pp.
- 39 • Jennings, R. D., and N. J. Scott, Jr. 1993. Ecologically correlated morphological variation
40 in tadpoles of the leopard frog, *Rana chiricahuensis*. Journal of Herpetology 27:285-293.
- 41 • Jennings, R.D ND. Activity and reproductive phenologies and their ecological correlates
42 among populations of the Chiricahua leopard frog, *Rana chiricahuensis*. Unpublished
43 report for New Mexico Department of Game and Fish Santa Fe, NM.
- 44 • Jennings, R.D. 1988. Ecological studies of the Chiricahua leopard frog, *Rana*
45 *chiricahuensis*, in New Mexico. Report to Share with Wildlife, New Mexico Department of
46 Game and Fish. Santa Fe, NM.

- 1 • Jennings, R.D. 1995. Investigations of recently viable leopard frog populations in New
2 Mexico: *Rana chiricahuensis* and *Rana yavapaiensis*. 36 p.
- 3 • Mecham, J.S. 1968. Evidence of reproductive isolation between two populations of the
4 frog, *Rana pipiens*, in Arizona. Southwestern Naturalist. 13:35-44.
- 5 • New Mexico Department of Game and Fish. 1996. October 1-last update. Fish and
6 Wildlife Information Exchange-VA Tech. Online. Available:
7 <http://www.fw.vt.edu/fishex/nm.html>. Accessed 1997, April 8.
- 8 • Platz, J. E. and J. S. Mecham. 1979. *Rana chiricahuensis*, a new species of leopard frog
9 (*Rana pipiens complex*) from Arizona. Copeia 1979:383-90.
- 10 • Platz, J.E. and Mecham, J.S. 1984. *Rana chiricahuensis*. Catalogue of American
11 Amphibians and Reptiles. 347:1-2.
- 12 • Rosen, P.C., Schwalbe, C.R., Parizek, D.A.J., Holm, P.A. and Lowe, C.H. 1995.
13 Introduced aquatic vertebrates in the Chiricahua region: effects on declining ranid frogs.
14 Biodiversity and management of the Madrean Archipelago: the sky islands of
15 southwestern United States and northwestern Mexico. DeBano, L.F., Gottfried, G.J.,
16 Hamre, R.H. and Edmi, C.B., editor. Rocky Mountain Forest and Range Experimental
17 Station. Fort Collins, CO.
- 18 • Schwalbe, C. R. 1993. Status of amphibians in Arizona. Park Science 13(4):10.
- 19 • Scott, N.J. and R.D. Jennings 1985. The tadpoles of five species of New Mexican leopard
20 frogs. Occasional papers the Museum of Southwest Biology. No. 3, Dec. 9, 1985. 21Pp.
- 21 • Sredl, M. 1993. Global amphibian decline: have Arizona's amphibians been affected?
22 Sonoran Herpetologist 6(2):14-21.
- 23 • Sredl, M. J., J. M. Howland, J. E. Wallace, and L. S. Saylor. 1997. Status and distribution
24 of Arizona's native ranid frogs. Pages 37-89 in M. J. Sredl, editor. Ranid frog
25 conservation and management. Arizona Game and Fish Department, Nongame and
26 Endangered Wildlife Program, Technical Report 121.
- 27 • Sredl, M.J. and Jennings, R.D. In press *Rana chiricahuensis* (Platz and Mecham, 1979)
28 Chiricahua Leopard Frogs. Status and Conservation of U.S. Amphibians. Volume 2:
29 Species Accounts. Lannoo, M.J., editor. University of California Press. Berkeley, CA.
- 30 • Sredl, M.J. and Saylor, L.S. 1998. Conservation and Management Zones and the role of
31 earthen cattle tanks in conserving Arizona leopard frogs on large landscapes.
32 Environmental, Economic, and Legal Issues Related to Rangeland Water Developments.
33 Feller, J.M. and Strouse, D.S., editor. 211-225. The Center for the Study of Law, Science
34 and Technology, Arizona State University. Tempe, AZ.
- 35 • Stebbins, R. C. 1985. A field guide to western reptiles and amphibians. Second edition.
36 Houghton Mifflin Company, Boston, MA. xiv + 336 pp.
- 37 • U.S. Fish and Wildlife Service (USFWS). 13 June 2002. Listing of the Chiricahua leopard
38 frog (*Rana chiricahuensis*). Federal Register 67(114):40790-40811.
- 39 • U.S. Fish and Wildlife Service (USFWS). 14 June 2000. Proposal to list the Chiricahua
40 leopard frog as threatened with a special rule. Federal Register 65:37343-37357.
- 41 • Wood, T. 1991. Results of 1991 amphibian monitoring on the Coronado National Forest.
42 The Nature Conservancy and U.S. Forest Service.

1 New Mexico Ridgenose Rattlesnake (*Crotalus willardi obscures*)

2 The New Mexico ridge-nosed rattlesnake was designated as a federally Threatened species on
3 August 4, 1978.

4 **Historic Range:** This snake occurs locally in Animas Mountains (New Mexico), Peloncillo
5 Mountains (Arizona and New Mexico), and Sierra de San Luis (Sonora and Chihuahua, Mexico)
6 (Campbell et al. 1989, Holycross and Smith 1997, Campbell and Lamar 2004).

7 **Basic Description:** A rattlesnake.

8 **Reproduction Comments:** Viviparous. Bears 2-9 young, August-September.

9 **Non-Migrant:** No

10 **Locally Migrant:** No

11 **Long Distance Migrant:** No

12 **Palustrine Habitat(s):** Riparian

13 **Terrestrial Habitat(s):** Bare rock/talus/scree, Woodland - Conifer, Woodland - Hardwood,
14 Woodland - Mixed

15 **Special Habitat Factors:** Burrowing in or using soil, Fallen log/debris

16 **Habitat Comments:** Primarily at high elevations in pine-oak woodland and pine-fir forest but
17 also found in foothill canyons in pinyon-juniper woodland. Inhabits canyon bottoms with canopies
18 of alder, box elder, maple, etc. (Stebbins 1985). Hides in leaf litter among cobbles and rocks;
19 frequently climbs into trees and shrubs (Matthews and Moseley 1990).

20 **Adult Food Habits:** Carnivore, Invertivore

21 **Immature Food Habits:** Carnivore, Invertivore

22 **Food Comments:** Preys on scorpions, centipedes, lizards, small mammals and birds.

23 **Adult Phenology:** Diurnal, Hibernates/aestivates

24 **Immature Phenology:** Diurnal, Hibernates/aestivates

25 **Phenology Comments:** Inactive in cold temperatures and extreme heat. Mainly diurnal but
26 probably at least partially nocturnal during hot summer weather; in summer, most active on warm
27 humid mornings; rains may stimulate late afternoon activity; in fall, active mainly in afternoon
28 (Ernst 1992). Most active during daylight hours from July through September.

29 **Length:** 61 cm

30 References:

- 31 • Armstrong, B. L., and J. B. Murphy. 1979. The natural history of Mexican rattlesnakes.
32 Univ. Kansas Museum Natural History Special Publication. No. 5. vii + 88 pp.
- 33 • Campbell, J. A., and E. D. Brodie, Jr., editors. 1992. Biology of the pit vipers. Selva,
34 Tyler, TX.

- 1 • Campbell, J. A., and W. W. Lamar. 1989. The venomous reptiles of Latin America.
2 Comstock Publ. Associates, Division of Cornell Univ. Press, Ithaca, NY. xii + 425 pp.
- 3 • Campbell, J. A., and W. W. Lamar. 2004. The venomous reptiles of the Western
4 Hemisphere. 2 volumes. Cornell University Press, Ithaca, NY.
- 5 • Collins, J. T. 1990. Standard common and current scientific names for North American
6 amphibians and reptiles. 3rd ed. Society for the Study of Amphibians and Reptiles.
7 Herpetological Circular No. 19. 41 pp.
- 8 • Ernst, C. H. 1992. Venomous reptiles of North America. Smithsonian Institution Press,
9 Washington, D.C. ix + 236 pp.
- 10 • Gloyd, H. 1940. The rattlesnakes. Chicago Academy Science, Special Publication No. 4.
- 11 • Holycross, A. T., and S. W. Smith. 1997. Geographic distribution: *Crotalus willardi*
12 *obscurus*. Herpetological Review 28:97.
- 13 • Klauber, L. M. 1972. Rattlesnakes: their habits, life histories, and influence on mankind.
14 Second edition. Two volumes. Univ. California Press, Berkeley, CA.
- 15 • Lowe, C. H., C. R. Schwalbe, and T. B. Johnson. 1986. The venomous reptiles of
16 Arizona. Arizona Game and Fish Department. ix + 115 pp.
- 17 • Matthews, J.R. and C.J. Moseley (eds.). 1990. The Official World Wildlife Fund Guide to
18 Endangered Species of North America. Volume 1. Plants, Mammals. xxiii + pp 1-560 +
19 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians,
20 Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180.
21 Beacham Publications, Inc., Washington, D.C.
- 22 • Stebbins, R. C. 1985. A field guide to western reptiles and amphibians. Second edition.
23 Houghton Mifflin Company, Boston, MA. xiv + 336 pp.
- 24

Western Yellow-billed Cuckoo (*Coccyzus americanus occidentalis*)

The western yellow-billed cuckoo was designated as a Federal Candidate species on September 12, 2006.

Historical Range: BREEDING: interior California to southern Idaho, southeastern Montana, the Dakotas, southern Manitoba (rarely), Minnesota, and New Brunswick, south to southern Baja California, southern Arizona, Coahuila, Chihuahua, Nuevo Leon, Tamaulipas, Gulf Coast, and Florida Keys; sporadically farther south in Mexico and in the Greater Antilles (AOU 1998). Uncommon on Cuba, Hispaniola, and Puerto Rico; rare in Virgin Islands, Jamaica, and northern Lesser Antilles (Saint Martin); possibly in Bahamas and Lesser Antilles (Raffaele et al. 1998). Bred formerly in British Columbia, Washington, and Oregon. NONBREEDING: southern Central America (rare and local in Costa Rica) and northern South America (and Trinidad and Tobago) south to eastern Peru, Bolivia, and northern Argentina (AOU 1998); rare in West Indies (Raffaele et al. 1998).

Basic Description: A bird (cuckoo).

Reproduction Comments: Breeding often coincides with the appearance of massive numbers of cicadas, caterpillars, or other large insects (Ehrlich et al. 1992). Clutch size is one to five (commonly two to three), largest when prey is abundant. Clutch sizes greater than six attributable to more than one female laying in nest (Hughes 1999). Incubation lasts 9-11, shared by male and female during day; male incubates at night (Hamilton and Hamilton 1965, Potter 1980, Potter 1981). Young are tended by both parents, climb in branches at seven-nine days. Sometimes lays eggs in the nests of Black-billed Cuckoo (*Coccyzus erythrophthalmus*) or (rarely) other species (Ehrlich et al. 1992).

Ecology Comments: Territory size averages 20-24 hectares (S. Laymon, in Riparian Habitat Joint Venture 2000). Known predators of adults include Aplomado Falcon (*Falco femoralis*), Red-shouldered Hawk (*Buteo lineatus*), and other raptors; of eggs and young include Blue Jay (*Cyanocitta cristata*), Common Grackle (*Quiscalus quiscula*), Black Racer (*Coluber constrictor*) and Eastern Chipmunk (*Tamias striatus*) (Hughes 1999). Occasional host for Brown-headed Cowbird (*Molothrus ater*), Bronzed Cowbird (*Molothreus aeneus*), and Black-billed Cuckoo (*Coccyzus erythrophthalmus*) (Hughes 1999).

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: Yes

Mobility and Migration Comments: Migrates regularly through the southern U.S., Middle America, and West Indies (sometimes large numbers in fall in Puerto Rico, Raffaele 1983). Birds from North America may migrate through Puerto Rico, but a small breeding population may be resident all year (Kepler and Kepler 1978). Migrants noted in April-May in Jamaica (Lack 1976). Migrates through Costa Rica mid-August to early November and late April-early June (Stiles and Skutch 1989). Arrives in California breeding grounds usually in early June (Biosystems Analysis 1989).

Estuarine Habitat(s): Scrub-shrub wetland

Palustrine Habitat(s): Riparian

Terrestrial Habitat(s): Forest - Hardwood, Forest - Mixed, Old field, Shrubland/chaparral, Suburban/orchard, Woodland - Hardwood, Woodland - Mixed

Habitat Comments: BREEDING: Open woodland (especially where undergrowth is thick), parks, deciduous riparian woodland; in the West, nests in tall cottonwood and willow riparian woodland. Nests in deciduous woodlands, moist thickets, orchards, overgrown pastures; in tree, shrub, or vine, an average of 1-3 meters above ground (Harrison 1979). Subspecies *occidentalis* requires patches of at least 10 hectares (25 acres) of dense riparian forest with a canopy cover of at least 50 percent in both the understory and overstory; nests typically in mature willows (Biosystems Analysis 1989). NON-BREEDING: forest, woodland, and scrub. Also mangroves in Puerto Rico (Raffaele 1983).

Adult Food Habits: Invertivore

Immature Food Habits: Invertivore

Food Comments: Eats mainly caterpillars; also other insects, some fruits, sometimes small lizards and frogs and bird eggs (Terres 1980). Gleans food from branches or foliage, or sallies from a perch to catch prey on the wing (Ehrlich et al. 1992).

Adult Phenology: Diurnal

Immature Phenology: Diurnal

Length: 31 cm

Weight: 64 grams

Stewardship Overview: Summer distribution throughout much of the eastern and Midwestern United States. Once common in the west, now rare and local, extirpated from British Columbia, Washington, Oregon, possibly Nevada. Winters primarily in South America east of the Andes, may breed in the tropics. Blue listed by Tate (1981). Western population currently under review for federal listing by USFWS; does not yet receive adequate federal due primarily to controversy surrounding the validity of its subspecies status. Listed as endangered in California, listed as threatened or endangered in every western state in which it occurs. From 1980 to 1994 eastern populations declined in all states except Louisiana and South Carolina. Highly significant declines in Alabama, Georgia, Illinois, Indiana, Michigan, New Jersey, New York, Ohio, Pennsylvania, Texas and Wisconsin, with the greatest decline in Connecticut. Main threats are habitat fragmentation, degradation of riparian woodland due to agricultural and residential development (Dobkin 1994), stochastic extinctions and low colonization rates, flood control (Laymon and Halterman 1987, 1989), riparian habitats invaded by less desirable salt cedar (*TAMARIX* spp.; Hughes 1999). Highly vulnerable to continued tropical deforestation (Morton 1992), but direct effects on population numbers not quantified. Preserves in the west should include riparian areas with dense stands of cottonwood and willow with an average tree height of 10-15 meters (Anderson and Laymon 1989). Preserves in the east should have open woodlands with clearings and low, dense, shrubby vegetation, associated with watercourses. Management should focus on acquiring and improving riparian habitats, and eliminating pesticide spraying near habitats.

Restoration Potential: May recolonize if suitable habitat is restored. On experimentally replanted sites (11 hectares) in southern California, foraged in second year and nested in third year following replanting, provided that cottonwood growth averaged 3 meters per year. Sites with growth of 2 meters per year or less not used for foraging or nesting by third year (Anderson and Laymon 1989).

Preserve Selection & Design Considerations: In California, Gaines (1974) defined habitat as willow and cottonwood forests below 1300 meters elevation, greater than 10 hectares in extent, and wider than 100 meters. Laymon and Halterman (1989) concluded that sites greater than 80 hectares (200 acres) in extent and wider than 600 meters (1950 feet) were optimal (100 percent

1 occupancy), sites 41-80 hectares (101-200 acres) in extent and wider than 200 meters (650 feet)
2 were suitable (58.8 percent), sites 20-40 hectares (50-100 acres) in extent and 100-200 meters
3 (325-650 feet) in width were marginal (9.5 percent), and sites less than 15 hectares (38 acres) in
4 extent and less than 100 meters (325 feet) in width were unsuitable. During a four-year study on
5 the Sacramento River, Halterman (1991) found that habitat patch area, the extent of habitat in a 8
6 kilometer (5 mile) section of river, and presence of low woody vegetation were the most important
7 variables in explaining the distribution of cuckoos. These variables combined explained 46
8 percent of the variation observed in the distribution of breeding pairs. Microhabitat requirements
9 are also important. Nesting groves at the South Fork Kern River are characterized by higher
10 canopy closure, higher foliage volume, intermediate basal area, and intermediate tree height
11 when compared to random sites (Laymon et al. 1997). Sites with less than 40 percent canopy
12 closure are unsuitable, those with 40 - 65 percent are marginal to suitable, and those with greater
13 than 65 percent are optimal (Laymon 1998). Lower nesting success for open-cup nesting birds
14 near edges in large habitats and in smaller habitat fragments (Chasko and Gates 1982, Gates
15 and Gysel 1978), and increased nest predation reaching up to 600 meters into forest interior
16 (Wilcove 1985) indicate that reserves less than 100 hectares are less valuable than larger
17 reserves (Wilcove et al. 1986). Simulation modeling demonstrates that populations of fewer than
18 10 pairs are very unstable and always become extinct in a short period of time (Richter-Dyn and
19 Goel 1972, Roth 1974); a minimum number of 25 pairs in a subpopulation with interchange to
20 other subpopulation should be reasonably safe from extinction by stochastic events (Hughes
21 1999). In the northeast and central U.S., and southern Canada, preserves should include
22 woodland, abandoned farmland, overgrown fruit orchards, successional shrubland, dense
23 thickets along streams and marshes (Johnsgard 1979, Peck and James 1983, Eaton 1988,
24 Jauvin 1996), shade trees, gardens (Oberholser 1974). In midwest U.S., also uses willow-
25 dogwood shrub wetlands, and successional hardwood forest with dense stands of small trees 1-7
26 meters in height; e.g., American Elm and or continuous stands of dense Hawthorn (Nolan 1963,
27 Eastman 1991). In southeastern U.S. occupies hammocks and hardwood forest, particularly
28 those crossed by streams, thickets, swamps, and fencerows (Stevenson and Anderson 1994).

29 **Management Requirements:** See California Department of Fish and Game (1990) for a listing
30 of management needs in California. In the west, conservation recommendations summarized in
31 Laymon (1980) include: determine numbers and locations of remnant populations; improve
32 existing, and acquire new riparian habitats; eliminate pesticide spraying in orchards adjacent to
33 riparian areas; and investigate feasibility of captive breeding and reintroduction to naturally
34 regenerated or reforested habitat. Riparian vegetation propagation and site management
35 techniques are outlined in Anderson and Laymon (1989). Grazing should be removed to allow
36 natural regeneration and encourage increased density of cottonwoods and willows.

37 **Monitoring Requirements:** Population densities may be highly variable locally (Eaton 1988)
38 depending on food availability; large localized influxes during times of insect abundances (Veit
39 and Petersen 1993). Estimates made over 1-2 year period must be assessed with caution
40 (Groschupf 1987). Population density may be underestimated due to quiet demeanor and
41 skulking behavior, easily overlooked when silent. Conventional observation, mist netting
42 (Rappole et al. 1993), or listening-post techniques are inadequate for estimating density; counting
43 responses to playback is preferable (Hamilton and Hamilton 1965). Overlapping territories
44 increase difficulty in monitoring and the only way to get a complete survey is to locate all or most
45 of the nests which is a very time-consuming and difficult task (Laymon, pers. comm.).

46 **Management Programs:** On the South Fork Kern River, an experimental study using riparian
47 restoration showed that the number of pairs is closely related to the amount of available habitat.
48 This site had a restoration program which began in 1996 and has established 125 hectares (310
49 acres) of willow-cottonwood habitat on the Kern River Preserve, all of which was being used by
50 cuckoos by the summer of 1996. An additional 510 hectares (1275 acres) of habitat was
51 established by natural regeneration in the South Fork Wildlife Area and the Isabella Reservoir
52 Draw-Down Zone between 1987 and 1992 (Laymon 1998).

Biological Research Needs: Need to determine cause(s) of declines in eastern and central populations.

References:

- Allen, Craig R., Stephen DeMarais, and R. Scott Lutz. 1994. Red Imported Fire Ant Impact on Wildlife: an Overview. Texas J. Sci. 46(1):51-59.
- American Ornithologists' Union (AOU). 1983. Check-list of North American Birds, 6th edition. Allen Press, Inc., Lawrence, KS. 877 pp.
- American Ornithologists' Union (AOU). 1998. Check-list of North American birds. Seventh edition. American Ornithologists' Union, Washington, DC. 829 pp.
- American Ornithologists' Union. 1957. Checklist of North American Birds. 5th edition. Baltimore: Lord Baltimore Press. 691 pp.
- Anderson, B.W., and S.A. Laymon. 1989. Creating habitat for the Yellow-billed Cuckoo (*Coccyzus americanus*). Pages 468-472 in USDA Forest Service, General Technical Report PSW-110.
- Aquin, P. 1999. Évaluation de la situation des groupes taxonomiques des oiseaux du Québec. Ministère de l'Environnement et de la Faune. 13 pages.
- Arnett, John H. Jr., E.A. Choat, R.T. Darby, P.W. Schwalbe, Andm. Weiss. 1954. A Field List of Birds of the Philadelphia Region. Delaware Valley Ornithological Club. 45 pp.
- Audubon Society. 1981-1985. Breeding Bird Atlas of New Hampshire. (unpublished).
- Baicich, P. J., and C. J. O. Harrison. 1997. A guide to the nests, eggs and nestlings of North American birds. Second edition. Academic Press, NY.
- Bancroft, G. T., A. M. Strong, and M. Carrington. 1995. Deforestation and its effects on forest-nesting birds in the Florida Keys. Conservation Biology 9: 835-844.
- Banks, R. C. 1988. Geographic variation in the yellow-billed cuckoo. Condor 90:473-477.
- Behle, W. H., E. D. Sorensen, and C. M. White. 1985. Utah birds: a revised checklist. Occas. Publ. No. 4, Utah Museum of Natural History, Univ. of Utah, Salt Lake City, UT. xv + 108 pp.
- Bent, A.C. 1940. Life histories of North American cuckoos, goatsuckers, hummingbirds, and their allies. Part I. U.S. National Museum Bulletin 176. 244 pp.
- Bierly, M.L. 1980. Bird Finding in Tennessee. 3825 Bedford Ave., Nashville, TN 37125.
- Biosystems Analysis, Inc. 1989. Endangered Species Alert Program Manual: Species Accounts and Procedures. Southern California Edison Environmental Affairs Division.
- BirdLife International. 2004. Threatened birds of the world 2004. CD ROM. BirdLife International, Cambridge, UK.
- Braun, M. J., D. W. Finch, M. B. Robbins, and B. K. Schmidt. 2000. A field checklist of the birds of Guyana. Smithsonian Institution, Washington, D.C.
- Burleigh, T. D. 1972. Birds of Idaho. The Caxton Printers, Ltd., Caldwell, Idaho. 467 pp.
- Cadman, M. D., P. F. J. Eagles, and F. M. Helleiner. 1987. The Atlas of Breeding Birds of Ontario. University of Waterloo Press, Waterloo, Canada. 617pp.
- California Department of Fish and Game. 1990. 1989 annual report on the status of California's state listed threatened and endangered plants and animals. 188 pp.
- Cannings, S. 2001. EO Specifications for Yellow-billed Cuckoo (*Coccyzus americanus*). NatureServe, Unpublished. 1 pp.

- 1 • Carter, M., G. Fenwick, C. Hunter, D. Pashley, D. Petit, J. Price, and J. Trapp. 1996.
2 Watchlist 1996: For the future. Field Notes 50(3):238-240.
- 3 • Chasko, G. G., and J. E. Gates. 1982. Avian habitat suitability along a transmission-line
4 corridor in an oak-hickory forest region. Wildlife Monographs, Vol. 82.
- 5 • Dickinson, Mary B., ed. 1999. Field Guide to the Birds of North America, 3rd ed. National
6 Geographic Society, Washington, D.C. 480 pp.
- 7 • DeGraaf, R.M., and J.H. Rappole. 1995. Neotropical migratory birds: natural history,
8 distribution, and population change. Comstock Publishing Associates, Ithaca, NY.
- 9 • DeCalesta, D. S. 1994. Effect of white-tailed deer on songbirds within managed forests in
10 Pennsylvania. Journal of Wildlife Manage. 58: 711-718.
- 11 • Desrosiers A., F. Caron et R. Ouellet. 1995. Liste de la faune vertébrée du Québec. Les
12 publications du Québec. 122 pp.
- 13 • Dionne C. 1906. Les oiseaux de la province de Québec. Dussault et Proulx.
- 14 • Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the
15 Northern Rockies and Great Plains. University of Idaho Press, Moscow, ID.
- 16 • Droege, S., and J.R. Sauer. 1990. North American Breeding Bird Survey, annual
17 summary, 1989. U.S. Fish and Wildlife Service, Biological Report 90(8). 22 pp.
- 18 • Dunn, E. H., C. M. Downes, and B. T. Collins. 2000. The Canadian Breeding Bird Survey,
19 1967-1998. Canadian Wildlife Service Progress Notes No. 216. 40 pp.
- 20 • Eastman, J. 1991. Yellow-billed Cuckoo. Pages 234-245 in R. Brewer, G.A. McPeck, and
21 R.J. Adams, Jr., editors. The atlas of breeding birds of Michigan. Michigan State
22 University Press, East Lansing, MI.
- 23 • Eaton, S. W. 1988. Yellow Billed Cuckoo. Pages 198-199 in: The Atlas of Breeding Birds
24 of New York State. Cornell Univ. Press, Ithaca, NY.
- 25 • Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to
26 the natural history of North American birds. Simon and Shuster, Inc., New York. xxx +
27 785 pp.
- 28 • Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1992. Birds in Jeopardy: the Imperiled and
29 Extinct Birds of the United States and Canada, Including Hawaii and Puerto Rico.
30 Stanford University Press, Stanford, CA. 259 pp.
- 31 • Erskine, A. J. 1992. Atlas of breeding birds of the Maritime Provinces. Nimbus Publishing
32 and the Nova Scotia Museum, Halifax, Nova Scotia.
- 33 • Franzreb, K. E. and S. A. Laymon. 1993. A reassessment of the taxonomic status of the
34 Yellow-billed Cuckoo. Western Birds 24:17-28.
- 35 • Gaines, D. 1974. Review of the status of the Yellow-billed Cuckoo in California:
36 Sacramento Valley populations. Condor 76:204-209.
- 37 • Gaines, D., and S.A. Laymon. 1984. Decline, Status and Preservation of the Yellow-billed
38 Cuckoo in California. Western Birds 15(2):49-80.
- 39 • Gates, J.E., and L.W. Gysel. 1978. Avian nest dispersion and fledgling success in field-
40 forest ecotones. Ecology 59:871-883.
- 41 • Godfrey, Earl W. 1986. Birds of Canada. National Museums of Canada, Ottawa, Ontario,
42 Canada. 595 p. + plates.
- 43 • Groschupf, K. 1987. Status of the Yellow-billed Cuckoo (*Coccyzus americanus*
44 *occidentalis*) in Arizona and west Texas. USFWS Rep. 20181-86-00731. Phoenix, AZ.

- 1 • Hagan, J. M., III, and D. W. Johnston, editors. 1992. Ecology and conservation of
2 neotropical migrant landbirds. Smithsonian Institution Press, Washington, D.C. xiii + 609
3 pp.
- 4 • Halterman, M.D. 1991. Distribution and habitat use of the Yellow-billed Cuckoo
5 (*Coccyzus americanus occidentalis*) on the Sacramento River, California, 1987-1990.
6 Masters Thesis, California State University, Chico.
- 7 • Hamilton, W.J., III, and M.E. Hamilton. 1965. Breeding characteristics of Yellow-billed
8 Cuckoos in Arizona. Proc. Calif. Acad. Sci. 32:405-432.
- 9 • Harrison, H. H. 1979. A field guide to western birds' nests. Houghton Mifflin Company,
10 Boston, MA. 279 pp.
- 11 • Hayward, C. L., C. Cottam, A. M. Woodbury, and H. H. Frost. 1976. Birds of Utah. Great
12 Basin Naturalist Memoirs 1: 229 pp.
- 13 • Heritage database. Washington Department of Fish and Wildlife. October 30, 2002.
- 14 • Howell, S. N. G., and S. Webb. 1995. A guide to the birds of Mexico and northern Central
15 America. Oxford University Press, Oxford, UK.
- 16 • Hughes, J. M. 1999. Yellow-billed Cuckoo (*Coccyzus americanus*). Birds of North
17 America 418: 1-27.
- 18 • Hughes, J.M. 1999. Yellow-billed Cuckoo (*Coccyzus americanus*). In A. Poole and F. Gill,
19 editors, The Birds of North America, No. 418. The Birds of North America, Inc.,
20 Philadelphia, PA. 28 pp.
- 21 • Hughes, J.M. 1999. Yellow-billed cuckoo (*Coccyzus americanus*). No. 318 IN A. Poole
22 and F. Gill, editors. The birds of North America. The Birds of North America, Inc.,
23 Philadelphia, PA. 28 pp.
- 24 • Hunter, W. C., R. D. Ohmart, and B. W. Anderson. 1988. Use of exotic saltcedar
25 (*Tamarix chinensis*) by birds in arid riparian systems. Condor 90:113-123.
- 26 • Jewett, S.G., W.P. Taylor, W.T. Shaw, and J.W. Aldrich. 1953. Birds of Washington
27 State. U. Washington Press. 767 pp.
- 28 • Johnsgard, P. A. 1979. Birds of the Great Plains: breeding species and their distribution.
29 Univ. Nebraska Press, Lincoln, NE. 539 pp.
- 30 • Johnsgard, P. A. 1986. Birds of the Rocky Mountains with particular reference to national
31 parks in the Northern Rocky Mountain region. Colorado Associated University Press,
32 Boulder, CO. xi + 504 pp.
- 33 • Johnson, D.H. and T.A. O'Neil. 2001. Wildlife-habitat relationships in Oregon and
34 Washington. Oregon State University Press. Corvallis. 736 pp.
- 35 • Juavin, D. 1996 Yellow-billed Cuckoo. Pages 582-583 in The Breeding birds of Quebec:
36 atlas of breeding birds of southern Quebec. Association Quebecoise des Groupes d'
37 Ornithologues, Province of Quebec Society for the Protection of Birds, Can. Wildl. Serv.,
38 Environ. Can./ Montreal.
- 39 • Kepler, C. B., and A. K. Kepler. 1978. Status and nesting of the yellow-billed cuckoo in
40 Puerto Rico. Auk 95:417-419.
- 41 • Lack, D. 1976. Island biology illustrated by the land birds of Jamaica. Studies in Ecology,
42 Vol. 3. Univ. California Press, Berkeley, CA. 445 pp.
- 43 • Larrison, E. J., J. L. Tucker, and M. T. Jollie. 1967. Guide to Idaho birds. Journal of the
44 Idaho Academy of Science 5: 1-220.

- 1 • Laughlin, S.B. and D.P. Kibbe, eds. 1985. The Atlas of Breeding Birds of Vermont. Univ.
2 Press of New England. 456pp.
- 3 • Laymon, S. A. 1998. Partners in Flight Bird Conservation Plan for Yellow-billed Cuckoo,
4 website.
- 5 • Laymon, S. A., and M. D. Halterman. 1987b. Can the western subspecies of Yellow-billed
6 Cuckoo be saved from extinction? Western Birds 18:19-25.
- 7 • Laymon, S.A. and M.D. Halterman. 1987. Can the Western Subspecies of the Yellow-
8 billed Cuckoo be Saved from Extinction? Western Birds 18:19-25.
- 9 • Laymon, S.A., P.L. Williams, and M.D. Halterman. 1997. Breeding status of the Yellow-
10 billed Cuckoo in the South Fork Kern River Valley, Kern County, California: Summary
11 report 1985 - 1996. Administrative Rep. U.S.D.A Forest Service, Sequoia National
12 Forest, Cannell Meadow Ranger District, Challenge Cost-Share Grant #92-5-13.
- 13 • Laymon, S.A., and M.D. Halterman. 1989. A proposed habitat management plan for
14 Yellow-billed Cuckoos in California. USDA Forest Service, Gen. Tech. Rep. PSW-110.
- 15 • Lowery, George H. 1974. The Birds of Louisiana. LSU Press. 651pp.
- 16 • McAtee W.L. 1959. Folk - names of Canadian birds. National Museum of Canada. Folk -
17 names of Canadian birds. National Museum of Canada. 74 pages.
- 18 • Montana Bird Distribution Online Database. 2001. Helena, Montana, USA. April-
19 September 2003. <http://nhp.nris.state.mt.us/mbd/>.
- 20 • Morton, E.S. 1992. What do we know about the future of migrant landbirds? Pages 579-
21 589 in J.M. Hagan III and D.W. Johnston, editors. Ecology and conservation of
22 neotropical migrant landbirds. Smithsonian Institution Press, Washington, DC.
- 23 • National Geographic Society (NGS). 1999. Field guide to the birds of North America.
24 Third edition. National Geographic Society, Washington, DC. 480 pp.
- 25 • Nicholson, C.P. 1997. Atlas of the breeding birds of Tennessee. The University of
26 Tennessee Press. 426 pp.
- 27 • Nolan, V., Jr. 1963. Reproductive success of birds in a deciduous scrub habitat. Ecology
28 44:305-313.
- 29 • Oberholser, H.C. 1974. The bird life of Texas. 2 vols. Univ. of Texas Press, Austin, TX.
- 30 • Ouellet H., M. Gosselin et J.P. Artigau. 1990. Nomenclature française des oiseaux
31 d'Amérique du Nord. Secrétariat d'État du Canada. 457 p.
- 32 • Parker III, T. A., D. F. Stotz, and J. W. Fitzpatrick. 1996. Ecological and distributional
33 databases for neotropical birds. The University of Chicago Press, Chicago, IL.
- 34 • Parks Canada. 2000. Vertebrate Species Database. Ecosystems Branch, 25 Eddy St.,
35 Hull, PQ, K1A 0M5.
- 36 • Peck, G. K., and R. D. James. 1983. Breeding birds of Ontario, nidology and distribution,
37 volume 1: nonpasserines. Royal Ontario Museum, Toronto, Canada.
- 38 • Pennsylvania Breeding Bird Atlas Project. 1984-1987. Unpublished.
- 39 • Poole, A. F. and F. B. Gill. 1992--. The birds of North America. The American
40 Ornithologists' Union, Washington, D.C. and The Academy of Natural Sciences,
41 Philadelphia, PA.
- 42 • Potter, E. F. 1980. Notes on nesting Yellow-billed Cuckoos. Journal of Field Ornithology
43 51:17-29.

- 1 • Potter, E. F. 1981. Effects of cool weather on nesting behavior and development in the
2 Yellow-billed Cuckoo. *Chat* 45:15-16.
- 3 • Price, J., S. Droege, and A. Price. 1995. The summer atlas of North American birds.
4 Academic Press, NY. x + 364 pp.
- 5 • Raffaele, H. A. 1983. A guide to the birds of Puerto Rico and the Virgin Islands. Fondo
6 Educativo Interamericano, San Juan, Puerto Rico. 255 pp.
- 7 • Raffaele, H., J. Wiley, O. Garrido, A. Keith, and J. Raffaele. 1998. A guide to the birds of
8 the West Indies. Princeton University Press, Princeton, NJ. 511 pp.
- 9 • Rappole, J.H., W.J. McShea, and J. Vega-Rivera. 1993. Evaluation of two survey
10 methods in upland avian breeding communities. *Journal of Field Ornithology* 64(1):55-70.
- 11 • Richter-Dyn, N. and N.S. Goel. 1972. On the extinction of a colonizing species.
12 *Theoretical Population Biology* 3:406-433.
- 13 • Ridgely, R. S. 2002. Distribution maps of South American birds. Unpublished.
- 14 • Ridgely, R. S. and J. A. Gwynne, Jr. 1989. A Guide to the Birds of Panama. 2nd edition.
15 Princeton University Press, Princeton, USA.
- 16 • Riparian Habitat Joint Venture. 2000. Version 1.0. The riparian bird conservation plan: a
17 strategy for reversing the decline of riparian associated birds in California. California
18 Partners in Flight.
- 19 • Riparian Habitat Joint Venture. 2000. Version 1.0. The riparian bird conservation plan: a
20 strategy for reversing the decline of riparian associated birds in California. California
21 Partners in Flight. Available at: <http://www.prbo.org/CPIF/Riparian/ripariandl.html>.
- 22 • Roth, D.A. 1974. The analysis of a population model demonstrating the importance of
23 dispersal in a heterogeneous environment. *Oecologia* 15:259-275.
- 24 • Ryke, N., D. Winters, L. McMartin and S. Vest. 1994. Threatened, Endangered and
25 Sensitive Species of the Pike and San Isabel National Forests and Comanche and
26 Cimarron National Grasslands. May 25, 1994.
- 27 • Sauer, J.R., J.E. Hines, G. Gough, I. Thomas, and B.G. Peterjohn. 1997. The North
28 American Breeding Bird Survey Results and Analysis. Version 96.3. Online. Patuxent
29 Wildlife Research Center, Laurel, MD. Available: <http://www.mbr.nbs.gov/bbs/bbs.html>.
- 30 • Sauer, J.R., and S. Droege. 1992. Geographical patterns in population trends of
31 Neotropical migrants in North America. Pages 26-42 in J.M. Hagan, III, and D.W.
32 Johnston, editors. *Ecology and conservation of Neotropical migrant landbirds*.
33 Smithsonian Institution Press, Washington, DC.
- 34 • Sibley, C.G., and B.L. Monroe, Jr. 1990. Distribution and Taxonomy of Birds of the World.
35 Yale University Press, New Haven, CT. xxiv + 1111 pp.
- 36 • Smith, M.R., P.W. Mattocks, Jr., and K.M. Cassidy. 1997. Breeding birds of Washington
37 State. Vol. 4. IN Washington State Gap analysis - Final report (K.M. Cassidy, C.E. Grue.,
38 M.R. Smith, and K.M. Dvornich, eds.). Seattle Audubon Society Publications in Zoology
39 No. 1, Seattle, 538 pp.
- 40 • Stevenson, H.M., and B.H. Anderson. 1994. The Birdlife of Florida. University Press of
41 Florida, 891 pp.
- 42 • Stiles, F. G. and A. F. Skutch. 1989. A guide to the birds of Costa Rica. Cornell University
43 Press, Ithaca, New York, USA. 511 pp.
- 44 • Tate, J. 1981. The Blue List for 1981: the first decade. *American Birds* 35:3-10.

- 1 • Taylor, D. 1985. Candidate threatened or endangered molluscs in Box Canyon ACEC.
2 Gooding County, ID. Report to Bureau of Land Management, Shoshone, ID, under Order
3 ID-050-PH5-0093. 19 pp.
- 4 • Terres, J. K. 1980. The Audubon Society encyclopedia of North American birds. Alfred A.
5 Knopf, NY.
- 6 • U. S. Fish and Wildlife Service (USFWS). 2000. Endangered and threatened wildlife and
7 plants; notice of 90-day finding for a petition to list the yellow-billed cuckoo as
8 endangered and commencement of a status review. Notice of 90-day petition finding.
9 Federal Register 65(33): 8104-8107.
- 10 • U. S. Fish and Wildlife Service (USFWS). 2000. Notice of 90-day finding for a petition to
11 list the Yellow-billed Cuckoo as Endangered and commencement of a status review.
12 Federal Register 65(33): 8104-8107.
- 13 • U. S. Fish and Wildlife Service (USFWS). 2000. Notice of 90-day finding for a petition to
14 list the Yellow-billed Cuckoo as Endangered and commencement of a status review.
15 Federal Register 65(33):8104-8107.
- 16 • U. S. Fish and Wildlife Service (USFWS). 2001. Endangered and Threatened Wildlife and
17 Plants; Re-opening of the Public Comment Period for Status Review of the Yellow-Billed
18 Cuckoo in the Western United States. Federal Register 66(6): 1633-1634.
- 19 • USGS. 2002. Yellow-billed cuckoo breeding bird survey results. [www.mbr-](http://www.mbr-pwrc.usgs.gov/cgi-bin/atlasa01.pl?03870)
20 [pwrc.usgs.gov/cgi-bin/atlasa01.pl?03870](http://www.mbr-pwrc.usgs.gov/cgi-bin/atlasa01.pl?03870).
- 21 • Veit, R. R., and W. R. Peterson. 1993. Birds of Massachusetts. Massachusetts Audubon
22 Society, Lincoln, MA. xv + 514 pp.
- 23 • Walters, R. E., and E. Sorensen (eds.). 1983. Utah bird distribution: latilong study. Publ.
24 No. 83-10, Utah Division of Wildlife Resources, Salt Lake City. 97 pp.
- 25 • Wilcove, D.S. 1985. Nest predation in forest tracts and the decline of migratory
26 songbirds. Ecology, 66:1211-1214.
- 27 • Wilcove, D.S., C.H. McLellan, and A.P. Dobson. 1986. Habitat fragmentation in the
28 temperate zone. Pages 237-256 in M.E. Soule, editor. Conservation biology: the science
29 of scarcity and diversity. Sinauer Associates, Sunderland, MA.
- 30 • Wood, Merrill. 1979. Birds of Pennsylvania. Pennsylvania State Univ., University Park,
31 PA. 133pp.
- 32 • Woodbury, A. M., C. Cottam, and J. W. Sugden. 1949. Annotated check-list of the birds
33 of Utah. Bull. Univ. Utah 39(16) [Biol. Ser. 11(2)]: 1-40.
- 34 • Zook, J. L. 2002. Distribution maps of the birds of Nicaragua, Costa Rica, and Panama.
35 Unpublished.
- 36

Southwestern Willow Flycatcher (*Empidonax traillii extimus*)

The southwestern willow flycatcher was designated as a federally Endangered species on February 27, 1995.

Historic Range: BREEDS: southwestern U.S. (southern California north to Independence, Arizona, southwestern New Mexico, southern Utah, and, at least formerly, southern Nevada) and possibly northern Baja California and Sonora (very rare if present). Sedgwick (2001) studied distributional limits using distinctive song types of *E.T. extimus* and *E.T. adastus*, and found intergradation or overlap in southwestern Colorado and northwestern New Mexico. In areas of intergradation, there was some sorting of song types by elevation; birds with songs attributable to *E.T. extimus* were found as far north as 37 deg N at low elevation, whereas birds attributable to *E.T. adastus* were found as far south as 33.7 deg N at high elevation. The latter population occurred at over 2,400 meters in eastern Arizona. Occurred at least formerly in western Texas (current status uncertain) and northern Sonora. Some isolated remnant populations in southern California were allocated to subspecies *extimus* by Unitt (1987), but not by Phillips (1948). Population along the lower Colorado River now limited to about 20 pairs at Havasu National Wildlife Refuge (M. Romich, pers. comm. 2003). Formerly widespread in Arizona; now persist only in several small, widely scattered locations. Unitt (1987) noted that there was little recent information from Nevada and Utah. Unitt (1987) and USFWS (1993, 1995) included populations in areas of intergradation in the range of *E.T. extimus*. Winters: probably central Mexico to northwestern Colombia (Stiles and Skutch 1989). Migrates: in southern California, migrates through desert regions and sometimes along the coast and onto the Channel Islands (Biosystems Analysis 1989).

Basic Description: A small bird (flycatcher).

General Description: A flycatcher with brownish-olive upperparts, a whitish throat that contrasts with the pale olive breast, a pale yellow belly, and two light wing bars; generally lacks a conspicuous eye ring; as in other flycatchers, the bill is depressed and wide at the base (NGS 1983).

Diagnostic Characteristics: The palest subspecies of *E. traillii*, adults most closely resemble subspecies *adastus* but are even paler above, especially on the head, and *extimus* has a less pronounced chest band and the belly and crissum are paler yellow (Phillips 1948). Song differs from that of other subspecies by being a more protracted, slurred "fit-a-bew" with a burry "bew" syllable rather than a crisp, sneezy "fitz-bew" (USFWS 1995).

Reproduction Comments: Nesting occurs usually from early June through the end of July, peak in mid-June (Unitt 1987); sometimes may lay eggs as early as late May. In Grand Canyon, Arizona, breeds from early June to mid-July or perhaps early August (Brown 1988). Clutch size usually is 3-4 (2-3 along Colorado River). Incubation lasts 12-15 days, by female. Young are tended by both parents, leave nest at 12-15 days, usually in early to mid-July. Typically raises one brood per year. May incur a high rate of cowbird parasitism, especially in low elevation populations (e.g., Harris 1991, Brown 1988). Sometimes polygynous.

Ecology Comments: Breeding territories are about 1.5 acres. Densities may be on the order of 9-14 pairs/100 acres.

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: Yes

- 1 **Mobility and Migration Comments:** Present in California from late April to September
2 (Biosystems Analysis 1989), in southern Arizona from early May to early or mid-September
3 (Phillips et al. 1964). Arrives in Grand Canyon, Arizona, in mid-May (Brown, in Unitt 1987).
4 Spring migration peaks in mid-May; fall migration extends from mid-August to early September
5 (Biosystems Analysis 1989).
- 6 **Palustrine Habitat(s):** FORESTED WETLAND, Riparian
- 7 **Terrestrial Habitat(s):** Old field, Shrubland/chaparral, Woodland - Hardwood, Woodland - Mixed
- 8 **Habitat Comments:** Thickets, scrubby and brushy areas, open second growth, swamps, and
9 open woodland (AOU 1983). Restricted to riparian habitat in Arizona (Brown 1988). Nests
10 primarily in swampy thickets, especially of willow, sometimes buttonbush (Phillips et al. 1964,
11 AOU 1983), tamarisk (Brown 1988), vines, or other plants, where vegetation is 4-7 m or more in
12 height. Tamarisk is commonly used in the eastern part of the range. Habitat patches as small as
13 0.5 ha can support one or two nesting pairs (see USFWS 1995). Nests in fork or on horizontal
14 limb of small tree, shrub, or vine, at height of 0.6-6.4 m (mean usually about 2-3 m) (Harris 1991),
15 with dense vegetation above and around the nest.
- 16 **Adult Food Habits:** Invertivore
- 17 **Immature Food Habits:** Invertivore
- 18 **Food Comments:** Eats mainly insects caught in flight, sometimes gleans insects from foliage;
19 occasionally eats berries. In breeding range, forages within and occasionally above dense
20 riparian vegetation.
- 21 **Adult Phenology:** Diurnal
- 22 **Immature Phenology:** Diurnal
- 23 **Length:** 15 cm
- 24 **Weight:** 11 grams
- 25 **Management Requirements:** In Oregon, willow flycatcher populations increased after reduction
26 in cattle grazing and cessation of poisoning and removal of riparian willows (Taylor and Littlefield
27 1986). Harris (1991) recommended habitat restoration and reduction in grazing as the best long-
28 term management strategies for reducing the rate of cowbird parasitism; trapping of cowbirds or
29 removal of cowbird eggs may be useful short-term strategies to provide immediate relief to
30 critical populations. Brown (1988) cautioned against activities that would reduce or eliminate
31 tamarisk (nesting habitat) in Grand Canyon, Arizona, and recommended that water releases from
32 Glen Canyon dam be managed in such a way as to minimize streambank erosion and
33 consequent reduction in riparian breeding habitat. See USFWS (1995) for further information.
- 34 **Monitoring Requirements:** Those doing field surveys should be aware that subspecies
35 *brewsteri* is present (in migration) in the range of *extimus* during most of the latter's breeding
36 season; surveys should encompass the period June 20 to July 15 and include repeated visits to
37 verify that observed birds are resident and territorial (Unitt 1987).
- 38 **References:**
- 39 • American Ornithologists' Union (AOU). 1983. Check-list of North American Birds, 6th
40 edition. Allen Press, Inc., Lawrence, KS. 877 pp.

- 1 • Behle, W. H. 1943. Birds of Pine Valley Mountain region, southwestern Utah. Bull. Univ.
2 Utah 34(2), Biol. Ser. 7(5); 1-85.
- 3 • Behle, W. H. 1985. Utah birds: geographic distribution and systematics. Utah Mus. Natur.
4 Hist., Occas. Pap. 5: 1-147.
- 5 • Bent, A.C. 1942. Life histories of North American flycatchers, larks, swallows, and their
6 allies. U.S. National Museum Bulletin 179. Washington, DC.
- 7 • Biosystems Analysis, Inc. 1989. Endangered Species Alert Program Manual: Species
8 Accounts and Procedures. Southern California Edison Environmental Affairs Division.
- 9 • Brown, B. T. 1988. Breeding ecology of a willow flycatcher population in Grand Canyon,
10 Arizona. Western Birds 19:25-33.
- 11 • Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1992. Birds in Jeopardy: the Imperiled and
12 Extinct Birds of the United States and Canada, Including Hawaii and Puerto Rico.
13 Stanford University Press, Stanford, CA. 259 pp.
- 14 • Grinnell, J. 1914. An account of the mammals and birds of the lower Colorado valley, with
15 especial reference to the distributional problems presented. Univ. California Publication
16 Zoology 12:51-294.
- 17 • Harris, J. H. 1991. Effects of brood parasitism by brown-headed cowbirds on willow
18 flycatcher nesting success along the Kern River, California. Western Birds 22:13-26.
- 19 • Harrison, C. 1978. A Field Guide to the Nests, Eggs and Nestlings of North American
20 Birds. Collins, Cleveland, OH.
- 21 • Harrison, H. H. 1979. A field guide to western birds' nests. Houghton Mifflin Company,
22 Boston, MA. 279 pp.
- 23 • Johnson, M.J. and M.K. Sogge. 1997. Southwestern Willow Flycatcher surveys along
24 portions of the San Juan River, Utah (Montezuma Creek-Mexican Hat and Clay Hills
25 Crossing): 1997. Final report to the Utah Division of Wildlife Resources. Colorado Plateau
26 Research Station/Northern Arizona University, Flagstaff, AZ. 40 pp.
- 27 • Koronkiewicz, T., M. K. Sogge, and C. A. Drost. 1998. A preliminary survey for wintering
28 willow flycatchers in Costa Rica. USGS Forest and Rangeland Ecosystem Science
29 Center, Colorado Plateau Field Station, Northern Arizona University, Flagstaff. 47 pp.
- 30 • LaRue, C.T. 1994. Birds of northern Black Mesa, Navajo County, Arizona. Great Basin
31 Naturalist 54(1):1-63.
- 32 • Lowther, P. E. 1993. Brown-headed cowbird. North Amer. Birds 47: 1-24.
- 33 • Marshall, R. M. 2000. Population status on breeding grounds. Chapter 3 *In*: D. Finch and
34 S. Stoleson, editors. Status, ecology, and conservation of the southwestern willow
35 flycatcher. USDA Forest Service, Rocky Mountain Research Station, Albuquerque, NM.
- 36 • McCabe, R. A. 1991. The little green bird: ecology of the willow flycatcher. Rusty Rock
37 Press, Madison, WI. xv + 171 pp.
- 38 • McDonald, K. P., J. Snider, L. C. Peterson, M. St. Germain, and S. Staats. 1995. Results
39 of 1995 southwestern willow flycatcher surveys in the Virgin River drainage and southern
40 Utah. Utah Division of Wildlife Resources Publ. No. 95-17.
- 41 • McDonald, K. P., L. C. Peterson, and M. St. Germain. 1997. Results of 1996 surveys for
42 southwestern willow flycatchers in the upper Virgin River drainage and southern Utah.
43 Utah Division of Wildlife Resources Publ. No. 97-3. iii + 29 pp + 2 appendices.
- 44 • National Geographic Society. 1983. Field guide to the birds of North America. National
45 Geographic Society, Washington, DC.

- 1 • New Mexico Department of Game and Fish. 1995. Recommended Changes: List of
2 Endangered Species in New Mexico. pp. 1-12.
- 3 • Ohmart, R. D., B. W. Anderson, and W. C. Hunter. 1988. The ecology of the lower
4 Colorado River from Davis Dam to the Mexico-United States international boundary: a
5 community profile. U.S. Fish and Wildlife Service, Biological Report 85(7.19). xvi + 296
6 pp.
- 7 • Paxton, E. H. 2000. Molecular genetic structuring and demographic history of the willow
8 flycatcher (*Empidonax traillii*). Master's Thesis. Northern Arizona University, Flagstaff. 42
9 pp.
- 10 • Phillips, A. R. 1948. Geographic variation in *Empidonax traillii*. Auk 65:507-514.
- 11 • Phillips, A., J. Marshall, and G. Monson. 1964. The birds of Arizona. The University of
12 Arizona Press, Tucson, AZ.
- 13 • Rosenberg, K.V., R.D. Ohmart, W.C. Hunter, and B.W. Anderson. 1990. Birds of the
14 lower Colorado River valley. University of Arizona Press, Tucson, AZ. 464 pp.
- 15 • Schreier, W. 1996. Bryce Canyon National Park 1996 endangered avian species survey
16 report--southwestern willow flycatcher. National Park Service, Colorado Plateau System
17 Support Office, Intermountain Field Area, Bryce canyon, UT. 21 pp.
- 18 • Sedgwick, J. A. 2001. Geographic variation in the song of Willow Flycatchers:
19 differentiation between *Empidonax traillii adastus* and *E. T. extimus*. Auk. 118:366-379.
- 20 • Sogge, M. K. 1995. Southwestern willow flycatcher surveys along the San Juan River,
21 Utah[:] 1994-1995. Final report to the Bureau of Land Management, San Juan Resource
22 Area. National Biological Service Colorado Plateau Research Station/Northern Arizona
23 University report. ii + 27 pp.
- 24 • Sogge, M. K., R. M. Marshall, S. J. Sferra, and T. J. Tibbets. 1997. A southwestern willow
25 flycatcher natural history summary and survey protocol. USDI National Park Service,
26 Colorado Plateau Research Station, Technical Report NPS/NAUCPRS/NRTR-97/12,
27 Flagstaff, AZ.
- 28 • Sogge, M. K., S. J. Sferra, T. D. McCarthey, S. O. Williams, III, and B. E. Kus. 2001.
29 Summary of southwestern willow flycatcher breeding site and territory characteristics -
30 1999. Prepared for the Southwestern Willow Flycatcher Recovery Team, U.S. Fish and
31 Wildlife Service, Region 2, Albuquerque, NM.
- 32 • Sogge, M. K., S. J. Sferra, T. McCarthey, S. O. Williams and B. E. Kus. 2002.
33 Southwestern W illow Flycatcher breeding site and territory summary - 2001. USGS
34 Forest and Rangeland Ecosystem Science Center, Colorado Plateau Field Station report
35 to the U.S. Fish and Wildlife Service Southwestern Willow Flycatcher Recovery Team.
- 36 • Sogge, M.K., et.al. 1995. Status of the southwestern willow flycatcher along the Colorado
37 River in Grand Canyon National Park - 1995. Summary Report. National Biological
38 Service Colorado Plateau Research Station/Northern Arizona University. 26 p.
- 39 • Spahr, R., L. Armstrong, D. Atwood, and M. Rath. 1991. Threatened, endangered, and
40 sensitive species of the Intermountain Region. U.S. Forest Service, Ogden, UT.
- 41 • Stiles, F. G. and A. F. Skutch. 1989. A guide to the birds of Costa Rica. Cornell University
42 Press, Ithaca, NY, USA. 511 pp.
- 43 • Taylor, D. M., and C. D. Littlefield. 1986. Willow flycatcher and yellow warbler response to
44 cattle grazing. American Birds 40:1169-1173.
- 45 • Terres, J. K. 1980. The Audubon Society encyclopedia of North American birds. Alfred A.
46 Knopf, NY.

- 1 • Tibbits, T.J. et.al. 1994. A Survey Protocol for the Southwestern Willow Flycatcher
2 (*Empidonax traillii extimus*). U.S. National Park Service Technical Report
3 NPS/NAUCPRS/NRTR-94/04, Colorado Plateau Research Station at Northern Arizona
4 University. 24 p.
- 5 • U.S. Fish and Wildlife Service. 1993. Proposed rule to list the southwestern willow
6 flycatcher as endangered with critical habitat. Federal Register 58(140):39495-39522. 23
7 July 1993.
- 8 • U.S. Fish and Wildlife Service (USFWS). 1989. Endangered and threatened wildlife and
9 plants; animal notice of review. Federal Register, Department of the Interior 54(4): 554-
10 579.
- 11 • U.S. Fish and Wildlife Service (USFWS). 2002. Southwestern Willow Flycatcher
12 (*Empidonax traillii extimus*). Final recovery plan. Prepared by Southwestern Willow
13 Flycatcher Recovery Team Technical Subgroup. U.S. Fish and Wildlife Service, Division
14 of Ecological Services, Albuquerque, NM.
- 15 • Unitt, P. 1987. *Empidonax traillii extimus*: an endangered subspecies. Western Birds
16 18:137-162.
- 17 • Woodbury, A. M., and H. N. Russell, Jr. 1945. Birds of the Navajo country. Univ. Utah
18 Bull. 35(14), Biol. Ser. 9(1): 1-157.
- 19 • Yeager, B. and B. Masslich. 1995. Biological assessment for the St. George road
20 projects. Prepared for Utah Department of Transportation. Bio/West, Inc., 1063 West
21 1400 North, Logan, UT 84321.

Northern Aplomado Falcon (*Falco femoralis*)

The northern aplomado falcon was designated as a federally Threatened species with a nonessential experimental population on February 25, 1986.

Historic Range: Historic breeding range: southeastern Arizona, southern New Mexico, and southern Texas south through Mexico (Tamaulipas, Chiapas, Campeche, Tabasco, Chihuahua, Coahuila, Sinaloa, Jalisco, Guerrero, Veracruz, Yucatan, and San Luis Potosi) to Guatemala (Pacific slope of Central American cordillera). Last verified breeding in the U.S. was in New Mexico in 1952 and in Texas in 1941 and 1995; unconfirmed report from Arizona in the late 1960s (AOU 1983); reintroduction is underway. Nests regularly only along Gulf Coast of Mexico in portions of northern and central Veracruz, northern Chiapas, western Campeche, and eastern Tabasco (Matthews and Moseley 1990). Unbanded individuals were recorded in New Mexico and Texas in the early 1990s. Historic winter range: Sinaloa, Chihuahua, and southern Tamaulipas south to southern Mexico; casual in Guatemala (AOU 1957).

Basic Description: A falcon.

Reproduction Comments: Egg-laying: January-June (mainly March-May, peak in April). Clutch size typically is 2-3. Both parents (mainly female) incubate, about 31-32 days (Cade 1982, Evans 1982). Young can fly at 4-5 weeks, may remain in nest area for several weeks more. Pairs remain together throughout the year (Palmer 1988).

Non-Migrant: Yes

Locally Migrant: Yes

Long Distance Migrant: No

Palustrine Habitat(s): Riparian

Terrestrial Habitat(s): Grassland/herbaceous, Savanna, Woodland – Conifer

Habitat Comments: Open rangeland and savanna, semiarid grasslands with scattered trees and shrubs; in U.S., was found in coastal prairies along sand ridges, in woodlands along desert streams, and in desert grasslands with scattered mesquite and yucca; has been found in open pine woodland in central Mexico (Matthews and Moseley 1990, Johnsgard 1990). Encroachment of thick tall grass or brush degrades habitat. Nests in old stick nests of other bird species (e.g., hawks, caracaras, ravens); in sites such as bromeliads in tropics. May sometimes nest on cliff.

Adult Food Habits: Carnivore, Invertivore

Immature Food Habits: Carnivore, Invertivore

Food Comments: Feeds primarily on birds (up to rock dove size), to a lesser extent on insects (moths, beetles, cicadas, orthopterans); uncommonly on small mammals, lizards, and snakes (Terres 1980, Cade 1982). Pairs often hunt together. Birds comprise most of diet biomass in eastern Mexico, but insects also are commonly consumed. Hunts from perch or air. See Palmer (1988) for further details. In eastern Mexico, hunted mainly within 1 km of nest site (Hector 1988).

Adult Phenology: Crepuscular, Diurnal

Immature Phenology: Crepuscular, Diurnal

Phenology Comments: Decidedly crepuscular in hunting habits, often catching prey after sunset; not very active in middle of day (Cade 1982). In eastern Mexico, preyed on birds mainly in the early morning, hawked insects later in the day (see Johnsgard 1990).

Length: 45 cm

Weight: 410 grams

References:

- Allen, Craig R., Stephen Demarais, and R. Scott Lutz. 1994. Red imported fire ant impact on wildlife: an overview. *Texas J. Sci.* 46(1):51-59.
- American Ornithologists' Union (AOU). 1957. *The A.O.U. Check-list of North American Birds*, 5th ed. Port City Press, Inc., Baltimore, MD. 691 pp.
- American Ornithologists' Union (AOU). 1983. *Check-list of North American Birds*, 6th edition. Allen Press, Inc., Lawrence, KS. 877 pp.
- Cade, T.J. 1982. *The falcons of the world*. Cornell University Press, Ithaca, NY. 192 pp.
- Campbell, L. 1995. *Endangered and Threatened Animals of Texas: Their Life History and Management*. Texas Parks and Wildlife Department, Endangered Resources Branch, Austin, TX. ix + 129 pp.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1992. *Birds in Jeopardy: the Imperiled and Extinct Birds of the United States and Canada, Including Hawaii and Puerto Rico*. Stanford University Press, Stanford, CA. 259 pp.
- Evans, D. L. 1982. Status reports on twelve raptors. U.S. Department of the Interior, Fish and Wildlife Service, Special Scientific Report No. 238. 68 pp.
- Hector, D. P. 1987. Decline of the aplomado falcon, U.S.A. *American Birds* 41:381-389.
- Hector, D. P. K. 1988. Vegetative cover, small bird abundance and patterns of aplomado falcon habitat quality in eastern Mexico. Pages 157-164 in Glinski et al., eds. *Proc. Southwest raptor management symposium and workshop*. Nat. Wildl. Fed. Sci. and Tech. Ser. No. 11.
- Johnsgard, P. A. 1990. *Hawks, eagles, and falcons of North America*. Smithsonian Inst. Press, Washington, D.C. xvi + 403 pp.
- Matthews, J.R. and C.J. Moseley (eds.). 1990. *The Official World Wildlife Fund Guide to Endangered Species of North America*. Volume 1. Plants, Mammals. xxiii + pp 1-560 + 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians, Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180. Beacham Publications, Inc., Washington, D.C.
- Palmer, R. S., ed. 1988. *Handbook of North American birds*. Vol. 5. Yale Univ. Press, New Haven. 465 pp.
- Terres, J. K. 1980. *The Audubon Society encyclopedia of North American birds*. Alfred A. Knopf, NY.

American Peregrine Falcon (*Falco peregrinus anatum*)

Historic Range: BREEDS: Across interior Alaska, south of the Brooks Range southeastward across Canada to Labrador, and south to Baja California and northern Mexico (Palmer 1988, Ambrose et al. 1988, Rowell 2002). Replaced on the coast of Alaska and outer coast of British Columbia by *F. p. pealei*. WINTERS: Those breeding in the boreal subarctic winter in South America; those at more southern latitudes exhibit variable migration behavior, and some are nonmigratory (USFWS 1999).

Basic Description: A medium-sized falcon.

General Description: A falcon with long pointed wings, a dark crown and nape, and a dark wedge extending below the eye; forehead is pale in immature, which are mainly brownish above rather than black or gray as in adults (NGS 1983).

Diagnostic Characteristics: Intermediate in coloration between the pale birds of the arctic (subspecies *tundrius*) and the very dark pergrines of the northwest coast of North America (subspecies *pealei*).

Reproduction Comments: Clutch size averages 4 at mid-latitudes, 3 in far north. Incubation lasts 32-35 days, mainly by female (male brings food). Young fledge at 39-49 days, gradually become independent. First breeds usually at 2-3 years, occasionally as yearling. Usually lifelong pair bond. Replaces lost clutches, usually at alternate site. Brood losses apparently caused mainly by bad weather. See many further details in Palmer (1988). In northwestern Arizona, mean distance between centers of nesting areas was around 6-8 km (Brown et al. 1992).

Ecology Comments: Great-horned Owl may be a serious nest predator in the U.S. Severe weather may result in high mortality in far north. Foraging range up to 27 kilometers (Martin 1979); home ranges in Great Britain varied from 44-65 square kilometers, and averaged 52 square kilometers (Brown and Amadon 1968). In Utah, home range radii varied from 0.3 to 29.8 kilometers, average 12.2 km (n = 19; Porter and White 1973).

Non-Migrant: No

Locally Migrant: No

Long Distance Migrant: Yes

Mobility and Migration Comments: Populations nesting in northern latitudes are highly migratory; those nesting in northern maritime climates, at mid-latitudes, and in the Southern Hemisphere much less so (Cade 1982). Tundra breeders migrate farthest, bypassing those farther south; a few winter in Florida, some in Caribbean, perhaps some in Central America, most in southern South America (Palmer 1988). Breeders from central Alaska migrated through central North America and wintered in southern Mexico, Central America, the Caribbean region, and South America (Britten et al. 1995). Two breeders from southern Utah migrated through western Mexico, and one continued to a wintering site in Nicaragua (Britten et al. 1995). In the U.S., the Atlantic coast from New Jersey to South Carolina and the barrier islands of the Texas Gulf Coast are important feeding areas for long-distance migrants. Arrives in northern breeding areas late April-early May; departure begins late August-early September (Johnson and Herter 1989). See Palmer (1988) for further information on timing of migration. From Padre Island, Texas, a northbound migrant reached south-central Canada in four days, and a southbound migrant passed through Mexico and reached Guatemala in six days (Chavez-Ramirez et al. 1994).

Estuarine Habitat(s): Bay/sound, Herbaceous wetland, Lagoon, River mouth/tidal river, Tidal flat/shore

Terrestrial Habitat(s): Bare rock/talus/scree, Cliff, Shrubland/chaparral, Urban/edificarian, Woodland - Conifer, Woodland - Hardwood, Woodland - Mixed

Habitat Comments: Various open situations from tundra, moorlands, steppe, and seacoasts, especially where there are suitable nesting cliffs, to mountains, open forested regions, and human population centers (AOU 1983). When not breeding, occurs in areas where prey concentrate, including farmlands, marshes, lakeshores, river mouths, tidal flats, dunes and beaches, broad river valleys, cities, and airports. Often nests on ledge or hole on face of rocky cliff or crag. River banks, tundra mounds, open bogs, large stick nests of other species, tree hollows, and man-made structures (e.g., ledges of city buildings) are used locally (Cade 1982). Nests typically are situated on ledges of vertical rocky cliffs, commonly with a sheltering overhang (Palmer 1988, Campbell et al 1990). Tundra populations nests typically on rocky cliffs, bluffs, or dirt banks. Ideal locations include undisturbed areas with a wide view, near water, and close to plentiful prey. Substitute man-made sites include tall buildings, bridges, rock quarries, and raised platforms. See Grebence and White (1989) for information on nesting along the Colorado River system.

Adult Food Habits: Carnivore

Immature Food Habits: Carnivore

Food Comments: Feeds primarily on birds (medium-size passerines up to small waterfowl); rarely or locally, small mammals (e.g., bats, lemmings), lizards, fishes, and insects (by young birds) may be taken. Prey pursuit initiated from perch or while soaring. May hunt up to several km from nest site (Skaggs et al. 1988). See Rosenfield et al. (1995) for information on food habits in Greenland.

Adult Phenology: Diurnal

Immature Phenology: Diurnal

Phenology Comments: In general, much hunting occurs in morning, and to lesser extent toward evening, but may hunt anytime during day.

Length: 51 cm

Weight: 1500 grams

References:

- Allen, Craig R., Stephen Demarais, and R. Scott Lutz. 1994. Red imported fire ant impact on wildlife: an overview. *Texas J. Sci.* 46(1):51-59.
- Ambrose, R.E., R.J. Ritchie, C.M. White, P.F. Schempf, T. Swem and R. Dittrock. 1988. Changes in the status of peregrine falcon populations in Alaska. Pp. 73-82. In: Cade, T.J., J.H. Enderson, C.G. Thelander and C.M. White (eds.). *Peregrine falcon populations: their management and recovery*. The Peregrine Fund, Inc., Boise, ID.
- Ambrose, S. 1993. Phone conversation with E. West, AKNHP zoologist regarding population level of F.P. Anatum. 4 October 1993. Ecological Services, U.S. Fish and Wildlife Service, Fairbanks AK.
- American Ornithologists' Union (AOU). 1957. *The A.O.U. Check-list of North American Birds*, 5th ed. Port City Press, Inc., Baltimore, MD. 691 pp.
- American Ornithologists' Union (AOU). 1983. *Check-list of North American Birds*, 6th edition. Allen Press, Inc., Lawrence, KS. 877 pp.

- 1 • Andrews, R. R. and R. R. Righter. 1992. Colorado Birds. Denver Museum of Natural
2 History, Denver. 442 pp.
- 3 • Audubon Society. 1981-1985. Breeding Bird Atlas of New Hampshire. (unpublished).
- 4 • Banasch, U., J. P. Goossen, A. E. Riez, C. Casler, and R. D. Barradas. 1992.
5 Organochlorine contaminants in migrant and resident prey of peregrine falcons, *Falco*
6 *peregrinus*, in Panama, Venezuela, and Mexico. Canadian Field-Naturalist 106:493-498.
- 7 • Bechard, M. J., D. Beig, and R. P. Howard. 1987. Historical nest sites of the peregrine
8 falcon (*Falco peregrinus anatum*) in Idaho. Unpublished report for U. S. Fish and Wildlife
9 Service. 34 pp.
- 10 • Berger, D.D., C.R. Sindelar, JR. and K.E. Gamble. 1969. The Status of Breeding
11 Peregrines in the Eastern United States. P.165-173. In: J.J. Hickey (Ed.). Peregrin Falcon
12 Populations Their Biology and Decline. University of Wisconsin Press, Madison. 596 p.
- 13 • Bollengier, R.M., JR. (Team Leader). 1979. Eastern Peregrine Falcon Recovery Plan.
14 U.S. Fish and Wildlife Service. 147 pp.
- 15 • Brown, B. T., et al. 1992. Density of nesting peregrine falcons in Grand Canyon National
16 Park, Arizona. Southwestern Naturalist 37:188-193.
- 17 • Brown, L. and D. Amadon. 1968. Eagles, Hawks, and Falcons of the World. McGraw-Hill,
18 NY.
- 19 • Burleigh, T. D. 1972. Birds of Idaho. The Caxton Printers, Ltd., Caldwell, Idaho. 467 pp.
- 20 • Committee on the Status of Endangered Wildlife in Canada (COSEWIC). n.d. Status
21 Report on Endangered Wildlife in Canada: Peregrine Falcon. COSEWIC. 48 pp.
- 22 • Cade, T. J., et al., eds. 1988. Peregrine falcon populations: their management and
23 recovery. The Peregrine Fund, Inc., Boise, ID. 949 pp.
- 24 • Cade, T.J. 1982. The falcons of the world. Cornell University Press, Ithaca, NY. 192 pp.
- 25 • California Department of Fish and Game (CDF&G). 1990. 1989 annual report on the
26 status of California's state listed threatened and endangered plants and animals. 188 pp.
- 27 • Campbell, R.W., N.K. Dawe, I. McTaggart-Cowan, J.M. Cooper, G.W. Kaiser, and M.C.E.
28 McNall. 1990. The Birds of British Columbia. Vol. 1. Nonpasserines: introduction and
29 loons through waterfowl. Royal British Columbia Museum, Victoria, British Columbia,
30 Canada. 514 pp.
- 31 • Cannings, S. 2001. EO Specifications for Peregrine Falcon (*Falco peregrinus*).
32 NatureServe, Unpublished. 1 pp.
- 33 • Colorado Bird Observatory. 1997. 1996 Reference Guide to the Monitoring and
34 Conservation Status of Colorado's Breeding Birds. Colorado Bird Observatory, Colorado
35 Division of Wildlife, Great Outdoors Colorado Trust Fund, and Partners, March 21, 1997.
- 36 • Court, G.S. 1993. A toxicological assessment of the American peregrine falcon (*Falco*
37 *peregrinus anatum*) breeding in Alberta, Canada - 1968 to 1992. Occasional Paper
38 Number 10, Department of Environmental Protection, Wildlife Management Division, Fish
39 and Wildlife Services, Government of Alberta. 28 pp.
- 40 • Craig, Gerald R. 1997. Recommended buffer zones and seasonal restrictions for
41 Colorado raptor nests. Unpublished report for Colorado Division of Wildlife. Updated
42 January 6, 1997.
- 43 • DeGraaf, R. M. and D. R. Rudis. 1986. New England wildlife: habitat, natural history, and
44 distribution. Univ. Mass. Press. Amherst, MA. 491 pp.

- 1 • Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1992. Birds in Jeopardy: the Imperiled and
2 Extinct Birds of the United States and Canada, Including Hawaii and Puerto Rico.
3 Stanford University Press, Stanford, CA. 259 pp.
- 4 • Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1992. Birds in jeopardy: the imperiled and
5 extinct birds of the United States and Canada, Including Hawaii and Puerto Rico.
6 Stanford University Press, Stanford, CA. 259 pp.
- 7 • Fuller, M. R., C. J. Henny, and P. B. Wood. 1995. Raptors. Pages 65-69 in E. T. LaRoe,
8 G. S. Farris, C. E. Puckett, P. D. Doran, and M. J. Mac, editors. Our living resources: A
9 report to the nation on the distribution, abundance, and health of U.S. plants, animals and
10 ecosystems. U.S. Department of Interior, National Biological Service, Washington, D.C.
- 11 • Godfrey, W.E. 1986. The Birds of Canada. Revised edition. National Museum of Natural
12 Sciences, Ottawa, Canada. 595 pp.
- 13 • Grebence, B. L., and C. M. White. 1989. Physiographic characteristics of peregrine falcon
14 nesting habitat along the Colorado River system in Utah. Great Basin Naturalist 49:408-
15 418.
- 16 • Hayes, G.E. and J.B. Buchanan. 2002. Washington State status report for the peregrine
17 falcon. Washington Dpt. Fish and Wildlife. Olympia, WA. 77 pp.
- 18 • Hickey, J. J. No date. Natural History of the Peregrine Falcon East of the Rockies.
19 Unpublished Manuscript. 178 pp.
- 20 • Holroyd, G. 1993. Phone conversation with E. West, AKNHP zoologist, regarding
21 population estimated for *Falco peregrinus tundrius* and *F.p. anatum* in Canada. 8 October
22 1993. Canadian Wildlife Service, Edmonton, Alberta.
- 23 • Holroyd, G. L., and U. Banasch. 1990. The reintroduction of the peregrine falcon, *Falco*
24 *peregrinus anatum*, into southern Canada. Canadian Field-Naturalist 104:203-208.
- 25 • Hubbard, J. P., and C. G. Schmitt. 1988. Organochlorine residues in avian prey of
26 peregrine falcons breeding in New Mexico. Pages 176-181 in Glinski et al., eds. Proc.
27 Southwest raptor management symposium and workshop. Nat. Wildl. Fed. Sci. and Tech.
28 Ser. No. 11.
- 29 • Jarvis, W. 1871. The Duck Hawk. Amer. Nat., 5:662.
- 30 • Jewett, S.G., W.P. Taylor, W.T. Shaw, and J.W. Aldrich. 1953. Birds of Washington
31 State. U. Washington Press. 767 pp.
- 32 • Johnsgard, P. A. 1986. Birds of the Rocky Mountains with particular reference to national
33 parks in the Northern Rocky Mountain region. Colorado Associated University Press,
34 Boulder. xi + 504 pp.
- 35 • Johnsgard, P. A. 1990. Hawks, eagles, and falcons of North America. Smithsonian Inst.
36 Press, Washington, D.C. xvi + 403 pp.
- 37 • Johnson, S. R. and D. R. Herter. 1989. The Birds of the Beaufort Sea. BP Exploration
38 (Alaska) Inc., Anchorage, AK. 372 pp.
- 39 • Johnson, T. H. 1988. Responses of breeding peregrine falcons to human stimuli. Pages
40 301-305 in Glinski et al., eds. Proc. Southwest raptor management symposium and
41 workshop. Nat. Wildl. Fed. Sci. and Tech. Ser. No. 11.
- 42 • Johnstone, R.M. 1997. Update of Status Report on the American Peregrine Falcon (*Falco*
43 *peregrinus anatum*). Department of Veterinary Anatomy, Western College of Veterinary
44 Medicine, University of Saskatchewan, Saskatoon, SK. 42 pp.
- 45 • Johnstone, R.M. 1999. Update COSEWIC Status Report on Anatum Peregrine Falcon
46 (*Falco peregrinus anatum*). Unpublished Report Prepared for the Committee on the
47 Status of Endangered Wildlife in Canada. 31 pp.

- 1 • King, W. B., compiler. 1979. Endangered birds of the world. The International Council for
2 Bird Preservation. Smithsonian Institution Press, Washington, D.C. [Reprinted in
3 handbook form in 1981.]
- 4 • Larrison, E. J., J. L. Tucker, and M. T. Jollie. 1967. Guide to Idaho birds. Journal of the
5 Idaho Academy of Science 5: 1-220.
- 6 • Lefranc, M. N., Jr., and R. L. Glinski. 1988. Southwest raptor management issues and
7 recommendations. Pages 375-392 in Glinski et al., eds. Proc. Southwest raptor
8 management symposium and workshop. National Wildlife Federation Science and Tech.
9 Ser. No. 11.
- 10 • Martin, M. 1979. Status report on peregrine falcon (*Falco peregrinus*) in Canada.
11 COSEWIC Report. B.C. Fish and Wildlife Branch, Victoria, B.C.
- 12 • National Geographic Society (NGS). 1983. Field guide to the birds of North America.
13 National Geographic Society, Washington, DC.
- 14 • National Geographic Society, 1987. Field Guide to the birds of North America, second
15 edition. The National Geographic Society, Washington, D.C.
- 16 • Nelson, D. 1993. Colorado Bird Atlas: Manual on Use of Breeding Codes. Denver
17 Museum of Natural History, Denver. 27 pp.
- 18 • New Hampshire Natural Heritage Inventory. Element File. Contact the New Hampshire
19 Natural Heritage Inventory: 603/271-3623.
- 20 • Palmer, R. S. (ed.). 1988. Handbook of North American Birds. Vol. 5, Diurnal raptors,
21 Part 2. Yale Univ. Press, New Haven, CT. 465 pp.
- 22 • Palmer, R. S., ed. 1988. Handbook of North American birds. Vol. 5. Yale Univ. Press,
23 New Haven. 465 pp.
- 24 • Peakall, D. B. 1990. Prospects for the peregrine falcon, *Falco peregrinus*, in the nineties.
25 Canadian Field-Naturalist 104:168-173.
- 26 • Porter, R. D. and C. M. White. 1973. Peregrine Falcon in Utah, emphasizing ecology and
27 competition with the Prairie Falcon. Brigham Young University Science Bulletin, Biol.
28 Series 18:1-74.
- 29 • Ratcliff, B. 2000. Peregrine Falcon Site Records. Unpublished Report Prepared by
30 Ontario Ministry of Natural Resources, North West Region. n.p.
- 31 • Rowell, P. A. 2002. Update COSEWIC status report on Anatum Peregrine Falcon (*Falco*
32 *peregrinus anatum*). Report submitted to the Committee on the Status of Endangered
33 Wildlife in Canada. Environment Canada, Ottawa. 23 pages + appendices.
- 34 • Rowell, Petra A. 2002. Update COSEWIC Status Report on Anatum Peregrin Falcon,
35 *Falco peregrinus anatum* (interim report 2nd draft, submitted April 2002). Committee on
36 the Status of Endangered Wildlife in Canada (COSEWIC) interim draft, unpublished. 29
37 pp.
- 38 • Sherrod, S. K., et al. 1982. Hacking: a method for releasing peregrine falcons and other
39 birds of prey. Second edition. The Peregrine Fund, Cornell Univ., Ithaca, NY. vi + 61 pp.
- 40 • Skaggs, R. W., et al. 1988. Peregrine falcon. Pages 127-136 in Glinski et al., eds. Proc.
41 Southwest raptor management symposium and workshop. Natural Wildlife Fed. Science
42 and Tech. Ser. No. 11.
- 43 • Smith, M.R., P.W. Mattocks, Jr., and K.M. Cassidy. 1997. Breeding birds of Washington
44 State. Vol. 4. IN Washington State Gap analysis - Final report (K.M. Cassidy, C.E. Grue.,
45 M.R. Smith, and K.M. Dvornich, eds.). Seattle Audubon Society Publications in Zoology
46 No. 1, Seattle, 538 pp.

- 1 • The Peregrine Fund. 1992. Peregrine falcon recovery program: status and
2 reommendations. Unpublished report.
- 3 • U. S. Fish and Wildlife Service. 1999. Final rule to remove the American Peregrine
4 Falcon from the federal list of endangered and threatened wildlife, and to remove the
5 similarity of appearance provision for free-flying Peregrines in the conterminous United
6 States. Federal Register 64 (164):46542-46558.
- 7 • U.S. Fish and Wildlife Service (USFWS). 1990. Endangered and threatened species
8 recovery program: report to Congress. 406 pp.
- 9 • U.S. Fish and Wildlife Service. 1980. Selected vertebrate endangered species of the
10 seacoast of the United States: American peregrine falcon. Tech. rep. FWS/OBS-
11 80/01.57. U.S. Dept. Interior, Fish and Wildlife Service, Biological Services Program,
12 Coastal Ecosystems Project, Slidell, LA. 14 p.
- 13 • U.S. Fish and Wildlife Service. 1991. Request for information on the Arctic and American
14 peregrine falcons. Federal Register 56(113):26969-26971.
- 15 • USFWS. n.d.. American Peregrine Falcon Proposed Monitoring Plan. U.S. Fish and
16 Wildlife Service. 5 pp.
- 17 • World Wildlife Fund. 1990. The official World Wildlife Fund guide to endangered species
18 of North America. D. W. Lowe, J. R. Matthews, and C. J. Moseley (eds.). Beacham
19 Publishing, Inc. Washington, D.C.
- 20

Bald Eagle (*Haliaeetus leucocephalus*)

Historic Range: BREEDING: central Alaska, northern Yukon, northwestern and southern Mackenzie, northern Saskatchewan, northern Manitoba, central Ontario, central Quebec, Labrador, and Newfoundland, south locally to the Commander and Aleutian Islands, southern Alaska, Baja California (both coasts), Sonora (Brown et al. 1988), New Mexico, Arizona, Texas Gulf Coast, and Florida (including the Keys); very local in Great Basin and prairie and plains regions in interior North America, where breeding range recently has expanded to include Nebraska and Kansas. NON-BREEDING: generally throughout the breeding range except in the far north (AOU 1983, Sibley and Monroe 1990), most commonly from southern Alaska and southern Canada southward. The Chilkat Bald Eagle Preserve, Alaska, supports the largest wintering population anywhere (Ehrlich et al. 1992). Winter concentrations occur in British Columbia-northwestern Washington, along the Missouri and Mississippi rivers, and in northern Arkansas. One of the largest fall (mid-October to mid-December) migrant concentrations (200-300 birds at any one time, close to a thousand individuals through the season) occurs at Hauser Lake near Helena, Montana.

Basic Description: Bald eagle. Mature adults have a white head and tail.

General Description: Adults have a white head, white tail, and a large bright yellow bill; elsewhere the plumage is dark. Immatures are dark with variable amounts of light splotching on the body, underwing coverts, flight feathers, and tail base; averages 79-94 cm long, 178-229 cm wingspan (NGS 1983).

Diagnostic Characteristics: Adults differ from other eagles in having both a white head and white tail (head of white-tailed eagle may look white at a distance). Bald eagle has a proportionately larger head and bill than does the golden eagle, in the immatures of which the white is confined to the base of the primaries and the base of the tail. Bald eagle lacks the long wedge-shaped tail of Steller's sea-eagle. Bald eagle's neck is shorter and tail is longer than in white-tailed eagle.

Reproduction Comments: Clutch size is 1-3 (usually 2). Incubation lasts about 5 weeks, by both sexes. Second hatched young often dies. Young first fly at 10-12.5 weeks, cared for by adults and may remain around nest for several weeks after fledging. Generally first breeds at about 5-6 years. Adults may not lay every year.

Ecology Comments: Commonly roosts communally, especially in winter. See Curnutt (1992) for information on the dynamics of a year-round communal roost in southern Florida. In Montana, the introduction of shrimp (*Mysis relicta*) had a cascading effect through the food chain, ultimately causing displacement of bald eagles (Spencer et al. 1991).

Non-Migrant: Yes

Locally Migrant: Yes

Long Distance Migrant: Yes

Mobility and Migration Comments: Most eagles that breed in Canada and the northern U.S. move south for winter. Migrates widely over most of North America (AOU 1983); moves generally E-SE across Canada and the Great Lakes region to the northeast coast of the U.S. In the northern Chesapeake Bay region, radio-tagged northern migrants arrived in late fall (mean date 21 December) and departed in early spring (mean date 27 March); radio-tagged southern migrants arrived throughout April-August and departed June-October (Buehler et al. 1991). See Palmer (1988) for fairly detailed review of seasonal movements in various regions. Defended territories are relatively small; 14 in Alaska varied from 11-45 hectares and averaged 23 ha (Hensel and Troyer 1964), and territory radius around active nests averaged 0.6 km in Minnesota

(Mahaffy and Frenzel 1987). Feeding home ranges surrounding active nests are undoubtedly much larger, depending on proximity to food sources and abundance of food. Minimum home range of breeding birds in Saskatchewan was 7 k² (Gerrard et al. 1992); on the Columbia River, Oregon, breeding home ranges averaged 21.6 k² (Garrett et al. 1993). Winter home ranges can be very large, especially for nonbreeding birds. An immature wintered in Arizona over an area of >40,000 k² and spent the summer in the Northwest Territories over a summer range of >55,000 k² (Grubb et al. 1994). Maximum distance between feeding area and night roost site was less than 16 km in winter in Missouri (Griffin et al. 1982). In north-central Arizona, February–April home range of immatures averaged 400 k²; birds moved frequently and roosted singly or in small groups (Grubb et al. 1989).

Marine Habitat(s): Near shore

Estuarine Habitat(s): Bay/sound, Lagoon, River mouth/tidal river, Tidal flat/shore

Riverine Habitat(s): BIG RIVER, MEDIUM RIVER

Lacustrine Habitat(s): Deep water, Shallow water

Palustrine Habitat(s): FORESTED WETLAND, Riparian

Terrestrial Habitat(s): Cliff, Forest - Conifer, Forest - Hardwood, Forest - Mixed, Woodland - Conifer, Woodland - Hardwood, Woodland - Mixed

Special Habitat Factors: Standing snag/hollow tree

Habitat Comments: Breeding habitat most commonly includes areas close to (within 4.0 km) coastal areas, bays, rivers, lakes, or other bodies of water that reflect the general availability of primary food sources including fish, waterfowl, and seabirds (Andrew and Mosher 1982, Green 1985, Campbell et al. 1990). Preferentially roosts in conifers or other sheltered sites in winter in some areas; typically selects the larger, more accessible trees (Buehler et al. 1991, 1992). Perching in deciduous and coniferous trees is equally common in other areas (e.g., Bowerman et al. 1993). Communal roost sites used by two or more eagles are common, and some may be used by 100 or more eagles during periods of high use. Winter roost sites vary in their proximity to food resources (up to 33 km) and may be determined to some extent by a preference for a warmer microclimate at these sites. Available data indicate that energy conservation may or may not be an important factor in roost-site selection (Buehler et al. 1991). In Saskatchewan lakes, density was positively correlated with abundance of large fishes (Dzus and Gerrard 1993). In winter, may associate with waterfowl concentrations or congregate in areas with abundant dead fish (Griffin et al. 1982); often roosts communally at night in trees that are used in successive years. Wintering areas are commonly associated with open water though in some areas eagles use habitats with little or no open water if other food resources (e.g., rabbit or deer carrion) are readily available. Avoids areas with nearby human activity (boat traffic, pedestrians) and development (buildings) (Buehler et al. 1991). Bald eagles usually nest in tall trees or on cliffs near water. Nest trees include pines, spruce, firs, cottonwoods, oaks, poplars, and beech. Ground nesting has been reported on the Aleutian Islands in Alaska, in Canada's Northwest Territories, and in Ohio, Michigan, and Texas. Nests located on cliffs and rock pinnacles have been reported historically in California, Kansas, Nevada, New Mexico, and Utah, but currently are known to occur only in Alaska and Arizona. Same nest may be used year after year, or may alternate between two nest sites in successive years. In British Columbia, nests with overhead canopy of foliage were most successful (Palmer 1988). See Livingston et al. (1990) for model of nesting habitat in Maine, Wood et al. (1989) for characteristics of nesting habitat in Florida (most nests in live pine trees). In Oregon, most nests were within 1.6 km of water, usually in largest tree in stand (Anthony and Isaacs 1989). In Colorado and Wyoming, forest stands containing nest trees varied from old-growth ponderosa pine to narrow strips of riparian vegetation surrounded by rangeland (Kralovec et al. 1992).

- 1 **Adult Food Habits:** Carnivore, Piscivore
- 2 **Immature Food Habits:** Carnivore, Piscivore
- 3 **Food Comments:** Feeds opportunistically on fishes, injured waterfowl and seabirds, various
4 mammals, and carrion (Terres 1980). See Haywood and Ohmart (1986), Kralovec et al. (1992),
5 Brown (1993), and Grubb (1995) for diet of inland breeding populations in Arizona, Colorado, and
6 Wyoming. Hunts live prey, scavenges, and pirates food from other birds (e.g., osprey) and, in
7 Alaska, sea otter (Watt et al. 1995, Condor 97:588-590). See Palmer (1988) for further
8 information on hunting methods. In the Columbia River estuary, tidal flats and water less than 4.0
9 meters deep were important foraging habitats (Watson et al. 1991). See Caton et al. (1992) for
10 information on foraging perches used in Montana. Sheep carcasses were significant food
11 sources in winter in Oregon (Marr et al. 1995, Wilson Bulletin 107:251-257).
- 12 **Adult Phenology:** Crepuscular, Diurnal
- 13 **Immature Phenology:** Crepuscular, Diurnal
- 14 **Phenology Comments:** In the Columbia River estuary, foraging activity was most common at
15 low tide and first daylight (Watson et al. 1991). In Arizona, foraging activity during the breeding
16 season peaked at 0800-1000 and 1600-1900 MST (Grubb 1995).
- 17 **Length:** 94 cm
- 18 **Weight:** 5244 grams
- 19 **Management Requirements:** Recovery has been assisted by intensive management that
20 included systematic monitoring, enhanced protection, captive breeding, relocation of wild birds,
21 and publicity (Matthews and Moseley 1990). Knight and Knight (1984) recommended a 450
22 meter buffer between a human in a canoe and a feeding eagle. For northern Chesapeake Bay,
23 Buehler et al. (1991) recommended a 1,360-meter-wide shoreline management zone that
24 extends 1,400 meters inland to encompass nonbreeding roost sites and provide a buffer from
25 human disturbance. Another study recommended a 250-m buffer between a human on land and
26 an eagle in a shoreline tree. A 500-m buffer around the nest may be adequate (see Fraser et al.
27 1985). In Michigan, 75 percent of all alert and flight responses to human activity occurred when
28 activity was within 500 m and 200 m, respectively; vehicles and pedestrians elicited the highest
29 response frequencies. Anthony and Isaacs (1989) made recommendations for Oregon: size of
30 areas for nest-site management should be 50-250 ha, with size and shape depending on
31 surrounding vegetation, topography, and eagle behavior; human activities within 800 m of nests
32 should be restricted from 1 January to 31 August; clearcut logging, road building, hiking trails, and
33 boat launch facilities should not be allowed within 400 m of nests. In Arizona, pedestrians were
34 the most disturbing human activity; eagles were more often flushed from perches than from nests
35 and were most easily disturbed when foraging; eagle response to disturbance frequencies were
36 64% at distances less than 216 m, 45% at 216-583 m, and 24 at distances greater than 583
37 meters (Grubb and King 1991). Along northern Chesapeake Bay, flush distances because of
38 approaching boats averaged 204 meters in winter, 176 meters in summer (Buehler et al. 1991,
39 see for further information on the effects of human activity). In the Columbia River estuary,
40 management of eagle foraging habitats should emphasize protection and enhancement of tidal
41 flats (Watson et al. 1991). See Busch (1988) for a discussion of management activities in the
42 southwestern U.S., Lefranc and Glinski (1988) for management recommendations.
43 Supplemental feeding can be used in efforts to replace diminished supplies of natural foods,
44 provide food free of environmental contaminants, provide essential nutrients, enhance survival of
45 subadults, manipulate distribution of populations, increase nesting success, support released
46 captive-bred birds, and/or afford opportunities for public viewing and education; potential
47 disadvantages of supplemental feeding include prohibitive costs, the loss of natural and cautious
48 behavior, dependence on these food supplies, which may alter migration patterns, and increased

potential for disease transmission (Knight and Anderson 1990). See Grubb (1980) for information on construction and use of an artificial nest structure.

Monitoring Requirements: See Fraser et al. (1983) for information on scheduling reproductive surveys. See Britten et al. (1995) for information on satellite telemetry.

References:

- Abbott, Jackson M. 1978. Chesapeake Bay Bald Eagles. Delaware Conservationist May: P.3-9.
- Abbott, Jackson M. 1982. "Our National Heritage." Delaware Conservationist, 25(1):6-11.
- Abbott, Jackson M. 1982. 1982 Bald Eagle Nest Survey – Chesapeake Bay Region. Alexandria, VA (Printed by author). 6 pp.
- Abbott, Jackson M. 1983. Bald Eagle Continues Comback in Bay. Audubon Naturalist News. October.
- Abbott, Jackson. 1977. Chesapeake Bay Bald Eagle Survey. Hawk Mtn Sanctuary Assn. Thirty-fourth Annual Report, pp 11-14. Alexandria, VA.
- Abbott, Jackson. 1980. Eagle Survey Results Encouraging. Audubon Naturalist News, July-Aug.
- Allen, Craig R., Stephen Demarais, and R. Scott Lutz. 1994. Red Imported Fire Ant Impact on Wildlife: an Overview. Texas J. Sci. 46(1):51-59.
- American Ornithologists' Union (AOU). 1983. Check-list of North American Birds, 6th edition. Allen Press, Inc., Lawrence, KS. 877 p.
- American Ornithologists' Union (AOU). 1983. Check-list of North American Birds, 6th edition. Allen Press, Inc., Lawrence, KS. 877 pp.
- American Ornithologists' Union (AOU). 1983. Checklist of North American birds. 6th ed. American Ornithologists' Union, Allen Press, Inc., Lawrence, KS. 877 pp.
- American Ornithologists' Union (AOU). 1998. Check-list of North American birds. Seventh edition. American Ornithologists' Union, Washington, DC. 829 pp.
- American Ornithologists' Union. 1957. Check-list of North American birds. Fifth edition. American Ornithologists' Union. Port City Press, Inc. Baltimore, MD. 691 pp. (Reprinted in 1961 by Port City Press, Inc., Baltimore, MD.)
- Andrew, J.M. and J.A. Mosher. 1982. Bald eagle nest site selection and nesting habitat in Maryland. J. Wildlife Management 46:382-390.
- Andrews, R. R. and R. R. Righter. 1992. Colorado Birds. Denver Museum of Natural History, Denver. 442 pp.
- Anthony, R. G., and F. B. Isaacs. 1989. Characteristics of bald eagle nest sites in Oregon. J. Wildlife Management. 53:148-159.
- Aquin, P. 1999. Évaluation de la situation des groupes taxonomiques des oiseaux du Québec. Ministère de l'Environnement et de la Faune. 13 pages.
- Audubon Society. 1981-1985. Breeding Bird Atlas of New Hampshire. (unpublished).
- Buckelew, A. and G. Hall. 1994. WV Breeding Bird Atlas. University of Pittsburg Press. BUCKELEW, A. AND G. HALL. 1994.
- Buehler, D.A. et al. 1991. Differences in distribution of breeding, nonbreeding, and migrant bald eagles on the northern Chesapeake Bay. The Condor 93:399-408.

- 1 • Bagley, Fred. 1986-04-15. Unpublished report: Bald Eagle Nest locations in LA.,
2 MS.,AR.,AL.
- 3 • Baicich, P. J., and C. J. O. Harrison. 1997. A guide to the nests, eggs and nestlings of
4 North American birds. Second edition. Academic Press, NY.
- 5 • Baker, R. J., and Y. A. Monstad. 2005. 2005 Minnesota Bald Eagle surveys. Nongame
6 Wildlife Program, Minnesota Department of Natural Resources Report. 4 pp.
- 7 • Baker, R., J. Galli, and S. Nelson. 2000. 2000 Minnesota Bald Eagle survey. Nongame
8 Wildlife Program, Minnesota Department of Natural Resources Report. 2 pp.
- 9 • Beebe, F.L. 1974. Field studies of the falconiformes of British Columbia. Vultures, eagles,
10 hawks, and falcons. Occasional Papers of the British Columbia Provincial Museum No.
11 17. Dept. Recreation and Conservation, Victoria, BC, Canada. 163 pp.
- 12 • Behle, W. H. 1985. Utah birds: geographic distribution and systematics. Utah Mus. Natur.
13 Hist., Occas. Pap. 5: 1-147.
- 14 • Bender, M. (ed.). 1994. Regional News - Region 2. End. Sp. Tech. Bull. 19(5): 18.
- 15 • Bengston, F., and L. D. Frenzel. 1984. Studies of lead toxicity in bald eagles at the Lac
16 Qui Parle Wildlife Refuge. Thesis, Master of Science, University of Minnesota. 95 pp.
- 17 • Bent, A.C. 1937. Life histories of North American birds of prey. Part 1. Bull. U.S. Natl.
18 Mus. 137. 409 pp.
- 19 • Bierly, M.L. 1980. Bird Finding in Tennessee. 3825 Bedford Ave., Nashville, TN 37125.
- 20 • BioSystems Analysis, Inc., Santa Cruz, CA. 1992. Ecology of bald eagles in Arizona.
21 Prepared for Bureau of Reclamation, Lower Colorado Region, Boulder City, NV, Contract
22 No. 6-CS-30-04470.
- 23 • Bird, D. M., editor. 1983. Biology and management of bald eagles and ospreys.
24 MacDonald. 325 pp.
- 25 • BirdLife International. 2004. Threatened birds of the world 2004. CD ROM. BirdLife
26 International, Cambridge, UK.
- 27 • Blood, D.A. and G.G. Anweiler. 1991. Status of the Bald Eagle in British Columbia.
28 Unpublished report to British Columbia Ministry of Environment, Wildlife Branch, Victoria.
29 160 p.
- 30 • Bowerman, W. W., T. G. Grubb, J. P. Giesy, A. J. Bath, and G. A. Dawson. 1993.
31 Population composition and perching habitat of wintering Bald Eagles in northcentral
32 Michigan. Canadian Field Naturalist 107: 273- 278.
- 33 • Britten, M. W., C. L. McIntyre, and M. Kralovec. Satellite radiotelemetry and bird studies
34 in national parks and preserves. Park Science 15(2):20-24.
- 35 • Brown, B. T. 1993. Winter foraging ecology of bald eagles in Arizona. Condor 95:132-
36 138.
- 37 • Brown, B. T., P. L. Warren, and L. S. Anderson. 1988. Status of bald eagles in the Rio
38 Yaqui drainage of Sonora, Mexico. Page 321 in Glinski et al., eds. Proc. Southwest raptor
39 management symposium and workshop. Nat. Wildl. Fed. Sci. and Tech. Ser. No. 11.
- 40 • Brown, B. T., and L. E. Stevens. 1992. Winter abundance, age structure, and distribution
41 of bald eagles along the Colorado River, Arizona. Southwestern Naturalist 37:404-435.
- 42 • Brown, B. T., et al. 1989. Changes in winter distribution of bald eagles along the
43 Colorado River in Grand Canyon, Arizona. J. Raptor Research 23:110-113.
- 44 • Brown, B.T., et.al. 1989. Changes in winter distribution of bald eagles along the Colorado
45 River in Grand Canyon, Arizona. J. Raptor Res. 23(3):110-113.

- 1 • Brownell, V. and M. Oldham. 1984. Status report on the Bald Eagle, *Haliaeetus*
2 *leucocephalus*, in Canada. COSEWIC report.
- 3 • Brownell, V.R. and M.J. Oldham. 1980. Status Report on the Bald Eagle (*Haliaeetus*
4 *leucocephalus*) in Canada. Ontario Ministry of Natural Resources, Toronto, and
5 Committee on the Status of Endangered Wildlife in Canada, Ottawa. 82 pp.
- 6 • Brownell, V.R. and M.J. Oldham. 1980. Status report on the Bale Eagle (*Haliaeetus*
7 *leucocephalus*) in Canada.. Draft Copy. 122 pp.
- 8 • Brownell, V.R. and M.J. Oldham. 1984. Status report on the Bald Eagle, *Haliaeetus*
9 *leucocephalus*, in Canada. Committee on the Status of Endangered Wildlife in Canada
10 (COSEWIC). 88 pp.
- 11 • Buehler, D. A. 2000. Bald Eagle (*Haliaeetus leucocephalus*). Number 506 in A. Poole and
12 F. Gill, editors. The birds of North America. The Birds of North America, Inc.,
13 Philadelphia, PA.
- 14 • Buehler, D. A., S. K. Chandler, T. J. Mersmann, J. D. Fraser, and J. K. D. Seegar. 1992.
15 Nonbreeding Bald Eagle perch habitat on the northern Chesapeake Bay. Wilson Bulletin
16 104:540-545.
- 17 • Buehler, D. A., et al. 1991. Differences in distribution of breeding, nonbreeding, and
18 migrant bald eagles on the northern Chesapeake Bay. Condor 93:399-408.
- 19 • Buehler, D. A., et al. 1991. Effects of human activity on bald eagle distribution on the
20 northern Chesapeake Bay. J. Wildlife Management 55:282-290.
- 21 • Buehler, D. A., et al. 1991. Nonbreeding bald eagle communal and solitary roosting
22 behavior and roost habitat on the northern Chesapeake Bay. J. Wildlife Management
23 55:273-281.
- 24 • Buehler, D. A., et al. 1991. Survival rates and population dynamics of bald eagles on
25 Chesapeake Bay. J. Wildlife Management 55:608-613.
- 26 • Buehler, D. A., et al. 1991. Winter microclimate of bald eagle roosts on the northern
27 Chesapeake Bay. Auk 108:612-618.
- 28 • Buehler, D.A. 2000. Bald Eagle (*Haliaeetus leucocephalus*). In The Birds of North
29 America, No. 506. (A. Poole and F. Gills, eds.). The Birds of North America, Inc.
30 Philadelphia, PA.
- 31 • Burleigh, T. D. 1972. Birds of Idaho. The Caxton Printers, Ltd., Caldwell, ID. 467 pp.
- 32 • Busch, D. E. 1988. Bald eagle. Pages 57-64 in Glinski et al., eds. Proc. Southwest
33 Raptor Manage. Symp. and Workshop. National Wildlife Federation Sci. and Tech. Ser.
34 No. 11.
- 35 • Byrd, M. A., and D. W. Johnston. 1991. Birds. Pages 477-537 in K. Terwilliger,
36 coordinator. Virginia's endangered species: proceedings of a symposium. McDonald and
37 Woodward Publ. Co., Blacksburg, VA.
- 38 • Chesapeake Bay Bald Eagle Recovery Team (Gary J. Taylor, Leader). 1980. The
39 Chesapeake Bay Region Bald Eagle Recovery Plan.
- 40 • Chesapeake Bay Bald Eagle Recovery Team 1982. The Chesapeake Bay Region Bald
41 Eagle Recovery Plan. Region 5, USFWS. 81 PP.
- 42 • Cline, K. 1985. Bald Eagles in the Chesapeake: a Management Guide for Landowners. .
43 U.S. FWS et al. 16 pp.
- 44 • Cline, Keith and William Clark. 1981. Progress Report: Chesapeake Bay Bald Eagle
45 Banding Project. Nat. Wildlife Federation, Raptor Information Center.

- 1 • California Department of Fish and Game (CDF&G). 1990. 1989 annual report on the
2 status of California's state listed threatened and endangered plants and animals. 188 pp.
- 3 • Campbell, R. W., N. K. Dawe, I. McTaggart-Cowan, J. M. Cooper, G. W. Kaiser, and M.
4 C. McNall. 1990. The birds of British Columbia. Volume 2. Nonpasserines: diurnal birds
5 of prey through woodpeckers. University of British Columbia Press, Vancouver, B.C. 636
6 pp.
- 7 • Campbell, R.W., N.K. Dawe, I. McTaggart-Cowan, J.M. Cooper, G.W. Kaiser, and M.C.E.
8 McNall. 1990. The Birds of British Columbia. Vol. 1. Nonpasserines: introduction and
9 loons through waterfowl. Royal British Columbia Museum, Victoria, British Columbia,
10 Canada. 514 pp.
- 11 • Cannings, S. 2001. EO Specifications for Bald Eagle (*Haliaeetus leucocephalus*).
12 NatureServe, Unpublished. 2 pp.
- 13 • Caton, E. L., B. R. McClelland, D. A. Patterson, and R. E. Yates. 1992. Characteristics of
14 foraging perches used by breeding Bald Eagles in Montana. Wilson Bulletin 104(1):136-
15 142.
- 16 • Caton, E. L., et al. 1992. Characteristics of foraging perches used by breeding bald
17 eagles in Montana. Wilson Bull. 104:136-142.
- 18 • Chester, D. N., et al. 1990. Habitat use by nonbreeding bald eagles in North Carolina. J.
19 Wildlife Management 54:223-234.
- 20 • Chippewa National Forest. 1979. Bald Eagle - Osprey Status Report, 1979. Chippewa
21 National Forest, Cass Lake, MN. 5 pp.
- 22 • Colorado Bird Observatory. 1997. 1996 Reference Guide to the Monitoring and
23 Conservation Status of Colorado's Breeding Birds. Colorado Bird Observatory, Colorado
24 Division of Wildlife, Great Outdoors Colorado Trust Fund, and Partners, March 21, 1997.
- 25 • Cory, C.B. 1909. The birds of Illinois and Wisconsin. Field Mus. Nat. Hist. Publ. 131,
26 Zool. Ser. 9:1-766.
- 27 • Craig, Gerald R. 1997. Recommended buffer zones and seasonal restrictions for
28 Colorado raptor nests. Unpublished report for Colorado Division of Wildlife. Updated
29 January 6, 1997.
- 30 • Craig, R.J. et al. 1988. Time and energy budgets of bald eagles wintering along the
31 Connecticut River. Jo. Field Ornith. 59:22-32.
- 32 • Curnutt, J. L. 1992. Dynamics of a year-round communal roost of bald eagles. Wilson
33 Bull. 104:536-540.
- 34 • Dickinson, Mary B., ed. 1999. Field Guide to the Birds of North America, 3rd ed. National
35 Geographic Society, Washington, D.C. 480 pp.
- 36 • DeGraaf, R. M. and D. R. Rudis. 1986. New England wildlife: habitat, natural history, and
37 distribution. Univ. Mass. Press. Amherst, MA. 491 pp.
- 38 • Deschamps, V. 1997a. Southwestern Ontario Bald Eagle Monitoring Project - 1996 Final
39 Report. Unpub. report, Ontario Birds At Risk Program, Long Point Bird Observatory, St.
40 Williams, Ontario. 8 pp.
- 41 • Deschamps, V. 1997b. Southwestern Ontario Bald Eagle Monitoring Project - 1997
42 Report. Unpub. report, prepared for OMNR, Southcentral Region, Aylmer District, Ontario
43 Birds At Risk Program, Long Point Bird Observatory, St. Williams, Ontario. 15 pp.
- 44 • Desrosiers A., F. Caron et R. Ouellet. 1995. Liste de la faune vertébrée du Québec. Les
45 publications du Québec.

- 1 • Detrich, Philip J. 1988. Evaluation of the Relationship Between the Morphoedaphic Index
2 and Bald Eagle Nesting Density in Minnesota. Funded by the Minnesota Department of
3 Natural Resources. Results in unpublished report.
- 4 • Dionne C. 1906. Les oiseaux de la province de Québec. Dussault et Proulx.
- 5 • Division of Natural Resources, Navajo Fish and Wildlife Department. 1995. Endangered
6 Species List for The Navajo Nation.
- 7 • Dugoni, Joseph A. 1980. Habitat utilization, food habits, and productivity or nesting
8 southern bald eagles in La. MS thesis, Louisiana State University - Baton Rouge.
- 9 • Dunn, E. H., C. M. Downes, and B. T. Collins. 2000. The Canadian Breeding Bird Survey,
10 1967-1998. Canadian Wildlife Service Progress Notes No. 216. 40 pp.
- 11 • Dzus, E., and J. Gerrard. 1993. Factors influencing bald eagle densities in northcentral
12 Saskatchewan. J. Wildlife Management 57:771-778.
- 13 • Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1992. Birds in Jeopardy: the Imperiled and
14 Extinct Birds of the United States and Canada, Including Hawaii and Puerto Rico.
15 Stanford University Press, Stanford, California. 259 pp.
- 16 • Evans, D. L. 1982. Status reports on twelve raptors. U.S. Department of the Interior, Fish
17 and Wildlife Service, Special Scientific Report No. 238. 68 pp.
- 18 • Evers, D. C. 1992. A guide to Michigan's endangered wildlife. Univ. Michigan Press, Ann
19 Arbor. viii + 103 pp.
- 20 • Frenzel, R.W. and R.G. Anthony. 1989. Relationship of diets and environmental
21 contaminants in wintering bald eagles. J. Wildl. Mgmt. 53(3):792-802.
- 22 • Field, M. 1986. 1995 Bald Eagle Nesting Report for Southwestern Ontario. Ontario Birds
23 At Risk Newsletter Vol. 3 (1): 3-4.
- 24 • Field, M. 2000. 2000 Elgin County Bald Eagle Nesting. The Cardinal, No. 180: 38-39.
- 25 • Field, M. 2001. Bald Eagle Nests in Elgin County, Ontario, 2001. The Cardinal. No. 185:
26 32-33.
- 27 • Finnley, D. (ed.). 1979. Endangered species: new challenge for the Navajo. Endangered
28 Species Technical Bulletin 4(6):1,7-11.
- 29 • Fischer, D.L. 1985. Piracy behavior of wintering bald eagles. The Condor 87:246-251.
- 30 • Fisher, A.K. 1893. The hawks and owls of the United States in their relation to agriculture.
31 Washington U.S. Dept. of Agriculture Bull. no. 6. 210 pp.
- 32 • Forbis, L. A. 1988. Status and trends of bald eagles breeding in Arizona, 1975-1986.
33 Pages 282-288 in Glinski et al., eds. Proc. Southwest raptor management symposium
34 and workshop. Nat. Wildl. Fed. Sci. and Tech. Ser. No. 11.
- 35 • Fraser, J. D., L. D. Frenzel, and J. E. Mathisen. 1985. The impact of human activities on
36 breeding bald eagles in north-central Minnesota. J. Wildlife Management 49:585-592.
- 37 • Fraser, J. D., et al. 1983. Scheduling bald eagle reproductive surveys. Wildlife Society
38 Bull. 11:13-16.
- 39 • Frazer, J.D. 1985. The impact of human activities on Bald Eagle populations - a review.
40 Pp. 68-84 in J.M. Gerrard and T.N. Ingram (eds.), The Bald Eagle in Canada. White
41 Horse Plains Publishers, Headingley, Manitoba.
- 42 • Frenzel, L.D. 1988-1989. MN DNR Bald Eagle Nesting Activity and Productivity Data by
43 County. Funded by the MN DNR, Section of Wildlife, Nongame Research Program.
44 Results in unpublished report.

- 1 • Frenzel, L.D. 1988. 1988 MN DNR bald eagle nesting activity and productivity data by
2 county. Draft final report to the Minnesota Department of Natural Resources. 7 pp.
- 3 • Frenzel, L.D. 1989. 1989 MN DNR bald eagle nesting activity and productivity data by
4 county. Draft final report to the Minnesota Department of Natural Resources. 6 pp.
- 5 • Garrett, M. G., J. W. Watson, and R. G. Anthony. 1993. Bald Eagle home range and
6 habitat use in the Columbia River estuary. Journal of Wildlife Management 57:19-27.
- 7 • Garrett, M. G., J. W. Watson, and R. G. Anthony. 1993. Bald Eagle home range and
8 habitat use in the Columbia River estuary. Journal of Wildlife Management 57:19-27.
- 9 • Gerrard, J. M. 1983. A review of the current status of bald eagles in North America. Pp.
10 5-21 in D. M. Bird (Ed.). Biology and management of bald eagles and ospreys. Harpell
11 Pr., Ste. Anne de Bellevue, Que.
- 12 • Gerrard, J. M., P. N. Gerrard, G. R. Bortolotti, and E. H. Dzus. 1992. A 24-year study of
13 Bald Eagles on Besnard Lake, Saskatchewan. Journal of Raptor Research 26:159-166.
- 14 • Gerrard, J. M., P. N. Gerrard, G. R. Bortolotti, and E. H. Dzus. 1992. A 24-year study of
15 Bald Eagles on Besnard Lake, Saskatchewan. Journal of Raptor Research 26:159-166.
- 16 • Gerrard, J. M., and G. R. Bortolotti. 1988. The bald eagle. Haunts and habits of a
17 wilderness monarch. Smithsonian Institution Press, Washington, D.C. 194 pp.
- 18 • Gerrard, J.M. 1983. A review of the current status of Bald Eagles in North America.
19 Pages 5-21 in D. M. Bird (Ed.). Biology and Management of Bald Eagles and Ospreys.
20 Proceedings of First International Symposium on Bald Eagles and Ospreys, Montreal, 28-
21 29 October 1981. Harpell Press. Ste. Anne de Bellevue, Quebec.
- 22 • Gibbs, J. P., and S. M. Melvin. 1992. American bittern, *Botaurus lentiginosus*. Pages 51-
23 69 in K. J. Schneider and D. M. Pence, editors. Migratory nongame birds of management
24 concern in the Northeast. U.S. Fish and Wildlife Service, Newton Corner, MA. 400 pp.
- 25 • Godfrey, Earl W. 1986. Birds of Canada. National Museums of Canada, Ottawa, Ontario,
26 Canada. 595 p. + plates.
- 27 • Green, N. 1985. The Bald Eagle. Pp 508-531 in R.L. DiSilvestro, ed., Audubon Wildlife
28 Report 1985. National Audubon Society, NY.
- 29 • Grier, J.W., T. Armstrong, P. Hunter, S. Lockhart and B. Ranta. 2001. Report on the
30 Status of Bald Eagles in Ontario. Unpublished draft (12 July). 47 pp.
- 31 • Griffin, C. R., T. S. Baskett, and R. D. Sparrowe. 1982. Ecology of bald eagles wintering
32 near a waterfowl concentration. U.S. Fish and Wildlife Service Special Science Report -
33 Wildlife No. 247:1-12.
- 34 • Grub, T.G., et.al. 1983. 1983 Arizona bald eagle research report. Rocky Mountain Forest
35 and Range Experiment Station, U.S. Forest Service.
- 36 • Grubb, T. G. 1980. An artificial bald eagle nest structure. USDA Forest Service, Rocky
37 Mountain Forest and Range Experimental Station, Research Note RM-383.
- 38 • Grubb, T. G. 1995. Food habits of bald eagles breeding in the Arizona desert. Wilson
39 Bulletin 107:258-274.
- 40 • Grubb, T. G., S. J. Nagiller, W. L. Eakle, and G. A. Goodwin. 1989. Winter roosting
41 patterns of bald eagles (*Haliaeetus leucocephalus*) in north-central Arizona.
42 Southwestern Naturalist 34:453-459.
- 43 • Grubb, T. G., W. W. Bowerman, J. P. Giesy, and G. A. Dawson. 1992. Responses of
44 breeding bald eagles, *Haliaeetus leucocephalis* [sic], to human activities in northcentral
45 Michigan. Canadian Field-Naturalist 106:443-453.

- 1 • Grubb, T. G., W. W. Bowerman, and P. H. Howey. 1994. Tracking local and seasonal
2 movements of wintering Bald Eagles *Haliaeetus leucocephalus* from Arizona and
3 Michigan with satellite telemetry. Pages 247-358 In B.U. Meyburg and R. D. Chancellor,
4 editors. Raptor conservation today. Pica Press, Moose Jaw, Saskatchewan.
- 5 • Grubb, T. G., W. W. Bowerman, and P. H. Howey. 1994. Tracking local and seasonal
6 movements of wintering Bald Eagles *Haliaeetus leucocephalus* from Arizona and
7 Michigan with satellite telemetry. Pages 247-358 IN B.-U. Meyburg and R. D. Chancellor,
8 editors. Raptor conservation today. Pica Press, Moose Jaw, Saskatchewan.
- 9 • Grubb, T. G., and R. M. King. 1991. Assessing human disturbance of breeding bald
10 eagles with classification tree models. J. Wildlife Management 55:500-511.
- 11 • Grubb, T.G. 1984. 1984 Arizona bald eagle research report. Annual progress report
12 prepared by U.S. Forest Service, Rocky Mountain Forest and Range Experiment Station,
13 Tempe, AZ, for Master Interagency Agreement No. IAG-43.
- 14 • Grubb, T.G. and C.E. Kennedy. 1982. RUN WILD, Wildlife/Habitat Relationships: 1978
15 bald eagle winter habitat on the National Forest System in the Southwest. Southwestern
16 Region, U.S. Forest Service Wildlife Unit Technical Series. 116 p.
- 17 • Hall, G. 1983. Birds of WV. Carnegie Museum of Natural History.
- 18 • Hancock, J. and J. Kushlan. 1984. The Herons Handbook.
- 19 • Harper, R.G. et al. 1988. Nonfish prey of wintering bald eagles in Illinois. Wilson Bull.
20 100:688-690.
- 21 • Harrison, C. 1978. A Field Guide to the Nests, Eggs and Nestlings of North American
22 Birds. Collins, Cleveland, Ohio.
- 23 • Hawk Watch International. 2003. An online search for migration watch results, 1999-2002
24 data. . Accessed 11 September 2003.
- 25 • Hayward, C. L., C. Cottam, A. M. Woodbury, and H. H. Frost. 1976. Birds of Utah. Great
26 Basin Naturalist Memoirs 1: 229 pp.
- 27 • Haywood, D. D., and R. D. Ohmart. 1986. Utilization of benthic-feeding fish by inland
28 breeding bald eagles. Condor 88:35-42.
- 29 • Heagy, A. 1996. Southwestern Ontario Bald Eagle Monitoring Program. Ontario Birds At
30 Risk Newsletter. Vol. 3 (1): 2.
- 31 • Hensel, R. J., and W. A. Troyer. 1964. Nesting studies of the bald eagle in Alaska.
32 Condor 66:282-286.
- 33 • Herkert, J. R., editor. 1992. Endangered and threatened species of Illinois: status and
34 distribution. Vol. 2: Animals. Illinois Endangered Species Protection Board. iv + 142 pp.
- 35 • Herkert, Jim. 1998. Proposed additions, deletions, and changes to the Illinois List of
36 Threatened and Endangered Animals. 101st ESPB Meeting, August 21, 1998. 16pp.
- 37 • Hoffman, S.W., et.al. 1992. Patterns and recent trends in counts of migrant hawks in
38 western North America: 1977-1991. Draft. Prepared by HawkWatch International,
39 Albuquerque, NM, for U.S. Fish and Wildlife Service, Office of Migratory Bird
40 Management, Washington, DC.
- 41 • Howell, S. N. G., and S. Webb. 1995. A guide to the birds of Mexico and northern Central
42 America. Oxford University Press, Oxford, UK.
- 43 • Hunter, P., and D. Baird. 1994/95. The bald eagle in Ontario's Great Lakes basin. Bird
44 Trends (Canadian Wildlife Service) (4):17-18.

- 1 • Hessel, D.J.T. and L. Brown. 1992. Population Changes in Diurnally-Migrating Raptors
2 and Duluth, Minnesota (1974-1989) and Grimsby, Ontario (1975-1990). Ontario Ministry
3 of Natural Resources.
- 4 • Imhof, T.A. 1976. Alabama Birds.
- 5 • Johnsgard, P. A. 1990. Hawks, eagles, and falcons of North America. Smithsonian Inst.
6 Press, Washington, D.C. xvi + 403 pp.
- 7 • Knight, R.L. and S.K. Skagen. 1988. Agnostic asymmetries and foraging ecology of bald
8 eagles. *Ecol.* 69(4):1188-1194.
- 9 • Kyle, Rober. 1982. Our National Bird Soars Again. *The Washington Post*, June 18, p. 57.
- 10 • King, W. B., compiler. 1979. Endangered birds of the world. The International Council for
11 Bird Preservation. Smithsonian Institution Press, Washington, D.C. [Reprinted in
12 handbook form in 1981.]
- 13 • Kirk, D. A., D. Hessel, and E. Dunn. 1995. Raptor population status and trends in
14 Canada. *Bird Trends (Canadian Wildlife Service)* 4:2-9.
- 15 • Kirk, D.A., and C. Hyslop. 1998. Population status and recent trends in Canadian raptors:
16 a review. *Biological Conservation* 83 (1): 91-118.
- 17 • Kjos, D.G. 1992. Bald eagle numbers continue to rise. *Endangered Species Technical*
18 *Bulletin* Vol 27, nos. 1-2. 4p.
- 19 • Knight, R. L., and D. P. Anderson. 1990. Effects of supplemental feeding on an avian
20 scavenging guild. *Wildl. Soc. Bull.* 18:388-394.
- 21 • Knight, R. L., and S. K. Knight. 1984. Responses of wintering bald eagles to boating
22 activity. *J. Wildlife Management* 48:999-1004.
- 23 • Kozie, K. D., and R. K. Anderson. 1991. Productivity, diet, and environmental
24 contaminants in bald eagles nesting near the Wisconsin shoreline of Lake Superior. *Arch.*
25 *Environ. Contam. Toxicol.* 20:41-48.
- 26 • Kralovec, M. L., R. L. Knight, G. R. Craig, and R. G. McLean. 1992. Nesting productivity,
27 food habits, and nest sites of bald eagles in Colorado and southeastern Wyoming.
28 *Southwestern Naturalist* 37:356-361.
- 29 • LaRue, C.T. 1994. Birds of northern Black Mesa, Navajo County, Arizona. *Great Basin*
30 *Naturalist* 54(1):1-63.
- 31 • Lagacé M., L. Blais et D. Banville. 1983. Liste de la faune vertébrée du Québec.
32 Première édition. Ministère du Loisir, de la Chasse et de la Pêche.
- 33 • Larrison, E. J., J. L. Tucker, and M. T. Jollie. 1967. Guide to Idaho birds. *Journal of the*
34 *Idaho Academy of Science* 5: 1-220.
- 35 • Leberman, R.C. 1987. A Field List of the Birds of Western Pennsylvania and Adjacent
36 Regions. Unpublished.
- 37 • Lefranc, M. N., Jr., and R. L. Glinski. 1988. Southwest raptor management issues and
38 recommendations. Pages 375-392 in Glinski et al., eds. *Proc. Southwest raptor*
39 *management symposium and workshop. National Wildlife Federation Science and Tech.*
40 *Ser. No. 11.*
- 41 • Lehman, R.N. 1979. A survey of selected habitat features of 95 bald eagle nest sites in
42 California. California Department of Fish and Game, Wildlife Management Branch,
43 Administrative Report 79-1.

- 1 • Lincer, J. L., W. S. Clark, and M. N. LeFranc, Jr. 1979. Working bibliography of the bald
2 eagle. Raptor Information Center, National Wildlife Federation, Washington, D.C. NWF
3 Scientific/Technical Series No. 2. 219 pp.
- 4 • Lincer, J.L., et.al. 1979. Working bibliography of the bald eagle. National Wildlife
5 Federation Scientific/Technical Series No. 2, Washington, DC.
- 6 • Lincer, J.L.; W.S. Clark and M.N. LeFranc, JR. 1979. Working Bibliography of the Bald
7 Eagle. Raptor Information Center, Nat. Wildlife Fed., Washington, D.C. USNWF
8 Scientific/Technician – CAL Series No. 2 219 pp.
- 9 • Livingston, S. A., et al. 1990. Habitat models for nesting bald eagles in Maine. J. Wildlife
10 Management 54:644-653.
- 11 • Lowery, George H. 1974. The Birds of Louisiana. LSU Press. 651pp.
- 12 • McKnight, J. 1989. Memorandum of November 3, to J. McKegg et al. RE: Somerville and
13 Associated Properties, Alliance Property Corporation, Calvert County, MD.
- 14 • Moser, J.A. and J.M. Andrew. 1981. Nesting Habitat and Nest Site Selection by the Bald
15 Eagle in Maryland. UMD, AEL, Frostburg State College, Frostburg, MD.
- 16 • Mahaffy, M. S., and L. D. Frenzel. 1987. Elicited territorial responses of northern bald
17 eagles near active nests. Journal of Wildlife Management 51:551-554.
- 18 • Maniscalco, J., compiler. 1992. *Haliaeetus leucocephalus* (Linnaeus) bald eagle: a
19 working bibliography. Available as a printed document or on diskette.
- 20 • Martell, M. 1988-1989. A study on wintering and released rehabilitated bald eagles in
21 Minnesota. A report submitted to the Minnesota Department of Natural Resources
22 Nongame Program. 34 pp.
- 23 • Martell, M. S., J. B. Nibe, P. T. Redig, and G. M. Buhl. 1991. A study of bald eagles
24 wintering along the Mississippi river between St. Paul and Red Wing, Min., and the St.
25 Croix river south of Hudson, Wis., 1987-1990. 1991. and Wisconsin rivers. Final report
26 submitted to the Minnesota Department of Natural Resources Nongame Program. 53 pp.
- 27 • Martell, M., C. Gieck, J. Nibe, D. Erickson, B. Mandernack, and P. T. Redig. 1991. Bald
28 eagle winter roosts on the Mississippi and Wisconsin rivers. A report submitted to the
29 Minnesota and Wisconsin Departments of Natural Resources. 42 pp.
- 30 • Martell, M., J. Nibe, and J. Galli. 1990. A report on bald eagle use of the Wacouta Bay,
31 (Mn.) area of the Mississippi River along with management recommendations. Final
32 report to the Minnesota Department of Natural Resources Nongame Wildlife Program. 15
33 pp.
- 34 • Martell, M., P. T. Redig., J. Nibe, and G. Buhl. 1990. Survival of released rehabilitation
35 bald eagles. Final report submitted to the Minnesota Department of Natural Resources.
36 73 pp.
- 37 • Martell, Mark; J. Nibe; P. Redig; and G. Buhl. 1987-1991. A Study on Wintering and
38 Released Rehabilitated Bald Eagles in Minnesota. Funded by the MN DNR, Section of
39 Wildlife, Nongame Research Program and Hormel, Incorporated. Results in unpublished
40 report.
- 41 • Mathisen, J. E. 1987. Bald eagle and osprey populations. Wildlife monitoring report,
42 Chippewa National Forest. 4+ pp.
- 43 • Matthews, J.R. and C.J. Moseley (eds.). 1990. The Official World Wildlife Fund Guide to
44 Endangered Species of North America. Volume 1. Plants, Mammals. xxiii + pp 1-560 +
45 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians,
46 Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180.
47 Beacham Publications, Inc., Washington, D.C.

- 1 • McAtee W.L. 1959. Folk - names of Canadian birds. National Museum of Canada. Folk -
2 names of Canadian birds. National Museum of Canada. 74 pages.
- 3 • Millsap, B. A. 1986. Status of wintering bald eagles in the coterminous 48 states. Wildlife
4 Society Bull. 14:433-440.
- 5 • Montana Bald Eagle Working Group. 1991. Habitat management guide for bald eagles in
6 northwestern Montana. 29pp.
- 7 • Montana Bald Eagle Working Group. 1994. Montana Bald Eagle management plan. 2nd
8 edition. Bureau of Reclamation. 104 pp.
- 9 • Montana Bird Distribution Online Database. 2001. Helena, Montana, USA. April-
10 September 2003. <http://nhp.nris.state.mt.us/mbd/>.
- 11 • Montopoli, G. J., and D. A. Anderson. 1991. A logistic model for the cumulative effects of
12 human intervention on bald eagle habitat. J. Wildlife Management 55:290-293.
- 13 • Newton, I. 1976. Population Limitation in Diurnal Raptors. Canadian field-Naturalist
14 90(3):274-300.
- 15 • Norman, D.M., A.M. Breault and J.E. Moul. 1989. Bald Eagle Incursions and Predations
16 at Great Blue Heron Colonies. Colonial Waterbirds. 12(2):215-216.
- 17 • National Geographic Society (NGS). 1983. Field guide to the birds of North America.
18 National Geographic Society, Washington, DC.
- 19 • National Geographic Society, 1987. Field Guide to the birds of North America, second
20 edition. The National Geographic Society, Washington, D.C.
- 21 • Natural Heritage and Nongame Research Program. 1996. Minnesota's list of
22 Endangered, Threatened, and Special Concern species. Effective 7/1/96. 16 pp.
- 23 • Nelson, D. 1993. Colorado Bird Atlas: Manual on Use of Breeding Codes. Denver
24 Museum of Natural History, Denver. 27 pp.
- 25 • Nicholson, C.P. 1997. Atlas of the breeding birds of Tennessee. The University of
26 Tennessee Press. 426 pp.
- 27 • Ohmart, R.D. and R.J. Sell. 1980. The bald eagle of the Southwest with special emphasis
28 on the breeding population of Arizona. U.S. Prepared by Arizona State University for U.S.
29 Water and Power Resources Service Contract No. BR-14-06-330-2674. 95 pp.
- 30 • Ouellet H., M. Gosselin et J.P. Artigau. 1990. Nomenclature française des oiseaux
31 d'Amérique du Nord. Secrétariat d'État du Canada. 457 p.
- 32 • Palmer, R. S., editor. 1988. Handbook of North American birds. Vol. 4. [Diurnal raptors,
33 part 1]. Yale University Press, New Haven, CT. vii + 433 pp.
- 34 • Parker III, T. A., D. F. Stotz, and J. W. Fitzpatrick. 1996. Ecological and distributional
35 databases for neotropical birds. The University of Chicago Press, Chicago.
- 36 • Parks Canada. 2000. Vertebrate Species Database. Ecosystems Branch, 25 Eddy St.,
37 Hull, PQ, K1A 0M5.
- 38 • Pendleton, B. A. G., B. A. Millsap, K. W. Cline, and D. M. Bird. 1987. Raptor
39 management techniques manual. National Wildlife Federation, Sci. and Tech. Ser. No.
40 10. 420 pp.
- 41 • Pennsylvania Breeding Bird Atlas Project. 1984-1987. Unpublished.
- 42 • Peterson, R. T. 1947. A Field Guide to the Birds. Houghton Mifflin Company, Boston. 230
43 pp.

- 1 • Peterson, R.T. 1980. A field guide to the birds of eastern and central North America.
2 Houghton Mifflin Company, Boston.
- 3 • Peterson, R.T. 1990. A field guide to western birds. Houghton Mifflin Company, Boston.
- 4 • Postupalsky, Sergej. 1968. The Status of the Bald Eagle at Lake Nipigon and the
5 Thunder Bay and Black Bay Area of Lake Superior, Ontario. report of an original field
6 study conducted for Investigations in Ornithology (Zoology 709), The University of
7 Michigan Biological Station. Not for Publication. 17 pp.
- 8 • Raffaele, H., J. Wiley, O. Garrido, A. Keith, and J. Raffaele. 1998. A guide to the birds of
9 the West Indies. Princeton University Press, Princeton, NJ. 511 pp.
- 10 • Redig, P. T., G. E., S. Schwartz, and E. Lawler. 1983. An investigation into the effects of
11 lead poisoning on bald eagles and other raptors: final report Submitted to the Minnesota
12 Department of Natural Resources. Unpagged.
- 13 • Redig, Patrick T., Gary E., Samuel Schwartz, and Ellen Lawler. 1980 - 1983. An
14 Investigation into the Effects of Lead Poisoning on Bald Eagles and Other Raptors: Final
15 Report. Funded by the MN DNR, Section of Wildlife, Nongame Research Program.
16 Results in unpublished report.
- 17 • Ridgway, R. 1889. The ornithology of Illinois. Vol. 1. Ill. State Lab. Nat. Hist. 520pp.
- 18 • Root, T. 1988. Atlas of wintering North American birds: An analysis of Christmas Bird
19 Count data. University of Chicago Press. 336 pp.
- 20 • Ryke, N., D. Winters, L. McMartin and S. Vest. 1994. Threatened, Endangered and
21 Sensitive Species of the Pike and San Isabel National Forests and Comanche and
22 Cimarron National Grasslands. May 25, 1994.
- 23 • Shea, D. S. 1973. A management-oriented study of bald eagle concentrations in Glacier
24 National Park. M.S. thesis. University of Montana, Missoula, 78 pp.
- 25 • Sibley, C.G., and B.L. Monroe, Jr. 1990. Distribution and Taxonomy of Birds of the World.
26 Yale University Press, New Haven, CT. xxiv + 1111 pp.
- 27 • Sibley, Charles G. and Burt L. Monroe, Jr. 1990. Distribution and Taxonomy of Birds of
28 the World. Yale University Press. New Haven, Connecticut and London. 1111pp.
- 29 • Snow, C. 1973. Southern bald eagle (*Haliaeetus leucocephalus leucocephalus*) and
30 Northern bald eagle (*Haliaeetus leucocephalus alascanus*). U.S Bureau of Land
31 Management Habitat Management Series for Endangered Species Report No. 5,
32 Portland, OR.
- 33 • Sogge, M.K., et.al. 1995. Monitoring winter bald eagle concentrations in the Grand
34 Canyon: 1993-1995. National Biological Service Colorado Plateau Research
35 Station/Northern Arizona University. 33 p.
- 36 • Spencer, C. N., B. R. McClelland, and J. A. Stanford. 1991. Shrimp stocking, salmon
37 collapse, and eagle displacement. BioScience 41:14-21.
- 38 • Stalmaster, M. V. 1987. The bald eagle. Universe. 227 pp.
- 39 • Stalmaster, M.V. 1987. The Bald Eagle.
- 40 • Steenhof, K. 1978. Management of Wintering Bald Eagles. U.S. Fish and Wildlife
41 Service, FWS/OBS-79/79, 55 pp.
- 42 • Steenhof, K. 1978. Management of wintering bald eagles. U.S. Fish and Wildlife Service,
43 FWS/OBS-79/79, 55 pp.
- 44 • Stupka, A. 1963. Notes on the Birds of Great Smoky Mountains National Park.

- 1 • Taylor, G.J. 1986. Memo to D.D. Boone et al. RE: 1986 Bald Eagle Nesting Productivity
2 in Maryland.
- 3 • Taylor, Gary J. AND Glenn D. Therres. 1981. A Computer Generated Description of
4 Potential Bald Eagle Nesting Habitat in Maryland. Wildlife Administration, DNR,
5 Annapolis, MD.
- 6 • Terres, J. K. 1980. The Audubon Society Encyclopedia of North American Birds. Alfred A.
7 Knopf, NY.
- 8 • Titus, K., and M. R. Fuller. 1990. Recent trends in counts of migrant hawks from
9 northeastern North America. Journal of Wildlife Management 54:463-470.
- 10 • Todd, R. 1978. Winter bald eagle in Arizona. Arizona Game and Fish Department,
11 Federal Aid in Wildlife Restoration Project W-53-R Work Plan 5 Job 1, Special Report.
- 12 • Todd, R. 1981. Multi-agency findings on the distribution of bald eagles for Arizona in the
13 January months of 1979-1981. Arizona Game and Fish Department, Federal Aid Project
14 W-53-R-31 Work Plan 5 Job 1, Special Report.
- 15 • U.S. Fish & Wildlife Service (USFWS). 1984. Draft Pacific states bald eagle recovery
16 plan.
- 17 • 1990. Endangered and threatened species recovery program: report to Congress. 406
18 pp.
- 19 • U.S. Fish and Wildlife Service. 1982. Bald eagle recovery plan (Southwestern
20 population). U.S. Fish and Wildlife Service, Albuquerque, NM. 65 p.
- 21 • U.S. Fish and Wildlife Service. 1999. Proposed rule to remove the Bald Eagle in the
22 lower 48 states from the list of endangered and threatened wildlife. Federal Register
23 64:36453-36464.
- 24 • Vermeer, K., and K. H. Morgan. 1989. Nesting population, nest sites, and prey remains of
25 bald eagles in Barkeley Sound, British Columbia. Northwestern Naturalist 70:21-26.
- 26 • Watson, J. W., M. G. Garrett, and R. G. Anthony. 1991. Foraging ecology of bald eagles
27 in the Columbia River estuary. J. Wildlife Management 55:492-499.
- 28 • Weekes, F.M. 1975. Behavior of a young Bald Eagle at a southern Ontario nest.
29 Canadian Field-Naturalist 89: 35-40.
- 30 • Wheye, D. 1992. Birds in Jeopardy: The Imperiled and Extinct Birds of the United States
31 and Canada, Including Hawaii and Puerto Rico. Stanford University Press. Stanford, CA.
32 259 pp.
- 33 • White, L. 1995. Bald eagle distribution and habitat use on the Animas, La Plata, and
34 Mancos rivers, Colorado and New Mexico, results of winter surveys: 1993–1995.
35 Technical Service Center, U.S. Bureau of Reclamation, Denver.
- 36 • Whittam, B. 2000. Southern Ontario Bald Eagle Monitoring Project - 2000 Report. Report
37 prepared by Bird Studies Canada, Ontario Birds At Risk Program for Ontario Power
38 Generation, Ontario Ministry of Natural Resources, Canadian Wildlife Service, and the
39 Edwards Charitable Foundation. 11 pp.
- 40 • Wiemeyer, S. N. et al. 1989. Environmental contaminants in blood of western bald
41 eagles. J. Raptor Res. 23:140-146.
- 42 • Witmer, G., and T. A. O'Neil. 1990. Assessing cumulative impacts to wintering bald
43 eagles in western Washington. Pages 144-150 in Mitchell et al., eds. Ecosystem
44 management: rare species and significant habitats. New York State Museum Bull. 471.
- 45 • Wood, Merrill. 1979. Birds of Pennsylvania. Pennsylvania State University, University
46 Park.133 pp.

- 1 • Wood, P. B., T. C. Edwards, Jr., and M. W. Collopy. 1989. Characteristics of bald eagle
2 nesting habitat in Florida. J. Wildlife Management 53:441-449.
- 3 • Woodliffe, P.A. 1992. Recent nesting history of the Bald Eagle in Rondeau Provincial
4 Park, Ontario. Ontario Birds 10(3):101-107.
- 5 • Yeager, B. and B. Masslich. 1995. Biological assessment for the St. George road
6 Projects. Prepared for Utah Department of Transportation. Bio/West, Inc., 1063 West
7 1400 North, Logan, UT 84321.
- 8

California Brown Pelican (*Pelecanus occidentalis californicus*)

The California brown pelican has an implied federally Endangered status because it is a subspecies of the federally endangered brown pelican (*Pelecanus occidentalis*), which was listed on November 15, 1994.

Historic Range: Breeds along Pacific coast of central and southern California (the Channel Islands south), on islands off Baja California and on islands in the Gulf of California (south to Isabella and the Tres Marias Islands); ranges regularly north of the breeding grounds to southern British Columbia (Johnsgard 1993, AOU 1998). [*Pelecanus occidentalis*: BREEDING: along Pacific coast from southern California to Peru and (where *thagus* is regarded as conspecific) central Chile, and along Atlantic, Gulf, and Caribbean coasts from Maryland south around Florida to southern Texas, Bahamas (Sprunt 1984), West Indies, off Yucatan Peninsula, and off Venezuela and Caribbean coast of Colombia. Ranges in Pacific coastal waters north to southern British Columbia (after breeding, before winter). NON-BREEDING: Ranges in Pacific coastal waters north to southern British Columbia (after breeding, before winter). In western North America, winters mainly from California south. In the southeastern U.S., primary winter range includes Florida and the Gulf Coast. Subspecies *carolinensis*: breeds locally in Maryland and Virginia and south to Florida (primary nesting range), also locally in Louisiana (where reintroduced) and in central coastal Texas; breeds locally also off northeastern Yucatan and Belize, and ranges southward through coastal Honduras and Costa Rica to Panama, where local breeding occurs off the Pacific coast; vagrants wander north to New England and occur casually inland to the Great Lakes and Great Plains states (Johnsgard 1993). Breeds also in the Bahamas (Sprunt 1984) (extirpated, according to Johnsgard 1993). Ranges throughout breeding range and along eastern shores of Mexico south along Central America to the Caribbean coasts of Colombia and Venezuela, and through the Greater and Lesser Antilles to Trinidad; and on the Pacific coast of Central America (AOU 1957). Subspecies *californicus*: breeds along Pacific coast in southern California (Anacapa Island), and in Mexico on islands off Baja California and on islands in the Gulf of California (south to Isabella and the Tres Marias Islands); possibly locally along the coast of Sonora and Sinaloa; vagrants have occurred north to British Columbia and Idaho (Johnsgard 1993).]

Basic Description: A large bird (brown pelican).

General Description: A large heavy water bird with a massive bill and huge throat pouch; wings and body are mostly grayish-brown; nonbreeding adult has a whitish head and neck, often washed with yellow; hindneck of breeding adult is dark chestnut; head and neck of juvenile is grayish brown; size varies greatly depending on location, with the smallest individuals in the West Indies, medium birds on the coasts of the U.S. (Atlantic and Gulf), Central America, and Colombia and Ecuador, large birds on the coasts of California, Mexico, and Galapagos Islands, and very large in Peru and Chile (NGS 1983, Palmer 1962).

Diagnostic Characteristics: Differs from subspecies *carolinensis* in being larger (e.g., average bill length 347 mm and 312 mm in males and females, respectively, vs. 319 mm and 294 mm) and, in definitive alternate plumage, the brown hindneck being much darker (sometimes almost black) (Palmer 1962). Differs from subspecies *occidentalis* in being much larger (average bill length of *occidentalis* 288 mm and 261 mm, for males and females, respectively) (Palmer 1962).

Reproduction Comments: Along the west coast of North America, egg laying may occur from late winter to early spring (peak usually in March or April but may vary among colonies and from year to year). Subspecies *carolinensis*: southern populations nest irregularly, usually beginning in late fall and extending through June; northernmost populations nest in spring and summer; intermediate populations nest, somewhat irregularly, in winter and spring. Clutch size averages between two and three. Incubation, by both sexes, lasts about 28-30 days. Young leave ground

1 nests at about 35 days, first fly at 71-88 days; leave nests in mangroves at about 63 days. May
2 Some first breed at two years in some colonies (e.g., newly formed ones), possibly not until about
3 four to seven years in stable populations (see Johnsgard 1993). Reproductive success varies
4 with level of disturbance by humans, starvation of young, and/or flooding of nests, but typically
5 the number of young fledged per nest averages one or less. See Johnsgard (1993) for
6 information on productivity. Long-lived; reproduction tends to be "boom or bust." Colonies
7 include up to 150 pairs in Trinidad.

8 **Ecology Comments:** Populations fluctuate considerably from year to year and from place to
9 place.

10 **Non-Migrant:** Yes

11 **Locally Migrant:** Yes

12 **Long Distance Migrant:** Yes

13 **Mobility and Migration Comments:** Many stay close to nesting areas in winter. A portion of the
14 eastern subspecies migrates to Florida, the Caribbean coasts of Colombia and Venezuela, and
15 the Greater Antilles for winter. During cold winters, some Texas breeders winter along the Gulf
16 Coast of Mexico. Individuals from breeding areas north of Florida winter mainly in Florida and
17 Cuba; young and adults from Florida breeding colonies are more sedentary (young generally do
18 not disperse more than 250 km from natal areas, adults may move up to 450-575 km from colony
19 during the nonbreeding season) (Johnsgard 1993).

20 **Marine Habitat(s):** Near shore

21 **Estuarine Habitat(s):** Bay/sound, Lagoon, River mouth/tidal river, Scrub-shrub wetland

22 **Terrestrial Habitat(s):** Bare rock/talus/scree, Cliff, Sand/dune

23 **Habitat Comments:** Mainly coastal, rarely seen inland or far out at sea. Feeds mostly in
24 shallow estuarine waters, less often up to 40 miles from shore. Makes extensive use of sand
25 spits, offshore sand bars, and islets for nocturnal roosting and daily loafing, especially by
26 nonbreeders and during the non-nesting season. Dry roosting sites are essential. Some roosting
27 sites eventually may become nesting areas. **BREEDING:** Nests usually on coastal islands, on
28 the ground or in small bushes and trees (Palmer 1962). Nests on middle or upper parts of steep
29 rocky slopes of small islands in California and Baja California; usually nests on low-lying islands
30 landward of barrier islands or reefs on Atlantic and Gulf coasts, where often nests in mangroves,
31 sometimes in Australian "pines," red-cedars, live oaks, redbays, or sea grapes. In the subtropics
32 and tropics, mangrove vegetation constitutes an important roosting and nesting substrate
33 (Collazo and Klaas 1985, Schreiber 1979, Schreiber and Schreiber 1982). May shift between
34 different breeding sites, apparently in response to changing food supply distribution (Anderson
35 and Gress 1983) and/or to erosion/flooding of nesting sites.

36 **Adult Food Habits:** Piscivore

37 **Immature Food Habits:** Piscivore

38 **Food Comments:** Eats mainly fishes, especially menhaden, mullet, sardines, pinfish, and
39 anchovies in U.S. waters; sometimes euphausiids; dives into water from air (USFWS 1980).
40 Feeds by diving in deeper water, by swimming, sometimes in cooperative groups, in shallower
41 water (Hilty and Brown 1986). Rarely reported scavenging or preying on eggs or young of water
42 birds. Forages in shallow estuarine and inshore waters mostly within 10 km of the coast
43 (Johnsgard 1993).

1 **Adult Phenology:** Crepuscular, Diurnal

2 **Immature Phenology:** Crepuscular, Diurnal

3 **Phenology Comments:** Most activity diurnal, little during twilight.

4 **Colonial Breeder:** Yes

5 **Length:** 122 cm

6 **Weight:** 3636 grams

7 **References:**

- 8 • American Ornithologists' Union (AOU). 1957. The A.O.U. Check-list of North American
9 Birds, 5th ed. Port City Press, Inc., Baltimore, MD. 691 pp.
- 10 • American Ornithologists' Union (AOU). 1983. Check-list of North American Birds, 6th
11 edition. Allen Press, Inc., Lawrence, KS. 877 pp.
- 12 • Anderson, D. W. 1988. Dose-response relationship between human disturbance and
13 brown pelican breeding success. Wildl. Soc. Bull. 16:339-345.
- 14 • Anderson, D. W., and F. Gress. 1983. Status of a northern population of California brown
15 pelicans. Condor 85:79-88.
- 16 • Biosystems Analysis, Inc. 1989. Endangered Species Alert Program Manual: Species
17 Accounts and Procedures. Southern California Edison Environmental Affairs Division.
- 18 • California Department of Fish and Game (CDF&G). 1990. 1989 annual report on the
19 status of California's state listed threatened and endangered plants and animals. 188 pp.
- 20 • Clapp, R. B., and P. A. Buckley. 1984. Status and conservation of seabirds in the
21 southeastern United States. Pages 135-155 in Croxall et al., eds. Status and
22 conservation of the world's seabirds. ICBP Tech. Pub. No. 2.
- 23 • Guravich, D, and J. E. Brown. 1983. The return of the brown pelican. Louisiana State
24 Univ. Press, Baton Rouge. viii + 118 pp.
- 25 • Johnsgard, P. A. 1993. Cormorants, darters, and pelicans of the world. Smithsonian
26 Institute Press, Washington, D.C. xiv + 445 pp.
- 27 • Matthews, J.R. and C.J. Moseley (eds.). 1990. The Official World Wildlife Fund Guide to
28 Endangered Species of North America. Volume 1. Plants, Mammals. xxiii + pp 1-560 +
29 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians,
30 Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180.
31 Beacham Publications, Inc., Washington, D.C.
- 32 • National Geographic Society (NGS). 1983. Field guide to the birds of North America.
33 National Geographic Society, Washington, DC.
- 34 • Palmer, R. S. (editor). 1962. Handbook of North American birds. Vol. 1. Loons through
35 flamingos. Yale University Press, New Haven. 567 pp.
- 36 • Sibley, C.G., and B.L. Monroe, Jr. 1990. Distribution and Taxonomy of Birds of the World.
37 Yale University Press, New Haven, CT. xxiv + 1111 pp.
- 38 • Spendelow, J. A. and S. R. Patton. 1988. National Atlas of Coastal Waterbird Colonies in
39 the Contiguous United States: 1976-1982. U.S. Fish and Wildlife Service, Biological
40 Report 88(5). x + 326 pp.

- 1 • Terres, J. K. 1980. The Audubon Society encyclopedia of North American birds. Alfred A.
2 Knopf, NY.
- 3 • U. S. Fish and Wildlife Service. 1980. Selected vertebrate endangered species of the
4 seacoast of the United States--brown pelican eastern and California subspecies.
5 FWS/OBS-80/01.40.
- 6 • U.S. Fish and Wildlife Service (USFWS). 1990. Endangered and threatened species
7 recovery program: report to Congress. 406 pp.
- 8

Mexican Spotted Owl (*Strix occidentalis lucida*)

The Mexican spotted owl was designated as a federally Threatened species on March 16, 1993.

Historic Range: Range extends from southern Utah (Kertell 1977, Marti 1979) and central Colorado (Webb 1983) south through the mountainous regions of Arizona (Ganey and Balda 1989), New Mexico, western Texas (Guadalupe Mountains), northern Sonora, Chihuahua, and Nuevo Leon south to Michoacan and Puebla (AOU 1983; USFWS 1994, 1995). Mexican occurrences documented during 1990-1993 were in the Sierra Madre Occidental, Sierra Madre Oriental, and Eje Neovolcanico, south to Aguascalientes; the Mexican portion of the range has not been thoroughly surveyed (USFWS 1995). Many populations in Arizona and New Mexico occur in relatively isolated mountain ranges, sometimes separated by large expanses of nonforested habitats; little is known of the populations in many of these mountain ranges; some ranges may include too little habitat to support spotted owl populations indefinitely without periodic immigration from neighboring ranges (Ganey, in Thomas et al. 1990). Abundance (density) is greatest in the central portion of the range; a little more than half of the U.S. population occurs in the Upper Gila Mountains Recovery Unit in Arizona and New Mexico (USFWS 1995). See USFWS (1995) for a spot map showing distribution based on observations made during 1990-1993.

Basic Description: Medium-sized, dark-eyed owl lacking ear tufts.

General Description: A large, dark-eyed, round-headed, brown owl with whitish spotting on the head, back, and underparts (spotted breast, barred belly).

Diagnostic Characteristics: Differs from other subspecies in being generally paler and having the lighter markings of the underparts more whitish (Ridgway 1914).

Reproduction Comments: Egg dates: peak in April in Arizona and New Mexico, sometimes as early as early March. Clutch size is 2-4, usually 2. Incubation, by female (fed by male), lasts about 30 days. Hatching generally occurs in early to mid-May. Young leave nest at about 5 weeks (June), fly at about 6-7 weeks, stay near nest for several weeks, fed by adults until late summer, independent by early fall (dispersal of young occurs in September-October). First breeds at 2-3 years; may not breed every year. Reproductive success generally is low (USFWS 1993); average number of young fledged per pair is about 1.0 (USFWS 1995).

Ecology Comments: Mostly solitary outside the breeding season. Home range size apparently varies with location and habitat; generally the smallest home ranges are a few hundred hectares and the largest ones are about 1500 ha (minimum convex polygon) (see USFWS 1995). In northern Arizona, mean home range of three pairs was 847 ha; owls shifted seasonally such that year-round home range was larger than the range used during any one season (Ganey and Balda 1989). Mean home range size of four pairs in the Lincoln National Forest was 1180 ha; mean home ranges in Utah varied from 242 ha in Zion National Park to 625 ha for two owls elsewhere (see USFWS 1993). In Utah, some home ranges shifted seasonally, others did not (see USFWS 1994). In general, fidelity to territories is apparently high (USFWS 1995). In Utah, seven juveniles dispersed 24-145 km (USFWS 1995). In New Mexico, five juvenile females dispersed 8-56 km (mean 22 km), five juvenile males dispersed 2-13 km (mean 6 km); some females, including an adult, made intermountain movements (Gutierrez et al. 1996). Density generally is less than 0.4/sq km (mostly about 0.1-0.2/sq km) (USFWS 1995). Annual survival rate appears to be about 80-90% in adults, 6-29% in juveniles (White et al. 1995, USFWS 1995).

Non-Migrant: Yes

Locally Migrant: Yes

Long Distance Migrant: No

Mobility and Migration Comments: In the southwestern U.S., apparently largely nonmigratory, with some vertical migration at higher elevations (Ganey et al. 1988) (i.e., owls move to lower elevations for winter, with some exceptions). Some owls remain year-round in the same general areas but exhibit seasonal shifts in habitat use pattern (USFWS 1995). Some migrate 20-50 km between summer and winter ranges (see USFWS 1995).

Palustrine Habitat(s): Riparian

Terrestrial Habitat(s): Cliff, Forest - Conifer, Forest - Hardwood, Forest - Mixed

Special Habitat Factors: Standing snag/hollow tree

Habitat Comments: Highest densities occur in mixed-conifer forests that have experienced minimal human disturbance (USFWS 1995, Ganey and Dick 1995). In the southwestern U.S., most common where unlogged closed canopy forests occur in steep canyons; uneven-aged stands with high basal area and many snags and downed logs are most favorable. In Arizona, occurs primarily in mixed-conifer, pine-oak, and evergreen oak forests; also occurs in ponderosa pine forest and rocky canyonlands (Ganey and Balda 1989). In Arizona, generally foraged more than or as frequently as expected (based on availability) in virgin mixed-conifer and ponderosa pine forests, and less than expected in managed forests; roosted primarily in virgin mixed-conifer forests; both foraging and especially roosting sites had more big logs, higher canopy closure, and greater densities and basal areas of both trees and snags than did random sites (Ganey and Balda 1994). In southern Utah, commonly used mesa tops, benches and warm slopes above canyons in fall and winter; relatively cool canyons were the primary summer habitat (see USFWS 1994). In New Mexico, breeding and roosting occurred in mixed-conifer forests that contained an oak component more frequently than expected by chance; generally did not use pinyon pine-alligator juniper woodlands for nesting or roosting; selected roost and nest sites in forests characterized by mature trees with high variation in tree heights and canopy closure greater than 75% (Seamans and Gutierrez 1995). Basically intolerant of even-age forest management practices (USFWS, Federal Register, 1 April 1994). Requires cool summer roosts (Barrows 1981, Ganey et al. 1993), such as near canyon bottoms, in dense forests, on shady cliffs or in caves (Ganey et al. 1988). Sometimes occurs in deep canyons in areas that lack extensive forests. Sometimes may winter in comparatively open habitats at lower elevations. Breeding formerly occurred in desert riparian habitat, but occurrences are rare in this habitat today. In general, foraging habitat requirements are not well known (USFWS 1995). See USFWS (1993, 1994, 1995) for further details on habitat. Nests on broken tree top, cliff ledge, in natural tree cavity, or in tree on stick platform, often the abandoned nest of hawk or mammal; sometimes in cave. In Utah and Colorado, most nests are in caves or on cliff ledges in steep-walled canyons; elsewhere, nests apparently most often are in trees, especially Douglas-fir (USFWS 1995, Seamans and Gutierrez 1995). Exhibits high level of nest site fidelity. Typically selects cool, shady sites with high canopy closure and at least a few old-growth trees, usually on moderate to steep slopes (USFWS 1993). In New Mexico, 61% of nest structures were on clumps of limbs caused by dwarf mistletoe infections; nest trees averaged 164 years old and 60.6 cm in diameter (Seamans and Gutierrez 1995). See also USFWS (1995).

Adult Food Habits: Carnivore

Immature Food Habits: Carnivore

Food Comments: Diet varies with location; woodrats, mice, and voles are common prey (USFWS 1995, Ward and Block 1995). Zion National Park, Utah: *Neotoma*, *Thomomys*, and beetles (Kertell 1977). Arizona: mainly cottontails, deer mice, woodrats, and voles (Ganey et al. 1988); also various birds, bats, lizards, and snakes (Duncan, 1992, Herpetol. Rev. 23:81). Arizona: mainly *Neotoma*, *Peromyscus*, *Microtus*, *Sylvilagus*, and *Thomomys* (Ganey 1992). Generally hunts from a perch. May cache prey.

1 **Adult Phenology:** Crepuscular, Nocturnal

2 **Immature Phenology:** Crepuscular, Nocturnal

3 **Phenology Comments:** Roosts during the day; hunts at dusk and at night. May leave roost
4 during day to capture prey beneath roost, retrieve cached prey, or to drink or bathe in stream. In
5 northern Arizona, calling peaked in late spring and during 2-hour period following sunset (Ganey
6 1990).

7 **Length:** 45 cm

8 **Restoration Potential:** Recovery plan (USFWS 1995) indicates that delisting could occur within
9 10 years (depends on results of monitoring over that period).

10 **Preserve Selection & Design Considerations:** Preserves should be distributed among the six
11 U.S. and five Mexican recovery units designated by USFWS (1995). This subspecies probably
12 exists as more or less discrete clusters of populations, reflecting the patchiness of the habitat;
13 each cluster of populations (e.g., the Mogollon Rim cluster and the Southern Rockies cluster)
14 apparently can be regarded as a classical metapopulation; owls disperse frequently within
15 clusters but only rarely between clusters (Keitt et al. 1995).

16 **Management Requirements:** Management initially should focus on the alleviation of major
17 threats: catastrophic wildfire and widespread use of even-aged silviculture; thereafter, other
18 priorities, such as creating replacement owl habitat, should be pursued (USFWS 1995, which see
19 for detailed management information). Manipulative experiments are needed to evaluate effects
20 of fire (or other forest management activities) on owls (Bond et al. 2002). See also Dawson et al.
21 (1987) and Lefranc and Glinski (1988) for management and research recommendations. See
22 USFWS (1994) for a review of management policies and practices by agencies and tribes.

23 **Monitoring Requirements:** Monitoring of the population and habitat over the next 10 years is
24 regarded as an essential part of the recovery plan (USFWS 1995). See USFWS (1995) for
25 detailed information on monitoring procedures. See also Bull (1987) for information on capture
26 techniques, Bosakowski (1987) and Forsman (1983) for census methods. See Ganey (1990) for
27 cautions on censusing owls through calling surveys. Paton et al. (1991) concluded that the use of
28 backpack-mounted radio tags should be avoided (due to impaired reproduction and survival of
29 radio-tagged owls).

30 **References:**

- 31 • Allen, Craig R., Stephen Demarais, and R. Scott Lutz. 1994. Red Imported Fire Ant
32 Impact on Wildlife: an Overview. Texas J. Sci. 46(1):51-59.
- 33 • American Ornithologists' Union (AOU). 1957. The A.O.U. Check-list of North American
34 Birds, 5th ed. Port City Press, Inc., Baltimore, MD. 691 pp.
- 35 • American Ornithologists' Union (AOU). 1983. Check-list of North American Birds, 6th
36 edition. Allen Press, Inc., Lawrence, KS. 877 pp.
- 37 • Andrews, R. R. and R. R. Righter. 1992. Colorado Birds. Denver Museum of Natural
38 History, Denver. 442 pp.
- 39 • Barrowclough, G. F., and R. J. Gutierrez. 1990. Genetic variation and differentiation in
40 the spotted owl (*Strix occidentalis*). Auk 107:737-744.
- 41 • Barrows, C.W. 1981. Roost selection by spotted owls: an adaptation to heat stress.
42 Condor 83:302-309.

- 1 • Bond, M. L., R. J. Gutiérrez, A. B. Franklin, W. S. LaHaye, C. A. May, and M. E.
2 Seamans. 2002. Short-term effects of wildfires on spotted owl survival, site fidelity, mate
3 fidelity, and reproductive success. *Wildlife Society Bulletin* 30:1022-1028.
- 4 • Bosakowski, T. 1987. Census of barred owls and spotted owls. Pages 307-308 in Nero,
5 R. W., et al., eds. *Biology and conservation of northern forest owls*. USDA For. Serv.,
6 Gen. Tech. Rep. RM-142.
- 7 • Bull, E. L., et al. 1987. Nest platforms for great gray owls. Pp. 87-90 in Nero, R. W., et al.,
8 eds. *Biology and conservation of northern forest owls*. USDA Forest Service, Gen.Tech.
9 Rep. RM-142.
- 10 • Bull, E.L. 1987. Capture techniques for owls. Pages 291-293 in R.W. Nero, R.J. Clark,
11 R.J. Knapton, and R.H. Hamre, editors. *Biology and conservation of northern forest owls*.
12 USDA Forest Service, Gen. Tech. Rep. RM-142.
- 13 • Campbell, L. 1995. *Endangered and Threatened Animals of Texas: Their Life History and*
14 *Management*. Texas Parks and Wildlife Department, Endangered Resources Branch,
15 Austin, Texas. ix + 129 pp.
- 16 • Dawson, W. R., J. D. Ligon, and J. R. Murphy. 1987. Report of the Scientific Advisory
17 Panel on the Spotted Owl. *Condor* 89:205-229.
- 18 • Division of Natural Resources, Navajo Fish and Wildlife Department. 1995. *Endangered*
19 *Species List for The Navajo Nation*.
- 20 • Fletcher, K.W. and H.E. Hollis. 1994. Habitats used, abundance, and distribution of the
21 Mexican spotted owl, *Strix occidentalis lucida*, on National Forest System lands in the
22 Southwestern Region. Southwestern Regional Office, U.S. Forest Service.
- 23 • Forsman, E. D. 1983. Methods and materials for locating and studying spotted owls.
24 U.S.D.A. Forest Service, Gen. Tech. Report PNW-162. Portland, Oregon. 8 pp.
- 25 • Forsman, E., and E. C. Meslow. 1986. The spotted owl. Pages 742-761 in R. L. Di
26 Silvestro, editor. *Audubon Wildlife Report 1986*. National Audubon Society, NY. 1094 pp.
- 27 • Ganey, J. L. 1990. Calling behavior of spotted owls in northern Arizona. *Condor* 92:485-
28 490.
- 29 • Ganey, J. L. 1992. Food habits of Mexican spotted owls in Arizona. *Wilson Bull.* 104:321-
30 326.
- 31 • Ganey, J. L. et al. 1988. Mexican spotted owl. Pages 145-150 in Glinski et al., eds. *Proc.*
32 *Southwest raptor management symposium and workshop*. Nat. Wildl. Fed. Sci. and Tech.
33 Ser. No. 11.
- 34 • Ganey, J. L., R. P. Balda, and R. M. King. 1993. Metabolic rate and evaporative water
35 loss of Mexican spotted and great horned owls. *Wilson Bull.* 105:645-656.
- 36 • Ganey, J. L., and J. L. Dick, Jr. 1995. Habitat relationships of the Mexican spotted owl:
37 current knowledge. Pages 1-42 in U.S. Fish and Wildlife Service. *Mexican spotted owl*
38 *recovery plan*. Volume II. Albuquerque, NM.
- 39 • Ganey, J. L., and R. P. Balda. 1989. Distribution and habitat use of Mexican spotted owls
40 in Arizona. *Condor* 91:355-361.
- 41 • Ganey, J. L., and R. P. Balda. 1989. Home-range characteristics of spotted owls in
42 northern Arizona. *J. Wildlife Management* 53:1159-1165.
- 43 • Ganey, J. L., and R. P. Balda. 1994. Habitat selection by Mexican spotted owls in
44 northern Arizona. *Auk* 111:162-169.
- 45 • Ganey, J.L. 1990. Calling behavior of spotted owls in northern Arizona. *The Condor*
46 92:485-490.

- 1 • Ganey, J.L. 1992. Food habits of Mexican spotted owls in Arizona. Wilson Bull.
2 104(2):321-326.
- 3 • Ganey, J.L. and R.P. Balda. 1989. Distribution and habitat use of Mexican spotted owls in
4 Arizona. The Condor 91:355-361.
- 5 • Ganey, J.L. and R.P. Balda. 1989. Home-range characteristics of spotted owls in
6 northern Arizona. J. Wildl. Manage 53(4):1159-1165.
- 7 • Ganey, J.L., et.al. 1993. Metabolic rate and evaporative water loss of Mexican spotted
8 and great horned owls. Wilson Bull. 105(4):645-656.
- 9 • Gutierrez, R. J., M. E. Seamans, and M. Z. Peery. 1996. Intermountain movement by
10 Mexican spotted owls (*Strix occidentalis lucida*). Great Basin Naturalist 56:87-89.
- 11 • Hayward, C.L., et.al. 1976. Birds of Utah. Great Basin Naturalist Memoirs, No. 1, Brigham
12 Young University, Provo, UT.
- 13 • Johnson, C. L. 1997. Distribution, Habitat, and Ecology of the Mexican Spotted Owl in
14 Colorado. Masters Thesis, University of Northern Colorado, Greeley.
- 15 • Keitt, T., A. B. Franklin, and J. P. Ward, Jr. 1995. Landscape analysis and
16 metapopulation structure. Pages 1-16 in U.S. Fish and Wildlife Service. Mexican spotted
17 owl recovery plan. Volume II. Albuquerque, NM.
- 18 • Kertell, K. 1977. The spotted owl at Zion National Park, Utah. Western Birds 8:147-150.
- 19 • LaRue, C.T. 1994. Birds of northern Black Mesa, Navajo County, Arizona. Great Basin
20 Naturalist 54(1):1-63.
- 21 • Lefranc, M. N., Jr., and R. L. Glinski. 1988. Southwest raptor management issues and
22 recommendations. Pages 375-392 in Glinski et al., eds. Proc. Southwest raptor
23 management symposium and workshop. National Wildlife Federation Science and Tech.
24 Ser. No. 11.
- 25 • Marti, C. D. 1979. Status of owls in Utah. Pages 29-35 in P. Schaeffer and S. Ehlers, eds.
26 Owls of the West: their ecology and conservation. National Audubon Society, Western
27 Education Center, Tiburon, CA.
- 28 • McDonald, C.B., et.al. 1991. Mexican spotted owl, *Strix occidentalis lucida*, status review.
29 U.S. Fish and Wildlife Service Endangered Species Report 20, Albuquerque, NM.
- 30 • Nelson, D. 1993. Colorado Bird Atlas: Manual on Use of Breeding Codes. Denver
31 Museum of Natural History, Denver. 27 pp.
- 32 • Paton, P. W. C., C. J. Zabel, D. L. Neal, G. N. Steger, N. G. Tilghman, and B. R. Noon.
33 1991. Effects of radio tags on Spotted Owls. J. Wildlife Management 55:617-622.
- 34 • Ridgway, R. 1914. The birds of North and Middle America. Part VI. U.S. National
35 Museum Bull. 50.
- 36 • Rinkevich, S. E. 1995. Mexican spotted owl draft recovery plan. Endangered Species
37 Bulletin 20(4):12-13.
- 38 • Rinkevich, S. E., J. L. Ganey, J. P. Ward, Jr., G. C. White, D. L. Urban, A. B. Franklin, W.
39 M. Block, and F. Clemente. 1995. General biology and ecological relationships of the
40 Mexican spotted owl. Pp. 19-35 in W. M. Block et al., Recovery plan for the Mexican
41 spotted owl (*Strix occidentalis lucida*). U. S. Fish and Wildlife Service, Albuquerque, NM.
42 172 pp.
- 43 • Ryke, N., D. Winters, L. McMartin and S. Vest. 1994. Threatened, Endangered and
44 Sensitive Species of the Pike and San Isabel National Forests and Comanche and
45 Cimarron National Grasslands. May 25, 1994.

- 1 • Seamans, M. E., R. J. Gutiérrez, C. A. May, and M. Z. Peery. 1999. Demography of two
2 Mexican spotted owl populations. *Conservation Biology* 13:744-754.
- 3 • Seamans, M. E., and R. J. Gutierrez. 1995. Breeding habitat of the Mexican spotted owl
4 in the Tularosa Mountains, New Mexico. *Condor* 97:944-952.
- 5 • Spahr, R., L. Armstrong, D. Atwood, and M. Rath. 1991. Threatened, endangered, and
6 sensitive species of the Intermountain Region. U.S. Forest Service, Ogden, Utah.
- 7 • Thomas, J. W., E. D. Forsman, J. B. Lint, E. C. Meslow, B. R. Noon, and J. Verner. 1990.
8 A conservation strategy for the northern spotted owl: a report to the Interagency Scientific
9 Committee to address the conservation of the northern spotted owl. U.S. Forest Service,
10 U.S. Fish and Wildlife Service, and National Park Service, Washington, D.C. 427 pp.
- 11 • U.S. Fish and Wildlife Service (USFWS). 1993. Notice of 90-day finding on petition to
12 remove the Mexican spotted owl from the list of endangered and threatened wildlife and
13 plants. *Federal Register* 58(183):49467-49468. 23 September 1993.
- 14 • U.S. Fish and Wildlife Service (USFWS). 31 August 2004. Final designation of critical
15 habitat for the Mexican spotted owl. *Federal Register* 69(168):53182-53231.
- 16 • U.S. Fish and Wildlife Service. 1993. Final rule to list the Mexican spotted owl as a
17 threatened species. *Federal Register* 58(49):14248-14271. 16 March 1993.
- 18 • U.S. Fish and Wildlife Service. 1994. Proposed determination of critical habitat for the
19 Mexican spotted owl. *Federal Register* 59(234):63162-63201. 7 December 1994.
- 20 • U.S. Fish and Wildlife Service. 1995. Draft recovery plan for the Mexican spotted owl.
21 xxvi + 646 pp.
- 22 • U.S. Fish and Wildlife Service. 1995. Recovery plan for the Mexican spotted owl (*Strix*
23 *occidentalis lucida*). Volume 1. Albuquerque, NM. 172 pp.
- 24 • Ward, J. P., Jr., A. B. Franklin, S. E. Rinkevich, and F. Clemete. 1995. Distribution and
25 abundance of Mexican spotted owls. Pages 1-14 in U.S. Fish and Wildlife Service.
26 Mexican spotted owl recovery plan. Volume II. Albuquerque, NM.
- 27 • Ward, J. P., Jr., and W. M. Block. 1995. Mexican spotted owl prey ecology. Pages 1-48 in
28 U.S. Fish and Wildlife Service. Mexican spotted owl recovery plan. Volume II.
29 Albuquerque, New Mexico.
- 30 • Webb, B. 1983. Distribution and nesting requirements of montane forest owls in
31 Colorado. Part IV. Spotted owl (*Strix occidentalis*). *Colorado Field Ornithologists' Journal*
32 17:2-8.
- 33 • White, G. C., A. B. Franklin, and J. P. Ward, Jr. 1995. Population biology. Pages 1-25 in
34 U.S. Fish and Wildlife Service. Mexican spotted owl recovery plan. Volume II.
35 Albuquerque, NM.
- 36 • Zarn, M. 1974. Spotted owl (*Strix occidentalis*). Habitat Management Series for Unique or
37 Endangered Species Report No. 10, U.S. Bureau of Land Management Technical Note
38 242, Denver, CO. 22 p.

Ocelot (*Felis pardalis*)

The ocelot was designated as a federally Endangered species on July 21, 1982.

Historic Range: Historical range: Texas, Louisiana, Arkansas, and Arizona south through Mexico, Central America, and South America to eastern Peru, eastern Bolivia, Paraguay, Uruguay, and northern Argentina. Occurs in the mountains of Colombia, Ecuador, and northern Peru, but not on the high plateaus of southern Peru and Bolivia (Kitchener 1991); recently recorded in Uruguay (see Kitchener 1991); to elevations of 1000 m. In the U.S., currently found regularly only in southern Texas (e.g., Laguna Atascosa National Wildlife Refuge, site of a recent radiotelemetry study). Occurrence in Arizona is based only on a few old records from the vicinity of Fort Verde and (b) (7)(E) (Hoffmeister 1986); documentation for these records is less than ideal.

Basic Description: A cat (ocelot).

General Description: A small spotted cat with a long tail; ground color ranges from whitish or tawny yellow to reddish gray and gray; dark markings form chainlike streaks, generally forming black-bordered elongated spots, which run obliquely down the sides; adult total length 92-137 cm, tail length 27-40 cm; mass 11-16 kg; greatest length of skull of adults, 120-158 mm (Hall 1981, Nowak 1991).

Diagnostic Characteristics: Differs from the jaguar in much smaller size (jaguar is 157-242 cm in total length) and pelage spots not forming distinct rosettes. Differs from *Felis wiedii* and *F. tigrina* in being larger (hind foot longer than 145 mm vs. shorter, greatest length of skull more than 120 mm vs. shorter, length of P4 more than 12.7 mm vs. shorter) (Hall 1981). Differs from young mountain lion in having spots arranged in rows or in a chainlike pattern.

Reproduction Comments: Texas: breeds in late summer. Births occur in fall and winter in Texas and Mexico (Leopold 1959). Tropics: breeds year-round. Gestation lasts about 70 days. Litter size is 2-4 (usually 2).

Ecology Comments: Population density in Costa Rica was estimated at 14-25/100 sq km (Kitchener 1991). In Brazil, Trolle and Kery (2003) used capture-recapture analysis of camera-trapping data to estimate density at 2.82 independent individuals per 5 sq km.

Non-Migrant: Yes

Locally Migrant: No

Long Distance Migrant: No

Mobility and Migration Comments: Home range in Texas reportedly is a few square kilometers (Kitchener 1991). In Peru, adult females occupied exclusive home ranges of about 2 sq km; male ranges were several times larger, exclusive of those of other males, and overlapped multiple female ranges; individuals often were solitary but appeared to make contact with others frequently (Emmons 1988).

Palustrine Habitat(s): FORESTED WETLAND, Riparian

Terrestrial Habitat(s): Forest - Hardwood, Savanna, Shrubland/chaparral, Woodland - Hardwood

Special Habitat Factors: Standing snag/hollow tree

Habitat Comments: Habitats with good cover; when active by day, tends to keep hidden in dense brush (Emmons and Feer 1990). Inhabits dense chaparral thickets in Texas. Elsewhere, occurs in humid tropical forests, mangrove forests, swampy savannas, brushland, and riverine scrub in deserts. Where not hunted, adapts well to disturbed habitats around villages; often uses man-made trails (Emmons and Feer 1990). Mainly terrestrial but climbs, jumps, and swims well (Nowak 1991). Dens are in caves, hollow trees, thickets, or the spaces between the closed buttress roots of large trees; rarely climbs but sometimes may sleep on tree branch.

Adult Food Habits: Carnivore

Immature Food Habits: Carnivore

Food Comments: Feeds on various small to moderate-sized vertebrates: rodents, rabbits, and other small mammals; young deer and peccaries; birds (sometimes including domestic poultry); snakes; lizards; fishes; etc. Hunts and captures prey on the ground (Emmons and Feer 1990).

Adult Phenology: Crepuscular, Nocturnal

Immature Phenology: Crepuscular, Nocturnal

Phenology Comments: Nocturnal and diurnal; mainly nocturnal (Emmons and Feer 1990).

Length: 125 cm

Weight: 14000 grams

References:

- Allen, Craig R., Stephen Demarais, and R. Scott Lutz. 1994. Red Imported Fire Ant Impact on Wildlife: an Overview. Texas J. Sci. 46(1):51-59.
- Albuja, L. 1991. Lista de vertebrados del Ecuador. Escuela Politécnica XVI:163-203.
- Anderson, S. 1997. Mammals of Bolivia, taxonomy and distribution. Bulletin of the American Museum of Natural History, No 231.
- Campbell, L. 1995. Endangered and Threatened Animals of Texas: Their Life History and Management. Texas Parks and Wildlife Department, Endangered Resources Branch, Austin, TX. ix + 129 pp.
- Carrillo, E., G. Wong, and A. Cuarón. 2000. Monitoring mammal populations in Costa Rican protected areas under different hunting restrictions. Conservation Biology 14:1580-1591.
- Davis, W. B. 1978. The mammals of Texas. Texas Parks and Wildlife Dept., Bull. No. 41. 294 pp.
- Emmons, L. H. 1988. A field study of ocelots (*Felis pardalis*) in Peru. Rev. Ecol. (Terre Vie) 43:133-157.
- Emmons, L. H. and F. Feer. 1990. Neotropical rainforest mammals: a field guide. Univ. of Chicago Press. xiv + 281 pp.
- Engstrom, M. and B. Lim. 2000. Checklist of the mammals of Guyana. Smithsonian Institution. <http://www.mnh.si.edu/biodiversity/bdg/guymammals.html>. \Escamilla, A., M. Sanvicente, M. Sosa, and C. Galindo-Leal. 2000. Habitat mosaic, wildlife availability, and hunting in the tropical forest of Calakmul, Mexico. Conservation Biology 14:1592-1601.
- Foster, M.L. and S.R. Humphrey. 1995. Use of highway underpasses by Florida panthers and other wildlife. Wildl. Soc. Bull. 23(1):95-100.

- 1 • Hall, E. R. 1981. The Mammals of North America, 2nd. Ed., 2 Vols. John Wiley and Sons,
2 New York, NY. 1181 pp.
- 3 • Hall, E. Raymond. 1981. The Mammals of North America, Vols. I & II. John Wiley & Sons,
4 New York, NY. 1181 pp.
- 5 • Hoffmeister, D. F. 1986. Mammals of Arizona. Univ. Arizona Press and Arizona Game
6 and Fish Dept. 602 pp.
- 7 • Jones, C., R. S. Hoffman, D. W. Rice, M. D. Engstrom, R. D. Bradley, D. J. Schmidly, C.
8 A. Jones, and R. J. Baker. 1997. Revised checklist of North American mammals north of
9 Mexico, 1997. Occasional Papers, Museum of Texas Tech University 173:1-20.
- 10 • Jones, J. K., Jr., R. S. Hoffman, D. W. Rice, C. Jones, R. J. Baker, and M. D. Engstrom.
11 1992. Revised checklist of North American mammals north of Mexico, 1991. Occasional
12 Papers, The Museum, Texas Tech University, 146:1-23.
- 13 • Kitchener, A. 1991. The natural history of the wild cats. Comstock Publishing Associates,
14 Cornell Univ. Press, Ithaca. xxi + 280 pp.
- 15 • Lowery, G. H., Jr. 1974. The mammals of Louisiana and its adjacent waters. Louisiana
16 State University Press, Baton Rouge. 565 pp.
- 17 • Mares, M. A., R. M. Báñez, J. K. Braun, and R. A. Ojeda. 1996. Observations on the
18 mammals of Tucumán Province, Argentina. I. Systematics, distribution, and ecology of
19 the Didelphimorphia, Xenarthra, Chiroptera, Primates, Carnivora, Perissodactyla,
20 Artiodactyla, and Lagomorpha. Annals of the Carnegie Museum 65:89-152.
- 21 • Matthews, J.R. and C.J. Moseley (eds.). 1990. The Official World Wildlife Fund Guide to
22 Endangered Species of North America. Volume 1. Plants, Mammals. xxiii + pp 1-560 +
23 33 pp. appendix + 6 pp. glossary + 16 pp. index. Volume 2. Birds, Reptiles, Amphibians,
24 Fishes, Mussels, Crustaceans, Snails, Insects, and Arachnids. xiii + pp. 561-1180.
25 Beacham Publications, Inc., Washington, D.C.
- 26 • Murray, J. L., and G. L. Gardner. 1997. *Leopardus pardalis*. Mammalian Species (548):1-
27 10.
- 28 • Nowak, R. M. 1991. Walker's mammals of the world. Fifth edition. Vols. I and II. Johns
29 Hopkins Univ. Press, Baltimore, MD. 1629 pp.
- 30 • Nowell, K., and P. Jackson. 1996. Wild cats: Status survey and conservation action plan.
31 Gland, Switzerland, IUCN.
- 32 • Pacheco, V., H. de Macedo, E. Vivar, C. Ascorra, R. Arana-Cardó, and S. Solari. 1995.
33 Lista anotada de los mamíferos peruanos. Conservation International, Washington, DC.
- 34 • Pacheco, V., S. Solari, E. Vivar, and P. J. Hocking. 1994. La riqueza biológica del Parque
35 Nacional Yanachaga - Chemillén. Magistri et Doctores 7:3-6.
- 36 • Russell, Dennis N. 1971?. History and status of the felids of Texas. Presentation at
37 "Symposium on the Status and Management of the Wild Cats of North America"; North
38 American Wildlife and Natural Resources Conference; Portland, Oregon; A Contribution
39 of Federal Aid in Wildlife Restoration Project Texas W-103-R.
- 40 • Sealander, J.A. and G.A. Heidt. 1990. Arkansas Mammals: Their Natural History,
41 Classification and Distribution. University of Arkansas Press, Fayetteville. 308 pp.
- 42 • Tirira, D. 1999. Mamíferos del Ecuador. Museo de Zoología, Pontificia Universidad
43 Católica del Ecuador, Quito.
- 44 • Trolle, M., and M. Kery. 2003. Estimation of ocelot density in the Pantanal using capture-
45 recapture analysis of camera-trapping data. Journal of Mammalogy 84:607-614.

- 1 • U.S. Fish and Wildlife Service (USFWS). 1990. Endangered and threatened species
2 recovery program: report to Congress. 406 pp.
- 3 • Wilson, D. E., and D. M. Reeder (editors). 1993. Mammal Species of the World: a
4 Taxonomic and Geographic Reference. Second Edition. Smithsonian Institution Press,
5 Washington, DC. xviii + 1206 pp. Available online at: <http://www.nmnh.si.edu/msw/>.
- 6 • Wilson, D. E., and S. Ruff. 1999. The Smithsonian book of North American mammals.
7 Smithsonian Institution Press, Washington, D.C. 750 pp.

Lesser Long-nosed Bat (*Leptonycteris curasoe*)

1 The lesser long-nosed bat was designated as a federally Endangered species on September 30,
2 1988.

3 **Distribution:** Central California (Constantine 1998), southern Arizona (e.g., Sidner and Davis
4 1988), and New Mexico to Honduras and El Salvador (Simmons, in Wilson and Reeder 2005).
5 U.S. populations apparently winter in Mexico.

6 **Habitat:** The habitat in Mexico is primarily tropical deciduous forest and thorn forest (Arita 1991).
7 In the United States, this bat roosts in old mines and caves at the base of mountains near alluvial
8 fans vegetated with agave, yucca, saguaro, and organ pipe cactus (Barbour and Davis 1969).
9 Young are born in maternity colonies in caves and mines.

10 **Diet:** Frugivore, Nectarivore

11 **Threats:** USFWS (1987, 1989) stated that the species was threatened by disturbance of roosts,
12 loss of food sources through land clearing and human exploitation, and direct killing by humans.
13 Overall, however, this species does not appear to be very threatened.

14 **References:**

- 15 • Arita, H. T. 1991. Spatial segregation in long-nosed bats, *Leptonycteris nivalis* and
16 *Leptonycteris curasoe*, in Mexico. *Journal of Mammalogy* 72:706-714.
- 17 • Arita, H. T., and S. R. Humphrey. 1988. Revisión taxonómica de los murciélagos
18 magueyeros del género *Leptonycteris* (*Chiroptera: Phyllostomidae*). *Acta Zoologica*
19 *Mexicana* (n.s.) 29:1-60.
- 20 • Cockrum, E. L., and Y. Petryszyn. 1991. The long-nosed bat, *Leptonycteris*: an
21 endangered species in the southwest? *Occas. Pap. Mus. Texas Tech Univ.* (142):1-32.
- 22 • Davis, W. B., and D. C. Carter. 1962. Review of the genus *Leptonycteris* (*Mammalia:*
23 *Chiroptera*). *Proceedings Biological Society Washington* 75:193-197.
- 24 • Fleming, T. H. 1989. Climb every cactus. *Bats* 7(3):3-6.
- 25 • Fleming, T. H. 1991. Following the nectar trail. *Bats* 9(4):4-7.
- 26 • Jones, J. K., Jr., R. S. Hoffman, D. W. Rice, C. Jones, R. J. Baker, and M. D. Engstrom.
27 1992. Revised checklist of North American mammals north of Mexico, 1991. *Occasional*
28 *Papers, The Museum, Texas Tech University*, 146:1-23.
- 29 • Nabhan, G. P., and T. Fleming. 1993. The conservation of New World mutualisms.
30 *Conservation Biology* 7:457-459.
- 31 • U.S. Fish and Wildlife Service (USFWS). 1994. Endangered and Threatened Wildlife and
32 Plants; Animal Candidate Review for Listing as Endangered or Threatened Species.
33 Proposed Rule. *Federal Register*, 59(219): 58982-59028.
- 34 • Wilson, D. E., and D. M. Reeder (editors). 2005. *Mammal species of the world: a*
35 *taxonomic and geographic reference*. Third edition. The Johns Hopkins University Press,
36 Baltimore. Two volumes. 2,142 pp.

37

Jaguar (*Panthera onca*)

The jaguar was designated as a federally Endangered species on July 22, 1997.

Historic Range: The jaguar once ranged throughout tropical lowlands of Mexico, Central America (now very rare except in Belize), and South America (to northern Argentina); in the United States, there are records from southern California, Arizona (Hoffmeister 1986, Johnson and Van Pelt 1997), New Mexico (Findley et al. 1975, Frey 2004), Texas (Schmidly 2004), and perhaps farther east in Louisiana; most records are from Arizona, where a minimum of 64 jaguars have been killed since 1900; some believe that a breeding population formerly existed in portions of the southwestern United States (Federal Register, 13 July 1994, 22 July 1997, which see for a state-by-state review of records). The species is now absent from much of the former range; it has been extirpated as a resident in most or all of the northern extent of the range in the southwestern United States and northern Mexico (see Federal Register, 13 July 1994, p. 35676, for discussion of recent records), El Salvador, Uruguay, developed areas of Brazilian coast, all but the northernmost parts of Argentina, and elsewhere. The largest remaining population is in Amazonian Brazil (Seymour 1989). In recent decades, jaguars occasionally have strayed into the United States in southern Arizona-New Mexico.

Basic Description: A large cat (jaguar).

Reproduction Comments: In tropical areas may breed throughout the year; births most common November-December in Paraguay, December-May in Brazil, March-July in Argentina, July-September in Mexico, June-August in Belize. Gestation lasts about 90-115 days. Litter size is 1-4 (average 2). Young begin to eat meat at about 10-11 weeks, though may suckle 5-6 months; remain in den about 1.5-2 months; stay with mother 1.5-2 year; females sexually mature in 2-3 years, males in 3-4 years (Seymour 1989).

Ecology Comments: Solitary and somewhat territorial, except during breeding season. Density estimated at 4/137 sq km in Brazil, 25-30 per 250 sq km in Belize (Seymour 1989). In Belize, daily home range may be only a few sq km, but may shift to new area every week or two. Home range in Brazil was estimated at 25-76 sq km (see Kitchener 1991). Major cause of mortality is hunting by humans.

Non-Migrant: Yes

Locally Migrant: No

Long Distance Migrant: No

Palustrine Habitat(s): Riparian

Terrestrial Habitat(s): Forest - Hardwood, Forest - Mixed, Grassland/herbaceous, Shrubland/chaparral, Woodland - Hardwood, Woodland - Mixed

Habitat Comments: Habitat includes a wide variety of situations, such as tropical and subtropical forests, lowland scrub and woodland, thorn scrub, pampas/llanos, desert, swampy savanna, mangrove swamps, lagoons, marshland, and floating islands of vegetation. At the southern extreme of the range, this cat inhabits open savanna, flooded grasslands, and desert mountains; at the northern extreme it may be found in chaparral and timbered areas. Young are born in a sheltered place such as a cave or thicket, under an uprooted tree, among rocks, or under a river bank (Seymour 1989).

Adult Food Habits: Carnivore

Immature Food Habits: Carnivore

Food Comments: Feeds on large and small mammals, reptiles and ground-nesting birds. Known to feed on peccaries, capybaras, tapirs, agoutis, deer, small crocodilians and turtles; opportunistic, see Seymour (1989) for further details. Hunts mostly on ground but may pounce on prey from tree or ledge.

Phenology Comments: Active throughout the year. Hunts primarily at night, but may be active day or night (Seymour 1989).

Length: 242 cm

Weight: 136000 grams

References:

- Allen, Craig R., Stephen Demarais, and R. Scott Lutz. 1994. Red Imported Fire Ant Impact on Wildlife: an Overview. Texas J. Sci. 46(1):51-59.
- Brown, D. E. 1983. On the status of the jaguar in the Southwest. Southwestern Naturalist 28:459-460.
- Cadena, A., R. P. Anderson, and P. Rivas-Pava. 1998. Colombian mammals from the chocoan slopes of Nariño. Occasional Papers Museum of Texas Tech University 180:1-15.
- Carrillo, E., G. Wong, and A. Cuarón. 2000. Monitoring mammal populations in Costa Rican protected areas under different hunting restrictions. Conservation Biology 14:1580-1591.
- Engstrom, M. and B. Lim. 2000. Checklist of the mammals of Guyana. Smithsonian Institute. <http://www.mnh.si.edu/biodiversity/bdg/guymammals.html>.
- Escamilla, A., M. Sanvicente, M. Sosa, and C. Galindo-Leal. 2000. Habitat mosaic, wildlife availability, and hunting in the tropical forest of Calakmul, Mexico. Conservation Biology 14:1592-1601.
- Findley, J. S., A. H. Harris, D. E. Wilson, and C. Jones. 1975. Mammals of New Mexico. Univ. of New Mexico Press, Albuquerque, NM. 360 pp.
- Fish and Wildlife Service. 1980. Selected vertebrate endangered species of the seacoast of the United States- the jaguar. FWS/OBS-80/01.41., Slidell.
- Frey, J. K. 2004. Taxonomy and distribution of the mammals of New Mexico: an annotated checklist. Museum of Texas Tech University Occasional Papers 240. 32 pp.
- Hall, E. Raymond. 1981. The Mammals of North America, Vols. I & II. John Wiley & Sons, New York, New York. 1181 pp.
- Hoffmeister, D. F. 1986. Mammals of Arizona. Univ. Arizona Press and Arizona Game and Fish Dept. 602 pp.
- Honacki, James H., et al, eds. 1982. Mammal Species of the World: A Taxonomic and Geographical Reference. Allen Press, Inc. and The Association of Systematics Collections, Lawrence, KS. 694 p.
- Johnson, T. B., and W. E. Van Pelt. 1997. Conservation assessment and strategy for the jaguar in Arizona and New Mexico. Technical Report 105, Nongame and Endangered Wildlife Program Program, Arizona Game and Fish Department, Phoenix, AZ. 24 pp.
- Jones, J. K., Jr., R. S. Hoffman, D. W. Rice, C. Jones, R. J. Baker, and M. D. Engstrom. 1992. Revised checklist of North American mammals north of Mexico, 1991. Occasional Papers, The Museum, Texas Tech University, 146:1-23.

- 1 • Kitchener, A. 1991. The natural history of the wild cats. Comstock Publishing Associates,
2 Cornell Univ. Press, Ithaca, NY. xxi + 280 pp.
- 3 • Lowery, George H. 1974. The mammals of Louisiana and its adjacent waters. Kingsport
4 Press, Inc. Kingsport, TN. 565 pp.
- 5 • Mammalian Species, nos. 1-604. Published by the American Society of Mammalogists.
- 6 • Medellín, R. A., C. Chetkiewicz, A. Rabinowitz, K. H. Redford, J. G. Robinson, E.W.
7 Sanderson, and A. Taber. 2000. El Jaguar en el nuevo milenio. Una evaluación de su
8 estado, detección de prioridades y recomendaciones para la conservación de los
9 jaguares en América. Universidad Nacional Autónoma de México/Wildlife Conservation
10 Society. México D. F., México.
- 11 • New Mexico Department of Game and Fish. 1996. October 1-last update. Fish and
12 Wildlife Information Exchange-VA Tech. Online. Available:
13 <http://www.fw.vt.edu/fishex/nm.html>. Accessed 1997, April 8.
- 14 • Nowak, R. 1994. Jaguars in the United States. Endangered Species Technical Bulletin
15 19(5):6.
- 16 • Nowak, R. M. 1991. Walker's mammals of the world. Fifth edition. Vols. I and II. Johns
17 Hopkins Univ. Press, Baltimore. 1629 pp.
- 18 • Nowak, R. M. 1999. Walker's mammals of the world. Johns Hopkins University Press,
19 Baltimore, MD. Two volumes, 1,936 pp.
- 20 • Nowell, K. and P. Jackson (compilers and editors). 1996. Wild cats. Status survey and
21 conservation action plan. IUCN/SSC Cat Specialist Group. IUCN, Gland, Switzerland.
- 22 • Pacheco, V., H. de Macedo, E. Vivar, C. Ascorra, R. Arana-Cardó, and S. Solari. 1995.
23 Lista anotada de los mamíferos peruanos. Conservation International, Washington, DC.
- 24 • Perry, R. 1970. The world of the jaguar. 168 pp.
- 25 • Phelps, J. Jaguar (*Panthera onca*). Unpublished article prepared for University of Arizona
26 Extension Service in house Beaver Transplant Meeting. Game Branch, Wildlife
27 Management Division, Arizona Game and Fish Department.
- 28 • Quigley, H. B., and P. G. Crawshaw, Jr. 1992. A conservation plan for the jaguar
29 *Panthera onca* in the Pantanal region of Brazil. Biological Conservation 61:149-157.
- 30 • Russell, Dennis N. 1971?. History and status of the felids of Texas. Presentation at
31 "Symposium on the Status and Management of the Wild Cats of North America"; North
32 American Wildlife and Natural Resources Conference; Portland, Oregon; A Contribution
33 of Federal Aid in Wildlife Restoration Project Texas W-103-R.
- 34 • Redford, K. H. and J. F. Eisenberg. 1992. Mammals of the Neotropics, The Southern
35 Cone, Volume 2, Chile, Argentina, Uruguay, Paraguay. Chicago, The University of
36 Chicago Press.
- 37 • Sanderson, E. W., K. H. Redford, C.-L. B. Chetkiewicz, R. A. Medellín, A. R. Rabinowitz,
38 J. G. Robinson, and A. B. Taber. 2002. Planning to save a species: the jaguar as a
39 model. Conservation Biology 16:58-72.
- 40 • Schmidly, D. J. 2004. The mammals of Texas. Revised edition. University of Texas
41 Press, Austin. xviii + 501 pp.
- 42 • Seymour, K. L. 1989. *Panthera onca*. Am. Soc. Mamm., Mammalian Species No. 340:1-
43 9.
- 44 • Silver, S. C., L.E.T. Ostro, L. K. Marsh, L. Maffei, A. J. Noss, M. J. Kelly, R. B. Wallace,
45 H. Gómez, and G. Ayala. 2004. The use of camera traps for estimating jaguar *Panthera*
46 *onca* abundance and density using capture/recapture analysis. Oryx 38:148-154.

- 1 • Swank, W. G., and J. G. Teer. 1989. Status of the jaguar - 1987. *Oryx* 23(1):14-21.
- 2 • Tirira, D. 1999. Mamíferos del Ecuador. Museo de Zoología, Pontificia Universidad
- 3 Católica del Ecuador, Quito.
- 4 • Van Pelt, W. E., and T. B. Johnson. [In prep.]. Conservation agreement for the jaguar in
- 5 Arizona and New Mexico. Nongame and Endangered Wildlife Program Technical Report
- 6 105. Arizona Game and Fish Department, Phoenix, Arizona.
- 7 • Wallace, R. B., H. Gomez, G. Ayala, and F. Espinoza. 2003. Camera trapping for jaguar
- 8 (*Panthera onca*) in the Tuichi Valley, Bolivia. *Mastozoología Neotropical* 10:133-139.
- 9 • Weber, W., and A. Rabinowitz. 1996. A global perspective on large carnivore
- 10 conservation. *Conservation Biology* 10:1046-1053.
- 11 • Whitaker, J. O., Jr. 1980. The Audubon Society field guide to North American mammals.
- 12 Alfred A. Knopf, New York. 745 pp.
- 13 • Wilson, D. E., and D. M. Reeder (editors). 1993. *Mammal Species of the World: a*
- 14 *Taxonomic and Geographic Reference*. Second Edition. Smithsonian Institution Press,
- 15 Washington, DC. xviii + 1206 pp. Available online at: <http://www.nmnh.si.edu/msw/>.
- 16 • Wilson, D. E., and D. M. Reeder (editors). 2005. *Mammal species of the world: a*
- 17 *taxonomic and geographic reference*. Third edition. The Johns Hopkins University Press,
- 18 Baltimore. Two volumes. 2,142 pp.
- 19

Cochise Pincushion Cactus (*Coryphantha robbinsorum*)

The Cochise pincushion cactus was designated as a federally Threatened species on January 9, 1986.

Historic Range: (b) (7)(E) Co., Arizona and Sonora, Mexico. Despite intensive searching, this species is known only from 1 population in southeastern Arizona and 1 in adjacent Sonora, Mexico. Most of the plants are concentrated in small pockets of this tiny range, making the species especially vulnerable to cactus poachers; also potentially threatened by pesticides and mining.
Habitat Comments: Grey limestone hills within a semidesert grassland, with small shrubs, other succulents, and grama grasses. About 1280 m elevation.

Threats: Habitat destruction from grazing, exploration and potential drilling for oil; collection; and off-road vehicles.

Reproduction: Lower reproduction rate than most cacti - estimated average annual production is 3 fruits with 20 seeds per plant.

References:

- Benson, L. 1982. The Cacti of the United States and Canada. Stanford Univ. Press, Stanford, California. 1044 pp.
- Kartesz, J.T. 1994. A synonymized checklist of the vascular flora of the United States, Canada, and Greenland. 2nd edition. 2 vols. Timber Press, Portland, OR.
- Kartesz, J.T., and R. Kartesz. 1980. A synonymized checklist of the vascular flora of the United States, Canada and Greenland. Vol. 2. The biota of North America. Univ. of North Carolina Press, Chapel Hill. 500 pp.
- Lopresti, V. 1984. *Coryphantha robbinsorum* en Mexico. Cactus and Succulent J. Mexico 29: 81.
- Rutman, S. 1992. Handbook of Arizona's endangered, threatened, and candidate plants. U.S. Fish and Wildlife Service. Phoenix, AZ. 30 pp.
- U.S. Fish and Wildlife Service. 1985. Proposal to determine *Coryphantha robbinsorum* (Cochise pincushion cactus) to be a threatened species. Federal Register 50(44): 9083-9086.
- U.S. Fish and Wildlife Service. 1986. Endangered and threatened status for *Coryphantha robbinsorum*. Federal Register 51(6): 952-956.

Lemmon Fleabane (*Erigeron lemmonii*)

The Lemmon fleabane was designated as a Federal Candidate species on December 6, 2007.

Historic Range: Known from a single canyon in the (b) (7)(E), (b) (7)(E) County, Arizona. Despite extensive searching for several years, only this one population is confirmed.

Habitat Comments: Crevices and ledges in limestone canyon walls and on vertical faces of large boulders along canyon bottoms. Surrounding vegetation is pine-oak woodland. 1920-2225 m elevation.

Threats: The greatest threat to *Erigeron lemmonii* is wildfire, which could be intense in the narrow canyon; an intense fire could directly kill individuals, desiccate plants, and alter habitat (Falk 2004). Measures have been taken to reduce the threat of wildfire (Falk 2004). Most plants are on cliff faces well above the heavy fuel loads (Stone 2003). Other potential threats include extended drought, major rock falls, and unauthorized rock climbing (Warren et al. 1991 cited by Stone 2003). Because the species is only known from one population, it is also especially vulnerable to catastrophic events. A management plan for the species has been developed. Rappelling, smoking, and leaving the trail are prohibited in Scheelite Canyon (Stone 2003).

Environmental Specificity: Narrow. Specialist or community with key requirements common.

Environmental Specificity Comments: Known only from crevices and ledges of west, south and north facing cliffs, and on vertical faces of large boulders along a single canyon. It is found on substrates of sandy silicate or granitic soils, and limestone outcrops, between 1920-2225 m (6,300-6,600 ft) elevation.

References:

- Arizona Ecological Services Field Office. 2001. September last update. Lemmon fleabane (*Erigeron lemmonii*) general species information. U.S. Fish and Wildlife Service, Southwest Region 2. Online. Available: <http://www.fws.gov/arizonaes/lemmon.htm> (accessed 19 January 2006).
- Arizona Game and Fish Department. 2004. *Erigeron lemmonii*. Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 5 pp.
- Falk, M. 2004. June-last update. Species Assessment and Listing Priority Assignment Form: *Erigeron lemmonii*. U.S. Fish and Wildlife Service, Endangered Species Program. Online. Available: http://ecos.fws.gov/docs/candforms_pdf/r2/Q0QA_P01.pdf. Accessed 2005, November 30.
- Kartesz, J.T. 1994. A synonymized checklist of the vascular flora of the United States, Canada, and Greenland. 2nd edition. 2 vols. Timber Press, Portland, OR.
- Kearney, T.H., R.H. Peebles, and collaborators. 1951. Arizona flora. 2nd edition with Supplement (1960) by J.T. Howell, E. McClintock, and collaborators. Univ. California Press, Berkeley, CA. 1085 pp.
- Rutman, S. 1992. Handbook of Arizona's endangered, threatened, and candidate plants. U.S. Fish and Wildlife Service, Phoenix, AZ.
- Stone, S. 2003. Management plan template for species at risk on DOD installations: *Erigeron lemmonii*. U.S. Army, Fort Huachuca, AZ.
- U.S. Fish and Wildlife Service. 2002. Candidate and listing priority assignment form: *Erigeron lemmonii*. USFWS, Arizona Ecological Services Field Office, Tucson Sub-office.

- 1 • U.S. Fish and Wildlife Service. 2005. Endangered and Threatened Wildlife and Plants;
2 Review of Native Species that are Candidates or Proposed for Listing as Endangered or
3 Threatened; Annual Notice of Findings on Resubmitted Petitions; Annual Description of
4 Progress on Listing Actions; Proposed Rule. Federal Register May 11, 2005, Vol. 70, No.
5 90: 24870-24934.
6

Huachuca Water Umbel (Cienega False Rush) (*Lilaeopsis schaffneriana* var. *recurva*)

The Huachuca water umbel was designated as a federally Endangered species on January 6, 1997.

Historic Range: The distribution of *Lilaeopsis schaffneriana* var. *recurva* ranges throughout southeastern Arizona and adjacent Sonora, Mexico from Sonoita Creek on the west to Rio San Bernadino on the east. Historically the species reached north to Tucson and south to Cananea, Sonora. The Tucson population along the Santa Cruz River (type locality) no longer exists, presumably due to the loss of perennial flow in this area. The taxon is restricted within this range to small sites with specific wetland requirements. As of 1991, a total of 10 locations in the U.S. and 6 in Sonora, Mexico are known.

Technical Description: An herbaceous semiaquatic perennial with slender erect leaves that grow from the nodes of creeping rhizomes. The rhizomes are usually very shallow, only 1-2 cm underground. They occasionally run along the bottom of still ponds and are generally white. Rhizomes branch freely and may form dense mats in the sand or mud streambed, making it impossible to identify individual plants. The cylindrical hollow leaves, which are typically borne two or three per node, have septa at irregular intervals. The pale green leaves are generally 1-3 mm in diameter, but their length varies depending upon the microhabitat in which they grow. When growing out of the water in wet soil near a stream, the leaves are often only 3-5 cm tall; growing in water that supports their weight, leaves up to 20 cm or more have been observed (Affolter 1985). Three to 10 tiny flowered umbels arise from root nodes. The inflorescence peduncles are typically 1-5 cm tall and always shorter than the leaves. Peduncle length also varies depending upon microhabitat; when growing out of the water they may be only 1-2 cm long, but when under water they may reach 6-7 cm. The flowers are 1-2 mm wide with tiny maroon-tinted petals. The fruits are globose, 1.5-2 mm in diameter, and usually slightly longer than wide.

Diagnostic Characteristics: *Lilaeopsis schaffneriana* var. *recurva* grows in perennial, shallow and slow-moving water. Such sites are rare in southeastern Arizona and northern Sonora, Mexico. *Lilaeopsis* is difficult to locate in the field, in part because it usually occurs with and resembles another small wetland species, *Eleocharis charibea*. *Lilaeopsis* has semisucculent leaves that are somewhat flexuous, whereas *Eleocharis* leaves are pithy, strictly straight and not at all succulent. The leaves of *Lilaeopsis* also appear to be a pale yellow-green compared to the darker green of most co-occurring herbaceous species.

Ecology Comments: Affolter (1985) observed flowering specimens from collections made in June and August and fruiting specimens from May and July through early September. Nature Conservancy botanists have observed *Lilaeopsis* flowering abundantly only once (April 1988), as the local conditions were drying out at Cottonwood Spring along Sonoita Creek. Flowering at low frequency has also been observed from March through October. Affolter (1985) suspects that other members of the genus *Lilaeopsis* self-pollinate. Seed germination from plants grown in an aquarium has been observed. The seeds stuck to the aquarium sides after falling from the parent plants and germinated within 1-2 weeks after ripening (Warren, pers. comm.). Although seeds from *Lilaeopsis schaffneriana* var. *recurva* appear to germinate easily, vegetative reproduction via rhizomatous spreading and dispersal of dislodged clumps is clearly important. Liz Ecker, curator of the Living Collection at the Desert Botanical Garden, has a living specimen which has flowered and born fruit, but she has done no germination studies with the taxon to date (L. Ecker, pers. comm.). An experimental transplant program for *Lilaeopsis* was conducted at the (b) (7)(E) in 1991 in order to establish a second secure population on the refuge that would be less vulnerable to destructive flooding than the existing population or (b) (7)(E). Aside from securing a population, the project allows us to learn more about the ecology and habitat of the species for future management (Warren 1991). Three transplant sites were chosen at perennial ponds. The first transplant took place August 26, 1990; the two subsequent

transplants were made on March 2, 1991, with follow-up monitoring of the transplants done on April 26, 1991. The three transplant sites yielded different results. *Lilaeopsis* could not be relocated at the first transplant site. Competition with other herbaceous plants appeared to have wiped out the transplanted colony. At the second site the *Lilaeopsis* transplant persisted, but due to a moderate amount of surrounding competitive vegetation, the patch did not grow beyond its original 5-inch diameter. However, the third site which was relatively free of competing vegetation, showed tremendous growth and vigor - increasing from 5 inches to approximately 2 feet in diameter over the 1.5-month period. The major conclusion is that *Lilaeopsis* can not survive where there is heavy competition from other herbaceous aquatic plants. Shallow standing water, in contrast to flowing streams, is grown in quickly with aquatic vegetation. Therefore *Lilaeopsis* grown at ponds may need special management to reduce density and accumulated litter from competing vegetation. *Lilaeopsis* is a vulnerable taxon which is easily destroyed by heavy flooding and scouring of habitat, although it also appears to need some amount of disturbance to the habitat in order to decrease surrounding competitive vegetation. *Lilaeopsis* appears to grow year round in the absence of killing frost whereas other aquatic plants tend to die off during the winter, allowing *Lilaeopsis* to more effectively colonize open space following low-level disturbance.

Census data: The Nature Conservancy established and monitored transects at three *Lilaeopsis* locations in 1989. Transects were established at (b) (7)(E) in the (b) (7)(E) and another at (b) (7)(E) near (b) (7)(E). The location (distance along the transect), length, and width of every *Lilaeopsis* patch along permanent transects was recorded. The density of leaves in each patch was also estimated using a rank scale. Lowest density patches received a 0.5 ranking, and highest density patches ranked 3.0. The rank-density value for a sample of patches was correlated with actual stem counts in 12cm x 12cm quadrants to calibrate the scale. Using these counts, a mean density (number of stems per 0.01 square meters) was calculated for each density rank (Gori et al. 1990). Density and coverage of *Lilaeopsis* varies greatly from site to site. Percent coverage of *Lilaeopsis* varied among the sites from 11.5% to 58.3%; of the total area occupied by *Lilaeopsis*, 10.4% to 75.3% had a density value of 2.0 or greater. For specific data see Gori et al. (1990). Together these data provide a profile of the distribution and density of *Lilaeopsis schaffneriana* var. *recurva* along transects in 1989. Similar measurements in subsequent years will indicate what changes have occurred in these streams. The fate of individual patches can also be tracked since the position, length, width and estimated leaf density of every patch is mapped along each transect.

Related species: The genus *Lilaeopsis* contains 13 species of perennial, rhizomatous herbs which live in temperate and alpine regions of North and South America and Australasia. These plants grow in damp, marshy and aquatic habits, often in brackish water. *Lilaeopsis schaffneriana* is one of 4 strictly freshwater species in the genus. It occurs in southeastern Arizona, central and northern Mexico and northwestern South America (Affolter 1985). There is a great deal of morphological variation within *Lilaeopsis schaffneriana*. Some is due to local environmental conditions, as Affolter (1985) showed when he reared plants from the same stock in different depths of water and got great differences in leaf length. Genetic differences, on the other hand, could easily arise among small populations which grow primarily by rhizomatous spreading. Affolter (1985) recognized the Arizona populations as a distinct subspecies based on differences in fruit shape as well as the major geographical gap across the continental divide between the ranges of the Arizonan and Mexican groups. *Lilaeopsis schaffneriana* var. *recurva* inhabits disjunct locations in southeastern Arizona and northern Sonora. Known locations for *Lilaeopsis schaffneriana* var. *schaffneriana* are similarly separated on the central plateau of central and southern Mexico. This kind of distribution is expected for an aquatic species surrounded by arid lands. Affolter expressed suspicion that the discontinuity between the subspecific ranges might reflect a lack of exploration for the plant. However, it is significant that the two subspecies of *Lilaeopsis* are found on opposite sides of the continental divide. So the predominant dispersal mechanism for the species, water, could not serve to mix populations (Warren 1991). *Lilaeopsis masonii* is a candidate category 2 species of northern California. It grows along the margins of rivers, sloughs, and islands of the San Joaquin-Sacramento River

1 delta (California Fish and Game 1988); there are approximately 30 known occurrences.
2 *Lilaeopsis masonii* differs from *L. schaffneriana* var. *recurva* in that it is found in intertidal zones of
3 brackish water marsh. It grows far enough inland so it does not grow directly in salt water as
4 some species of *Lilaeopsis*, but the water is brackish and the plants do experience tidal
5 fluctuation (R. Bittman, pers. comm.). *Lilaeopsis masonii* grows in dense mats at water margins.
6 Associated species are: marsh pennyworts (*Hydrocotyle umbellata* and *H. verticillata*), three-
7 ribbed arrow grass (*Triglochin striata*), mudwort (*Limosella subulata*), tules (*Scirpus* spp.), rushes
8 (*Juncus* spp.), and Suisun marsh aster (*Aster chilensis* var. *lentus*). *Lilaeopsis masonii* grows
9 from an elevation of sea level to 25 feet. It flowers from April to October. Little is known about
10 the ecology/biology of this species (R. Bittman, pers. comm.). The primary threats to *Lilaeopsis*
11 *masonii* are proposed water projects which involve dredging, rip-rapping, levee construction, and
12 other alterations to natural banks and river channels. Heavy cattle grazing also occurs at some of
13 the sites. Petroleum processing plants exist in the area and the species is vulnerable to oil spills.
14 One spill impacted two populations in 1988. The long-term effects of oil on the species is
15 unknown (California Fish and Game 1988). No recovery programs are currently necessary for
16 *Lilaeopsis masonii*, but the species is of interest here because proposed management for the
17 species includes an experimental transplant program. Rip-rap work has been proposed along
18 Barker's Slough in Solano County. This could potentially destroy dense colonies of *Lilaeopsis*
19 *masonii*. A project to transplant all *Lilaeopsis masonii* at the rip-rap sites to suitable habitat has
20 been proposed. Information gained from California's transplant program may prove useful to our
21 efforts at managing *Lilaeopsis schaffneriana* var. *recurva*.

22 **Habitat Comments:** Cienegas (mid-elevation wetland communities), riverine systems, and
23 springs at about 1150-2130 m elevation. Usually in wet soils along the periphery of a channel, in
24 backwaters, or in small openings in the understory near springs. Does not tolerate much
25 competition with other species, but will quickly colonize open habitat created by scouring floods
26 and persist there until interspecific plant competition becomes too great. In order for populations
27 to expand, some plants must remain in areas that escape the effects of periodic scouring floods.
28 *Lilaeopsis schaffneriana* var. *recurva* is restricted to cienega habitats, which are marshy or
29 meadow-like wetlands surrounded by semiarid vegetation (Warren 1991). Hendrickson and
30 Minckley (1984) describe three different types of cienegas based on elevation: low, mid, and high
31 elevation cienegas. Low elevation cienegas or subtropical marshes occur mostly along major
32 perennial rivers below 3000 feet. The low elevation *Lilaeopsis* sites have experienced the most
33 disturbance both human and natural. Low elevation cienega habitats were probably river
34 backwaters and floodplain seeps. These locations are very unstable, experiencing cycles of
35 flooding and drying due to varying climatic patterns. Human influence including groundwater
36 pumping and diversion of water for irrigation have eliminated perennial flow in most southeastern
37 Arizona rivers. Perennial flow is essential for wetland formation. This loss of habitat is evident in
38 the disappearance of 4 historic locations of the taxon. Grazing has added to the problem by
39 contributing to watershed deterioration, which exacerbates erosive flooding and further
40 destabilizes cienega habitats. There are 2 known sites occurring at low elevation on the same
41 stream; one is in the San Bernadino National Wildlife Refuge, in the U.S., and the other in
42 Sonora, Mexico near the border along the Rio San Bernadino. Mid-elevation cienegas occur
43 between 3000-6000 feet. This elevation range fits Hendrickson and Minckley's (1984) definition
44 of true cienega habitat. Permanent water is available and a unique wetland community has
45 developed at these sites (Warren 1991). Flooding potential is lower at these cienega sites
46 because they have smaller drainage areas. Also, the gradients are gentler at these mid-elevation
47 sites as opposed to the higher elevation cienegas. There are 6 current U.S. locations for
48 *Lilaeopsis* at mid-elevation sites, and 4 in Sonora; they are: Bear Canyon, Lone Mountain
49 Canyon, Cottonwood Spring, San Rafael Valley (3 springs) and Turkey Creek in Arizona, and Ojo
50 de Agua de Cananea, Rio San Rafael, Arroyo Los Fresnos and along the Rio Magdalena in
51 Sonora. Flooding, however, is still a potential problem at this elevation range as demonstrated by
52 the population at Cottonwood Spring, which was seriously reduced by flooding from Hog Canyon
53 in 1988. Grazing also has a negative impact on this watershed. High elevation cienegas occur at
54 elevations over 6000 feet. They are described by Hendrickson and Minckley (1984) as "marshy
55 to bog-like alpine and cold temperate meadowland." They may form in surface depressions that

fill with water or at stream headwaters. There are few potential sites for *Lilaeopsis* at these elevations because usually these higher sites are in canyons with stream gradients too steep to support cienega wetlands. Three high elevation sites of *Lilaeopsis* are known in the (b) (7)(E) [REDACTED]. One is in upper Scotia Canyon and another in upper Garden Canyon. An additional *Lilaeopsis* population is reported in Sunnyside Canyon from 6050-6200 feet (S. McLaughlin, pers. comm.). The surrounding vegetation of the cienega communities varies with elevation. Willow (*Salix* spp.) and cottonwood (*Populus* spp.) trees, cattails (*Tyogys* spp.), large reeds, bulrush (*Scirpus* spp.), and halophytes in nearby saline areas are typical of desert-scrub communities of the low elevation cienega sites. Rushes, grasses, fewer cattails, semiaquatic sedges, watercress (*Nasturtium officinale*), water pennywort (*Hydrocotyle americana*), halophytes in adjacent saline areas, and trees (not as common with willows being the most common) are the dominant species of the grassland/oak woodland habitat of mid-elevation cienegas. Finally, the high elevation community is conifer forest including cold-resistant sedges and rushes, semiaquatic and terrestrial grasses, and low, woody alder (*Alnus* spp.) and willow (*Salix* spp.) shrubs. Physical factors, particularly hydrological conditions such as watershed area and stream gradient, appear to limit the distribution of *Lilaeopsis schaffneriana* var. *recurva*. The taxon appears to have specific requirements which limit its distribution to perennial water, gentle stream gradients, small- to medium-sized drainage areas and mild winters. Weather and precipitation data (NOAA 1986) from stations within the range of *Lilaeopsis*: At Canelo 1 NW station in Santa Cruz County, the data are summarized as follows: elevation 5010'; N latitude 31 33'; W longitude 110 32'; mean annual precipitation 17.06"; January mean temperature (F) 42.2; July mean temperature (F) 74.2; and annual mean temperature (F) 57.2.

At (b) (7)(E) FAA station in (b) (7)(E) County, the data are summarized as follows: elevation 4098'; N latitude 31 28"; W longitude 109 36'; mean annual precipitation 12.16"; January mean temperature (F) 44.9; July mean temperature (F) 79.1; and annual mean temperature (F) 61.6. Because the *Lilaeopsis* sites are so dispersed, climatic data provided here are relatively non-specific. Populations inhabit the physiographic province known as the Sonoran Desert Section of Basin and Range. The hydrologic regime appears to be a critical aspect of *Lilaeopsis* habitat. In an effort to characterize hydrologic conditions at each site, we estimated site substrate stability and watershed gradient above the site based on visual observations of the sites. We have made a somewhat arbitrary classification of stream channel conditions at each site as "stable" or "unstable" based on the condition of herbaceous vegetation along the stream bank and channel: stable sites are those where the stream banks, and part or all of the channel, are well stabilized by herbaceous vegetation; unstable sites are those where the channel and much of the banks are unconsolidated, shifting alluvium. Under present watershed conditions, 10 square miles appears to be a watershed size threshold above which flooding is too severe for *Lilaeopsis* to persist, although larger watershed area may be mitigated by low gradient, as at San Bernadino. This taxon does not tolerate much competition with other species, but will quickly colonize habitat disturbed by scouring floods and persist there until interspecific plant competition becomes too great. In order for populations to expand, some plants must remain in areas that escape the effects of periodic floods (Rutman and Rorabaugh 1995).

Stewardship Overview: High priority needs include protecting perennial stream flow through acquisition of water rights; management of the watershed to assume a good vegetative cover by perennial grasses to prevent scouring floods and monitoring known populations to detect downward trends if they occur. Working with private landowners is a high priority since several sites are on private land.

Restoration Potential: At present there is not enough evidence of a decline in the populations to require a recovery program. However, it is important to maintain existing populations at their present levels to guard against any possible future decline. The species shows evidence of successful reproduction at all known sites indicating a high recovery potential. The experimental transplant program at San Bernadino shows a high survivorship rate given suitable growing conditions (ie. few surrounding competitive species).

Preserve Selection & Design Considerations: Adequate protection of *Lilaeopsis* populations requires consideration of the direct site impacts as well as indirect effects of water supply and watershed condition. Therefore, primary site boundaries may be relatively small and include only the wetland habitat where *Lilaeopsis* is found. Secondary site boundaries should include key portions of the watershed to be managed for maintenance of water supply and erosion control. An important protection consideration is the acquisition of water rights to ensure stable future water levels. Various privately owned sites should be protected through continuing land owner education and assistance. These sites should be put in protective ownership if the opportunity presents itself.

Management Requirements: The primary management need of *Lilaeopsis schaffneriana* var. *recurva* is to protect the cienega habitat that supports known populations. Management procedures include protecting water supplies by acquiring instream flow water rights and managing watersheds to reduce flood frequency and intensity. Continued monitoring of the known populations and surveys for other potential locations should also be part of the management procedure. Recreation management may be necessary at some local populations. Prescribed burns may be essential for certain populations to reduce the density of accumulated litter from competing vegetation.

Monitoring Requirements: Continued monitoring every other year of existing populations is needed in order to determine whether the populations are stable, increasing or declining and subject to nearby threats. Three of the 12 known sites have been monitored by The Nature Conservancy since 1989. The percent coverage and density of the species were determined along transects (Gori et al 1990).

Management Programs: This element is not being actively managed.

Monitoring Programs: One program underway since 1989. Contact: Peter Warren, Public Lands Protection Planner, The Nature Conservancy, Arizona Field Office, Tucson, Az.

Management Research Programs: One research program involving transplant populations was conducted by The Nature Conservancy in 1991 and is being monitored at the San Bernadino National Wildlife Refuge.

References:

- Affolter, J. M. 1985. A monograph of the genus *Lilaeopsis* (Umbelliferae). Systematic Botany Monographs. Vol. 6. American Society of Plant Taxonomists. 140 pp.
- Affolter, J.M. 1985. A monograph of the genus *Lilaeopsis* (Umbelliferae). Systematic Botany Monographs 6: 1-140.
- California Department of Fish and Game Endangered Plant Project and the Natural Diversity Data Base. 1988. California Native Plant Status Report: *Lilaeopsis masonii*. California Dept. of Fish and Game, Sacramento, CA.
- Correll, D.S., and H.B. Correll. 1972. Aquatic and wetland plants of southwestern United States. 2 volumes. Stanford Univ. Press, Stanford, CA. 1777 pp.
- Gori, D.F., P.L. Warren and L.S. Anderson. 1990. Population studies of sensitive plants of the Huachuca, Patagonia and Atacosa Mountains, Arizona. Submitted to Coronado National Forest in completion of P. O. 40-8197-9-0119. The Arizona Nature Conservancy, Tucson, AZ.
- Hendrickson, D. A. and W.L. Minckley. 1984. Cienegas: Vanishing climax communities of the American Southwest. Desert Plants 6(3): 1-175.
- Hill, A. W. 1927. The genus *LILAEOPSIS*: a study in geographical distribution. Journal of the Linnaeus Society of Botany 47:525-551.

- 1 • Kartesz, J.T. 1994. A synonymized checklist of the vascular flora of the United States,
2 Canada, and Greenland. 2nd edition. 2 vols. Timber Press, Portland, OR.
- 3 • Kartesz, J.T. 1999. A synonymized checklist and atlas with biological attributes for the
4 vascular flora of the United States, Canada, and Greenland. First edition. In: Kartesz,
5 J.T., and C.A. Meacham. Synthesis of the North American Flora, Version 1.0. North
6 Carolina Botanical Garden, Chapel Hill, NC.
- 7 • Kartesz, John T. 1991. Synonym names from 1991 checklist, as extracted by Larry
8 Morse, TNC, June 1991.
- 9 • Kearney, T.H., R.H. Peebles, and collaborators. 1951. Arizona flora. 2nd edition with
10 Supplement (1960) by J.T. Howell, E. McClintock, and collaborators. Univ. California
11 Press, Berkeley, CA. 1085 pp.
- 12 • NOAA. 1986. Climatological data annual summary: Arizona 90 (13). U.S. Department of
13 Commerce, National Environmental Satellite, Data and Information Service, National
14 Climatic Data Center, Asheville, NC.
- 15 • Rutman, S. 1992. Handbook of Arizona's endangered, threatened, and candidate plants.
16 U.S. Fish and Wildlife Service, Phoenix, AZ.
- 17 • Rutman, S., and J. Rorabaugh. 1995. Endangered and threatened wildlife and plants:
18 Proposal to determine endangered status for three wetland species found in southern
19 Arizona and northern Sonora. Federal Register 60(63): 16836-16846.
- 20 • Shreve, F., and I.L. Wiggins. 1964. Vegetation and flora of the Sonoran Desert. 2
21 volumes. Stanford Univ. Press, Stanford. 1740 pp.
- 22 • U.S. Fish and Wildlife Service. 1993. Endangered and threatened wildlife and plants;
23 review of plant taxa for listing as endangered or threatened species. Federal Register, 50
24 CFR Part 17, 58(188):51144-51190.
- 25 • Warren, P. L., D.F. Gori, L.S. Anderson and B.S. Gebow. 1991. Status report: *Lilaeposis*
26 *schaffneriana* var. *recurva*. Submitted to : U.S. Fish and Wildlife Service. The Arizona
27 Nature Conservancy, Tucson, AZ.
- 28 • Warren, P.L. 1991. Reconvry of *Lilaeposis schaffneriana* var. *recurva* at San Bernadino
29 National Wildlife Refuge. Submitted to : U.S. Fish and Wildlife Service. The Arizona
30 Nature Conservancy, Tucson, AZ.
- 31 • Warren, P.L., L.S. Anderson and P.B. Shafroth. 1989. Population studies of sensitive
32 plants of the Huachuca and Patagonia Mountains, Arizona. Submitted to Coronado
33 National Forest in completion of P.O. No. 40-8197-9-431. The Arizona Nature
34 Conservancy, Tucson, AZ.

1 **Madrean (Canelo Hills) Ladies Tresses (*Spiranthes delitescens*)**

2 The Madrean ladies tresses was designated as a federally Endangered species on January 6,
3 1997.

4 **Historic Range:** Four populations of *Spiranthes delitescens* have been found in Arizona:
5 (b) (7)(E) County, above the dam in Babocomari Cienega; Santa Cruz County, along Turkey
6 Creek, at O'Donnell Cienega, and on a slope below Sheehy Spring (Sheviak 1990). This species
7 most likely exists in Mexico; however, to date, no plants have been located south of Arizona
8 (Sheviak 1990). The occurrence of other populations of this species of Ladies' Tresses in the
9 United States is probably unlikely due to the limited available sites possessing the specific habitat
10 parameters which appear to be required by these plants (Sheviak 1990).

11 **Diagnostic Characteristics:** *Spiranthes delitescens* can be distinguished from other Mexican
12 and southwestern United States *Spiranthes* species by the shape of its medium-sized flowers: the
13 floral tube curving into a horizontal apex and an ascending base, and the sepals curving outward
14 and downward. In addition, the pubescence is distinct; the trichomes are glandular-capitate and
15 taper at the apex. Cytological differences between *S. delitescens* and other *Spiranthes* species
16 also exist (Sheviak 1990).

17 **Reproduction Comments:** Flowers, Pollination and Hybridization: Orchid flowers have a unique
18 morphology which has coevolved with their pollinators (van der Pijl and Dodson 1966). A large
19 petal, called the labellum or lip, acts as a landing platform for many pollinators. In *Spiranthes*
20 spp., balls of sticky pollen grains, pollinia, are positioned near the column (the partially united
21 stamen and pistil) in such a way that when the pollinator enters the floral tube, on its way to the
22 nectaries, it inadvertently triggers the rostellum causing the pollinia to be deposited on the
23 pollinator (van der Pijl and Dodson 1966). Bees are the primary pollinator for *Spiranthes*, with
24 *Bombus* being the most common genus; other pollinating organisms include flies, moths, and
25 butterflies (van der Pijl and Dodson 1966; Dressler 1981). Within three days of successful
26 pollination, *Spiranthes* flowers dehydrate and become discolored (Catling 1982). One pollinia
27 contains over 10,000 pollen grains. This allows for efficient fertilization of the thousands of ovules
28 in the ovaries of most orchids (van der Pijl and Dodson 1966). Some orchids are self-fertile, but
29 most often fertilization is the result of outcrossing. Self-pollination is advantageous when plants
30 have extended their range into areas not previously inhabited by the species (Dressler 1981).
31 *Spiranthes* are often self-fertilized, and individuals that require cross-pollination are receptive for
32 only 10 to 40 days (Catling 1982). Flowers older than 40 days contain dead ovules (Catling
33 1982). Within three weeks of pollination the seeds are fully developed and the ovary splits.
34 Usually 100% of *Spiranthes* ovaries expand, but often only 50% of them contain seeds (Catling
35 1982). Orchids easily hybridize; both inter-specific and inter-generic hybrids occur in the wild
36 (Sanford 1974). In dry climates, flowering often occurs during the rainy season. Flowering of the
37 *Spiranthes* occurs in July, when temperatures range from 60°F at night to 100°F during the day
38 and when the majority of the year's 15 to 20 inches of precipitation falls (Merrigan 1990; The
39 Nature Conservancy Arizona Field Office, pers. comm.). In some cases too much rain, possibly
40 causing a decrease in pollinator activity, results in a decrease in the number of flowers and
41 consequently the number of fruits (Dressler 1981). Most nontropical species release their seeds
42 in the fall at the beginning of the dormant period (Dressler 1981). In addition to moisture
43 dependency, flowering of some species of *Spiranthes* is photoperiodically induced (Catling 1982).
44 The age of sexual maturity is dependent on the species and can range from several years to over
45 twenty years (Stoutamire 1974; B. Jennings, pers. comm. 25 Jan. 1990). Inflorescences first
46 develop in *Spiranthes spiralis* thirteen to fifteen years after seed germination (Wells 1981). Once
47 reproductively mature, the age of the plant is not a factor in flowering, whereas temperature and
48 precipitation appear to be significantly related to the percentage of flowering plants (Wells 1981).
49 *Spiranthes diluvialis* will not bloom in dry years when precipitation levels are atypically low (B.
50 Jennings, pers. comm. 25 Jan. 1990). *S. spiralis* plants which have reverted back to the
51 saprophytic stage are capable of flowering during the initial year of resuming above-ground
52 growth (Wells 1967). The average percentage of flowering *S. spiralis* plants over a thirteen year

1 period was 33, ranging from 73% in 1966, 19% in 1970, 43% in 1973, down to 1% after the 1976
2 drought, and recovering to 31% the following year (Wells 1981).

3 **Ecology Comments:** *Spiranthes* of Canelo Hills Cienega and Turkey Creek: The grass-like
4 leaves of the orchid, growing low in the sedge and horsetail fields, are difficult to see for most of
5 the year. The inconspicuous plants are visible July and August when the roughly 20 cm tall
6 inflorescences develop (P. Sundt, pers. comm. 23 Jan. 1990; The Nature Conservancy Arizona
7 Field Office, pers. comm.). The fruits mature approximately three weeks after the flowers form,
8 usually during the end of August, releasing hundreds of tiny seeds from each capsule to be
9 dispersed, probably via the wind (McClaran and Sundt 1992; P. Sundt, pers. comm. 23 Jan.
10 1990). Many inflorescences are damaged during the summer; Sundt (pers. comm. 23 Jan. 1990)
11 feels that grasshoppers may be responsible for the broken stalks and the chewed capsules. The
12 life-cycle of the plants is unclear. Most likely these orchids are perennial; however, no dormant
13 underground structures have been identified (McClaran and Sundt 1992). Determining the over-
14 wintering structure is difficult without disturbing the plants. The plant may remain below-ground
15 most of the year or, common to many *Spiranthes*, small, inconspicuous leaf rosettes may grow
16 throughout the cool months, hidden by the tall vegetation (McClaran and Sundt 1992; B.
17 Jennings, pers. comm. 25 Jan. 1990). In February, an inspection of approximately 40 flagged
18 areas (presumably indicating the previous year's orchids) revealed no above-ground orchid
19 structures (Newman 1990). Plants rarely flower in consecutive years and the relationship
20 between the flowering plants cannot be elucidated since the growth pattern of the subterranean
21 structures is unknown (McClaran and Sundt 1992). Censusing of the *Spiranthes* at Canelo Hills
22 began in 1978; however, accurate assessment of the demographic patterns is difficult because
23 varying techniques were used during the first eight years of monitoring. With this caveat in mind,
24 the total number of plants in O'Donnell Canyon fluctuated from 40 in 1978, 196 in 1979, dropping
25 to 30 in 1982 through 1984, and then increasing to roughly 80 plants in 1988 (McClaran and
26 Sundt 1992). These data suggests that the number of flowering plants has declined since 1979.
27 Few conclusions can be drawn from the data, considering that the early measurements were
28 based on the number of flowering plants and the later censusing was based on the total number
29 of plants (flowering and not flowering), and that individual plants would appear one year, not
30 appear the following year (no visible above-ground structures), and then reappear in subsequent
31 years. In fact, it is difficult to estimate population size based on counts of aboveground plants
32 due to the lack of information concerning the life-cycle and environmental requirements of
33 *Spiranthes delitescens*. Other species of *Spiranthes* grow initially underground saprophytically
34 for many years, revert back to saprophytic growth when environmental conditions are not
35 favorable and flower irregularly. Population declines followed by recoveries are characteristic of
36 many *Spiranthes*. The plants growing at Turkey Creek appear to be in a plant community
37 characterized by shorter plant height and greater alpha diversity than at Canelo Cienega.
38 Grazers have been excluded from the latter location since 1969, when this part of Canelo Hills
39 was bought by The Nature Conservancy; Turkey Creek is currently grazed (McClaran and Sundt
40 1992; P. Sundt, pers. comm. 23 Jan. 1990). Thus, although soils and topography of the two sites
41 differ, grazing is also a likely factor differentiating the two sites. The population in Turkey Creek,
42 ranging from hundreds to thousands of plants, appears healthier and more vigorous than the
43 Canelo Hills' population (McClaran and Sundt 1992; M. Heitlinger, pers. comm. 8 Jan. 1990; P.
44 Warren, pers. comm. 25 Jan. 1990). Sundt (pers. comm. 23 Jan. 1990) proposes that the Turkey
45 Creek plants have always been more vigorous than the O'Donnell Creek plants, due to the
46 different characteristics of the particular sites, and that little significant change has occurred in the
47 two populations over time.

48 **Spiranthes and Other Terrestrial Orchids**

49 **Seeds and Fruits:** Terrestrial orchid fruit are usually thin-walled, dry, and papery (Dressler
50 1981). Depending upon the species, *Spiranthes* fruit may mature within a few days after
51 fertilization or may take as long as one year to completely develop (Luer 1975). Seeds of
52 terrestrial orchids tend to mature and are dispersed at the end of the plants' growing season,
53 which often coincides with the time of maximum germination (Stoutamire 1974). When fully

1 mature, the valves on the capsule open and the wind-borne seeds are dispersed (Luer 1975).
2 Water and humans have also been implicated in orchid seed dispersal; there is no evidence
3 supporting the involvement of non-human animals (Sanford 1974). Orchid seeds have been
4 found 400 miles from the parent plant; without human intervention, however, dispersal rarely
5 occurs this far (Sanford 1974). Orchid seeds are rudimentary when dispersed; the sole protection
6 of the undifferentiated embryo is the seed coat, and no endosperm or other form of nourishment
7 surrounds the embryonic plant (Luer 1975; B. Jennings, pers. comm. 25 Jan. 1990). Due to the
8 naked, unprotected seed structure a dormancy period is highly unlikely and the period of viability
9 relatively short (Stoutamire 1974; B. Jennings, pers. comm. 25 Jan. 1990). The rapid dispersal,
10 lack of dormancy, requirement for specific fungi, and necessity of precise environmental
11 conditions explains the extremely low seed survival rate of an estimated one in a million
12 (Stoutamire 1974; Luer 1975; B. Jennings, pers. comm. 25 Jan. 1990). Cultivated Orchids:
13 Terrestrial orchids are difficult to grow due to the specific symbiotic associations often required.
14 Dimmitt (pers. comm. 22 Jan. 1990) does not know of any amateur orchidist having successfully
15 germinated and cultivated any member of the genus *Spiranthes*. Although limited, laboratory and
16 greenhouse experiments have uncovered some information on the germination and growth of
17 terrestrial orchids. The seeds of many *Spiranthes* species retain their viability for three years
18 when stored in a refrigerator (Stoutamire 1974). *Spiranthes cernua* seeds germinate readily in
19 sterile water; *S. orchioides* seeds swell with imbibition but fail to germinate (Stoutamire 1974).
20 When placed under a light source after germination, several *Spiranthes* species produce
21 chlorophyll; this indicates an ability to grow autotrophically in the absence of a mycorrhizal
22 associate. However, other species require sterile agar media, containing mineral salts and an
23 external source of organic carbon, indicating an obligate heterotrophic (required mycorrhizal
24 associates) stage (Stoutamire 1974; Dressler 1981; Arditti 1982). Arditti (1982) lists the specific
25 media requirements for laboratory growth of many *Spiranthes* species. No information is
26 available about the early growth requirements of *S. porrifolia* and *S. vernalis*, the putative parents
27 of the southern Arizona plants. When plants are grown in sterile laboratory conditions, light is
28 required for normal development of many early photosynthesizing species, but it may inhibit the
29 germination of the late-photosynthesizing species (Stoutamire 1974). A protocorm develops from
30 the undifferentiated embryo and is the initial external structure when seed germination
31 commences (Sanford 1974; Stoutamire 1974). Two stages of high mortality are found in agar-
32 grown seedlings: the first stage occurs shortly after the protocorm emerges from the seed coat,
33 when it reaches 1 mm to 2 mm in length, and the second stage occurs shortly after the roots
34 develop. In the wild this later stage correlates with the transitional period when the seedling
35 changes from an obligate mycorrhizal dependent to a partly autotrophic organism (Stoutamire
36 1974). In the lab, seedling growth initially occurs in the downward direction and after several
37 centimeters of growth the apical meristem turns and grows upward (Stoutamire 1974); in
38 *Spiranthes* the protocorm initially forms into the tubercle (Sanford 1974). During the first year of
39 growth, short thickened corms or modified lateral buds, called sinkers, are formed in most
40 terrestrial orchids (Stoutamire 1974). *Spiranthes spiralis* development is expedited by laboratory
41 conditions and within 18 months after the seeds are sown, four green leaves and a 5 mm long
42 tuber are produced (Wells 1981). Enlarged primary structures develop concurrently with the first
43 seedling leaves. Adventitious buds on the stem of some *Spiranthes* species are capable of
44 vegetative reproduction (Stoutamire 1974). In the greenhouse, *S. cernua* and *S. sinensis*
45 develop from a protocorm to a flowering plant in 35 months and 29 months, respectively
46 (Stoutamire 1974).

47 **Germination and Mycorrhizal Associations:** Mycorrhizal penetration into the seed and embryo
48 is required for successful germination of most terrestrial orchid seeds; the seedlings are obligate
49 mycorrhizal dependents until aerial shoots and photosynthesizing apparatuses have developed
50 (Dressler 1981). The abundance of hair-like projections on the non-photosynthesizing
51 protocorms may allow for rapid mycorrhizal association (Stoutamire 1974). Results from
52 laboratory studies suggest a more rapid germination and development period in the early
53 photosynthesizing species than in the late photosynthesizing species, possibly due to a
54 facultative, rather than obligate, relationship of the former species with the fungus (Stoutamire
55 1974). Most often chlorophyll does not develop for several months even in the early

1 photosynthesizing species (Dressler 1981). Wells' (1981) results indicate that juvenile orchids
2 remain underground and thus without chlorophyll for greater than one year and maybe as long as
3 fifteen years. As the plant ages, the dependency on fungi is reduced; however, most mature
4 terrestrial orchid roots are associated with endophytic fungi (Warcup 1975, Dressler 1981). Most
5 of the rapidly photosynthesizing protocorm species require sunlight to germinate and often grow
6 in sunny wet areas, characteristic of open marshes and bogs (Stoutamire 1974; Dressler 1981).
7 Whereas germination of most of the non-photosynthesizing protocorm species is inhibited by
8 light, these species grow in well-drained forest soils or open, seasonally dry grasslands
9 (Stoutamire 1974; Dressler 1981). Thus species that grow in cienegas, such as the southern
10 Arizona plants, are presumably early photosynthesizers. Mycorrhizal fungi are required to supply
11 the embryo with needed enzymes and nutrients early in the growth of the seedling; minerals,
12 vitamins and an available organic carbon source are essential to the development of the plant
13 (Stoutamire 1974; Luer 1975; Dressler 1981). The species-specificity of the fungi-orchid
14 symbiosis is ambiguous and is thought to decrease as the plant ages (Warcup 1975). Several
15 different species of fungi are associated with most roots, and taxonomic relationships between
16 fungi and orchid species appear to exist (Warcup 1975; Dressler 1981). Environmental
17 conditions will affect the fungi-orchid relationship; high levels of nitrogen and low soil pH may
18 reduce the likelihood of fungal penetration into the seed, thus decreasing the germination rate
19 (Warcup 1975). The absence of visible growth of an orchid plant does not imply dormancy or
20 death of the plant (Stoutamire 1974). Often orchids grow below-ground for several years without
21 emerging from the soil, receiving nourishment from fungal assimilates (Stoutamire 1974). Some
22 terrestrial orchids have grown saprophytically and remained underground for fifteen years
23 (Sanford 1974). *Spiranthes spiralis* grows saprophytically, solely as a mycorrhizal-rhizome type
24 structure, for eight years before a tuber is produced and a total of eleven years before aerial
25 stems are produced (Wells 1981).

26 **Vegetative Growth and Population Fluctuations:** Orchids may grow vegetatively for many
27 years before flowering. *Cypripedium candidum* requires more than twelve years to reach
28 reproductive maturity (Bender 1986) and some *Spiranthes* only bloom every twenty years (B.
29 Jennings, pers. comm. 25 Jan. 1990). Underground structures include tubers, corms, sinkers,
30 roots, and storage roots (Stoutamire 1974). Vegetative propagation occurs through the growth of
31 buds on lateral underground stems, and newly formed plants eventually separate from the parent
32 plant (Wells 1967). Orchids do not produce typical primary roots and most growth occurs in the
33 secondary root system (Stoutamire 1974). The roots of most terrestrial orchids which grow in
34 moist areas occur above the water-line, allowing for the provision of sufficient amounts of oxygen
35 (Dressler 1981). Depending on the species, above-ground vegetative growth may continue year-
36 round or only during the warm growing season. The normally slow growth rate often decreases in
37 the cool season and small over-wintering leaf rosettes may form (B. Jennings, pers. comm. 25
38 Jan. 1990). *Spiranthes spiralis*, growing in the grasslands of England, are green year-round; leaf
39 rosettes are present when the plants are not in bloom (Wells 1981). In January, a mature plant
40 will contain two mature tubers produced the previous year and a small protuberance, which will
41 develop into the following year's tuber. Plants of this species produce no roots, thus the tuber
42 and fungi are responsible for obtaining the necessary nutrients and water. In July the leaf
43 rosettes die and by August new leaves are formed and a flowering stalk develops (Wells 1981).
44 Stable communities, with a relatively fixed number of mature plants, often have high seedling
45 mortality (Stoutamire 1974). However, terrestrial orchid populations often display great
46 fluctuation within several year periods (Luer 1975). Colonies of many *Spiranthes* species are
47 often labile and above-ground parts may appear and disappear in alternating years (Luer 1975).
48 Population size can alternate from several to hundreds to thousands and back down to several
49 plants in a few years (B. Jennings, pers. comm. 25 Jan. 1990). Plants of *Spiranthes diluvialis* in
50 one location fluctuated from 5500 visible flowering plants in 1986 to 200 plants in 1989, whereas
51 another population, experiencing similar weather conditions and apparently no different
52 management practices, did not have a large flux in population size (B. Jennings, pers. comm. 25
53 Jan. 1990). One population of *Spiranthes spiralis* went from 420 plants in 1963 to 1050 plants in
54 1969 (Wells 1981); however, the population size of *Spiranthes spiralis* usually remains relatively
55 constant (Wells 1967). Sheviak (1974) attributes the pronounced changes in population size and

1 distribution to both climatic fluctuations and edaphic factors which influence the
2 saprophytic/autotrophic state of the orchid. Due to the narrow pH tolerance, specific temperature
3 and moisture requirements of the fungi-orchid association, changes in the environment will lead
4 to altered states of the orchid (Sheviak 1974, B. Jennings, pers. comm. 25 Jan. 1990). In
5 horticultural conditions, *S. cernua* and *S. magnicamporum* can revert from an autotrophic state to
6 a saprophytic state (Sheviak 1974). Some orchids, such as *Triphora trianthophora*, grow
7 underground saprophytically for most of their life and only occasionally produce aerial stems
8 (Sheviak 1974). *Habenaria leucophaea* and some other very rare orchids, may produce
9 hundreds of plants in a location where it was previously rare and then one or two seasons later
10 disappear back to the saprophytic state where it remains for many years (Sheviak 1974). The
11 various negative slopes in the linear survivorship curves (number of plants versus survival years)
12 of different *Spiranthes spiralis* cohorts (same age plants) indicate that the chance of survival is
13 dependent on the year in which the cohorts were produced and is not significantly affected by
14 varying environmental conditions (Wells 1981). The mean expected life for all the cohorts was 53
15 years and the calculated time until only one plant remained for each cohort ranged from 23 years
16 to 67 years (Wells 1981).

17 **Habitat Comments:** Cienegas (mid-elevation wetland communities) at about 1525 m elevation.
18 Soils are highly organic and seasonally or continuously water-saturated, but are not subject to
19 scouring floods. Associated plants are mostly tall grasses, sedges, and rushes. Most members
20 of the genus *Spiranthes* require a moist habitat (B. Jennings, pers. comm. 25 Jan. 1990). *S.*
21 *graminea* grows abundantly in cienegas (permanently wet meadows in desert foothills) and in the
22 mountains of central Mexico (Luer 1975). *Spiranthes diluvialis*, in Colorado and Utah, grow in
23 flood plains, old stream channels and along streambeds, in densely vegetated open sites and
24 under willow trees (B. Jennings, pers. comm. 25 Jan. 1990). The four populations of *Spiranthes*
25 *delitescens* occur above a dam in Babocomari Cienega, in marshy meadows, seeps and
26 hummocks along Turkey Creek, in marshy meadows and seeps at O'Donnell Cienega, and on a
27 seeping slope below Sheehy Spring (Sheviak 1990). The dominant vegetation in the cienegas
28 near the *Spiranthes* include grasses, sedges (*Carex* spp.), rushes (*Juncus* spp.), spike-rush
29 (*Eleocharis* spp.), cat-tails (*Typha* spp.), and horsetails (*Equisetum* spp.) (Grater 1973, Merrigan
30 1990, The Nature Conservancy Arizona Field Office, pers. comm.). Johnson grass (*Sorghum*
31 *halepense*), a potential threat to the orchid, appears to be spreading into the marshy meadow
32 (McClaran and Sundt 1992). The cienegas are at approximately 1500 m elevation and contain
33 fine grained, highly organic, saturated soils (Merrigan 1990, The Nature Conservancy Arizona
34 Field Office, pers. comm.). The orchid grows in both saturated soil and the surrounding drier
35 sites.

36 **Stewardship Overview:** *Spiranthes delitescens* is a newly identified species known from four
37 populations in southern Arizona; it is distinct from *Spiranthes graminea*. The population at one
38 site, ranging from hundreds to thousands of plants, appears healthier and more vigorous than
39 another one. Possibly the greatest threat to the survivability and fecundity of the orchid is the
40 dense vegetation surrounding the small orchid plants. Possible effective management practices
41 such as grazing, fires, and control of competing native and non-native plants have not been
42 researched enough to determine the best practice or the best combination of practices. Due to
43 the possible fluctuation in population size resulting from the reversion from a partially autotrophic
44 plant back to a saprophytic plant, a characteristic common to many *Spiranthes* species, the status
45 of these plants cannot be determined. Extensive research on the life-cycle and environmental
46 requirements of this species is required before management plans should be discussed; burning
47 experiments are being planned for one population.

48 **Restoration Potential:** Recovery of the *Spiranthes* is dependent on determining the optimum
49 habitat conditions required for successful flowering, fruiting, germination, and maturation. Most
50 probably this will relate to reduction in the density of the vegetation cover of the marsh. A
51 prescribed burn at one protected site in 1991 failed to increase orchid numbers that year. But
52 because saprophytic individuals in other *Spiranthes* species take at least one year to revert to
53 aboveground plants and because germinated seeds must spend one to twelve years as obligate

saprophytes, the response of the population to the prescribed burn is not known at this time. Therefore, until the ecological requirements are known and optimal conditions can be produced through management actions, we can only speculate as to the recovery potential of the populations.

Management Requirements: Discussions on Natural Occurrence and Management Implications of Fire and Grazing at Two Sites and the Response of Orchids to Habitat Alterations: Heitlinger (pers. comm. 8 Jan. 1990) and McClaran (pers. comm. 24 Jan. 1990) feel that historically fires occurred naturally in the cienegas when lightning-caused fires in the uplands spread down into the marshes and burned at cool temperatures. Suppression activities and roads are factors resulting in the reduction of the spread of natural fires. Most likely the fires would have occurred in the late spring (April through June) before moist, green vegetation developed (Merrigan 1990). In this case, fires would have periodically removed the dense vegetation surrounding the orchids prior to maximum orchid growth. However, Gehlbach (1986) and Sundt (pers. comm. 23 Jan. 1990) feel that little evidence exists to support the assumption that fires frequently swept through the marshes; they believe the wet marsh would not support fires. Perhaps fire was restricted to drought years or occurrences of winter lightning storms. The possibility of burning having a detrimental effect on the orchid does exist if the fire occurs during a crucial growth phase or if the fleshy surface tubers are damaged by fire (P. Sundt, pers. comm. 23 Jan. 1990; The Nature Conservancy Arizona Field Office, pers. comm.). Controlled burning maintains the appropriate habitat for some orchid species (Dressler 1981). Some species in South Africa and Australia flower only after fires, some flower more prolifically without fire, and the flowering of some other species is unaffected by fire (Dressler 1981). Several prairie orchids, such as *Spiranthes cernua*, sand-prairie ecotype, and *Spiranthes lacera*, appear to increase the number of flowering plants after burns (conditions of the burns were not indicated); possibly, fire physiologically triggers the bloom stage (Sheviak 1974; Sanford 1974; B. Jennings, pers. comm. 25 Jan. 1990). Orchids with protected underground buds tend to benefit (increase in number of flowering plants) or be unaffected by fires, whereas species with surface pseudobulbs require protective rocky spots in order to survive fires (Sanford 1974). Most likely, the timing of a fire is extremely critical. A burn at one site conducted in April 1979 resulted in the increase in orchid number from 40 to 196 in August following the fire (McClaran and Sundt 1992; The Nature Conservancy Arizona Field Office, pers. comm.). However, the number of plants growing in unburned locations also increased during this period, so possibly other environmental conditions were responsible for the significant increase in number of orchid plants (McClaran and Sundt 1992; The Nature Conservancy Arizona Field Office, pers. comm.). A fire conducted in May 1986 resulted in a decrease in population size from 97 (flags, presumably indicating orchids from the previous year) to 8 plants (McClaran and Sundt 1992; The Nature Conservancy Arizona Field Office, pers. comm.). The difference in the effect of the second fire compared to the 1979 fire may be due to the more advanced, vulnerable growth stage of the orchid in May. These results indicate the importance of determining the most beneficial time of burning. The effects of high fuel loads and temperature of burns in the cienega should be determined in order to prevent damage to the tuber by hot fires. Gehlbach (1986) emphasizes the importance of grazing on marsh vegetation. Over the past 10,000 years periodic exposure of southern Arizona cienegas to mammoths, Spanish cattle and Anglo livestock have resulted in trampling and grazing. Gehlbach (pers. comm. 25 Feb. 1990) feels that short durations of heavy grazing, analogous to the conditions of migratory animals, may be a natural and efficient means of managing the cienega. Livestock possibly aids in the survival of the orchid by tilling the soil, providing appropriate microsites for seedling establishment, and decreasing the litter accumulation. McClaran and Sundt (1992) suggest that grazing at one site and the exclusion of grazing at another site may explain the more abundant orchid plants at the former location. *Spiranthes* at the first site grow in a more open and less crowded vegetative (not necessarily more natural) setting than those at the second site (P. Sundt, pers. comm. 23 Jan. 1990). Possibly, cattle grazing may aid in the orchid growth by reducing the competition of neighboring grasses for space and nutrients (Fernald 1987). However, the populations at both of these sites are both described as decreasing in the number of flowering plants over the past ten years (M. McClaran, pers. comm. 24 Jan. 1990), thus damaging the argument of the effectiveness of grazing. Due to the absence of grazers for

1 thousands of years, between the period of mammoths and cattle, Heitlinger (pers. comm. 8 Jan.
2 1990) feels that a non-grazing disturbance was most likely associated with the recent evolution of
3 this orchid. Management experiments on *Spiranthes spiralis* indicate that grazing by rabbits
4 cleared the vegetation and provided sites for seed germination eleven years prior to the study.
5 This is evident by the increase in number of autotrophic seedlings of a species that requires
6 eleven years of saprophytic development prior to emergence (Wells 1981). This experiment
7 suggests the long-term time span required to assess the response of a *Spiranthes* species to a
8 particular management technique. The rare orchid *Spiranthes magnicamporum* increases
9 significantly in lightly grazed areas, but apparently the benefit from grazing is not due to increases
10 in light level; the optimum grazing level is so low that there is no significant reduction in
11 vegetation (Sheviak 1974). Casual observations indicate a high concentration of several
12 *Spiranthes* species in grazed areas. The rare *S. parksii*, which grows in open, grassy woodland
13 sites in Texas, is most abundant in areas exposed to heavy cattle grazing; the *S. romanzoffiana*
14 growing in Alaska is especially abundant along moose trails; Gehlbach (pers. comm. 25 Feb.
15 1990) suggests the possibility of the hoof-turned soil benefitting the establishment and/or survival
16 of the plants. Higher concentrations of *S. cernua* and *S. gracilis* are found growing beside horse
17 trails than in areas distant from horse trails; the plants occur close to the trail where the effects of
18 the hooves are present, but far enough from the trail to be out of reach of the grazers (F.
19 Gehlbach, pers. comm. 25 Feb. 1990). Detrimental effects of grazing are illustrated by the
20 apparent (but not confirmed) extirpation of a population of *Spiranthes diluvialis* plants in Utah in a
21 heavily grazed field (Sheviak 1984). The species may have a number of additional management
22 needs although the research needed to identify these needs has not been completed. These needs
23 include: (1) maintenance of the hydrologic regime; (2) control of exotics like Johnson grass; and
24 (3) reduction of accumulated litter to increase light and water availability to orchids. Maintenance
25 of the hydrologic regime may require the retirement or reduction of grazing in the watershed to (i)
26 stabilize spring flows and (ii) reduce the probability of a scouring flood and channel erosion, thus
27 ensuring that water table depths remain near the surface. Flooding of marshy species has most
28 likely resulted in the apparent decline or extirpation of *Spiranthes* populations in southern Arizona
29 and Utah (Sheviak 1984, McClaran and Sundt 1992). However, Gehlbach (pers. comm. 25 Feb.
30 1990) speculates on a beneficial scheme of periodic flash floods playing a historical role in
31 restoring favorable conditions for the orchid by removing the dense vegetation cover. Control of
32 exotic species like Johnson grass can be accomplished by (i) frequent mowing in areas that are
33 completely dominated by Johnson grass and too dry to support *Spiranthes* and (ii) hand-
34 application of herbicides to weeds in areas that are dominated by native species. Many orchid
35 species cannot compete with fast growing, large herbaceous plants. The population size of
36 *Spiranthes spiralis* growing in areas where land is frequently disturbed (mowed, plowed, etc.)
37 decreases when tall grasses or dense short grasses increase in abundance (Sanford 1974).
38 *Spiranthes ovalis* is a rare plant under undisturbed conditions; however, it readily invades areas
39 that have been altered, particularly abandoned wooded pastures and old fields (Sheviak 1974).
40 *Cypripedium candidum* and *Spiranthes lacera* thrive in sites where annual mowing occurs (Curtis
41 1946; Sheviak 1974). A recovery in the number of *Cypripedium candidum* plants was seen within
42 five years of initiation of mowing practices which reduced the amount of shrubs (Curtis 1946). In
43 mowed sites, flowering of *Spiranthes lacera* is directly dependent (the dependency was not
44 explained) on the clipping regime (Sheviak 1974). Reduction of accumulated litter can be
45 accomplished by prescribed burning, grazing, mowing, or clipping. Disagreement over the most
46 natural management regime for the *Spiranthes* exists, with several individuals suggesting burning
47 (M. Heitlinger, pers. comm. 8 Jan. 1990; P. Warren, pers. comm. 25 Jan. 1990) and others
48 recommending grazing (Gehlbach 1986; P. Sundt, pers. comm. 23 Jan. 1990). Manipulations
49 which alter the soil characteristics should be avoided in the fall when the seeds are most likely
50 beginning to germinate and commence the mycorrhizal relationship; in many orchid species the
51 initial orchid-fungi association is extremely precarious (Wells 1981). More information on the
52 orchid's life-cycle and environmental requirements and experimentation on the effect of different
53 management practices (grazing, fire, mowing, clipping) are needed to identify the most effective
54 management procedures.

Monitoring Requirements: Monitoring *Spiranthes delitescens* at all known sites is needed to assess the current status of the species. There is some background information on population numbers of aboveground plants at two well-studied sites. Both populations appear to be declining; the declines have been most dramatic at one site. There are no estimates of population size for the other two populations which are known only from collection records. Monitoring can also be used to understand the developmental processes and ecological requirements of this species, thereby increasing our ability to accurately forecast and interpret population fluctuations. A permanent marking system should be employed, allowing for continual monitoring of individual plants. The position of each plant should be labelled with respect to the perimeter of the specific plot in which the plant is contained. Labelled stakes, indicating the precise location, should be placed consistently on one side (i.e. due north) of each plant. McClaran and Sundt (1992) use a 1 m X 1 m square placed over permanent corner stakes to mark the plot boundary, and each plant is labelled with both the distance to each stake and the direction (E or W) relative to the line connecting the two stakes. Yearly vegetative and floral measurements should be taken consistently in August, during the period of flower and fruit development. Measurements on each individual plant should include presence or absence of vegetative and floral growth, height of shoot and inflorescence and number of flowers and fruits. The percentage of mature fruits which contain seeds is valuable information, since some *Spiranthes* species develop fruit without producing seeds (Catling 1982). Along with the yearly detailed monitoring, visual observations of the vegetative conditions (presence or absence of leaf rosettes) throughout the year should be noted. The environmental requirements for germination, growth, survivorship and reproduction are unknown for *Spiranthes delitescens*. If research indicates that one or more of the following environmental parameters are important, then this parameter(s) should also be monitored on a monthly or biweekly basis throughout the growing season. Potentially important environmental parameters may include: soil temperature, moisture, pH, light intensity at soil level and 10 cm above the soil level (orchid leaf level), and precipitation. Possibly, complete soil analyses should be performed periodically in order to determine differences in mineral availability and microorganism diversity at the various sites.

Management Programs: Burning experiments are being planned for one protected site. The study site will be divided into thirds and the three treatments will include a control, burns conducted every two years and every seven years. Contact: Mark Heitlinger, Director of Stewardship, The Nature Conservancy, Arizona Field Office, Tucson, Arizona.

Monitoring Programs: Several monitoring programs are currently underway at one protected site.

Contacts: Peter Warren/Dave Gori, The Nature Conservancy, Arizona Field Office, Tucson, Arizona 85705; (602) 622-3861. The monitoring plan for *Spiranthes* includes counts of vegetative and reproductive individuals in eleven experimental plots that were randomly assigned one of three prescribed burn treatments. Dave Gori has received funds from The Nature Conservancy to develop a monitoring plan for *S. delitescens* in 1992.

Mitchel McClaran and Peter Sundt, Department of Range Management, University of Arizona, Tucson, Arizona 85721; (602) 621-1673. Vegetative and floral parameters of the *Spiranthes* have been monitored by various people from 1978 to 1989 (McClaran and Sundt 1992).

Judy Davis, Department of Hydrology, University of Arizona, Tucson, Arizona 85721; (602) 621-1723. Monitoring of several hydrological features at the cienega have been conducted from 1988 to 1990 (J. Davis, pers. comm. 29 Jan. 1990).

Spiranthes spiralis was monitored from 1963 until 1980 at the following location: Knocking Hoe National Nature Reserve, Bedfordshire, England (Wells 1981).

Management Research Programs: The Nature Conservancy is now conducting a long-term study to assess the effect of prescribed burns and burn frequency on the structure and

composition of cienega vegetation and *Spiranthes*. For more information about this study, contact: Dave Gori, The Nature Conservancy, 300 E. University Blvd., #230, Tucson, Arizona 85705; (602) 622-3861.

Cytological and morphological studies have been performed by: Charles Sheviak, Botanist, New York State Museum, Albany, New York.

References:

- Arditti, J. 1982. Orchid biology, reviews, and perspectives. Cornell University Press, Ithaca, NY. 390p.
- Arizona Game and Fish Department. 2000. *Spiranthes delitescens*. Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- Bender, J. 1986. Element stewardship abstract for *Cypripedium candidum*. The Nature Conservancy. Midwest Regional Office, Minneapolis, MN.
- Catling, P. 1982. Breeding systems of northeastern North American *Spiranthes*. Canadian Journal of Botany 60: 3017-3034.
- Catling, P. and V. Catling. 1988. *Spiranthes nebularum* a new species from southern Mexico and Guatemala. Rhodora 90:139-146.
- Curtis, J. 1946. Use of mowing in management of white ladyslipper. Journal of Wildlife Management 10(4): 303-308.
- Dressler, R. 1981. The Orchids: Natural History and Classification. Harvard Univ. Press, Cambridge, MA. 332 pp.
- Fernald, A. 1987. Plant community ecology of two desert marshes in southeastern Arizona. M.A. thesis, University of Colorado, Boulder, CO.
- Gehlbach, F. 1986. Professor, Department of Biology, Baylor University. Correspondence, TNC files, Tucson, AZ.
- Grater, R. 1973. Canelo Hills Cienega, an evaluation. The Nature Conservancy files, Tucson, AZ.
- Kartesz, J.T. 1994. A synonymized checklist of the vascular flora of the United States, Canada, and Greenland. 2nd edition. 2 vols. Timber Press, Portland, OR.
- Luer, C.A. 1975. The native orchids of the United States and Canada excluding Florida. New York Botanical Garden. 361 pp.
- McClaran, M. and P. Sundt. 1992. Population dynamics of the rare orchid, *Spiranthes delitescens*. In press, Southwestern Naturalist.
- Merrigan, S. 1990. Element stewardship abstract - Cienega (in preparation). The Nature Conservancy, Tucson, AZ.
- Rutman, S. 1992. Handbook of Arizona's endangered, threatened, and candidate plants. U.S. Fish and Wildlife Service, Phoenix, AZ.
- Rutman, S., and J. Rorabaugh. 1995. Endangered and threatened wildlife and plants: Proposal to determine endangered status for three wetland species found in southern Arizona and northern Sonora. Federal Register 60(63): 16836-16846.
- Sanford, W. 1974. The ecology of orchids. In C. Withner, editor, The orchids, scientific studies. John Wiley & Sons, New York, NY. 1-101p.

- 1 • Sheviak, C.J. 1984. *Spiranthes diluvialis* (Orchidaceae), a new species from the western
2 United States. *Brittonia* 36:8-14.
- 3 • Sheviak, C.J. 1990. A new *Spiranthes* (Orchidaceae) from the cienegas of southernmost
4 Arizona. *Rhodora* 92(872):213-231.
- 5 • Sheviak, C.T. 1974. An Introduction to the Ecology of the Illinois Orchidaceae, Illinois
6 State Museum Scientific Papers XIV, Springfield, IL, U.S.A. B74SHE01PAUS
- 7 • Stoutamire, W. 1974. Terrestrial orchid seedlings. In C. Withner, editor, *The Orchids,*
8 *scientific studies.* John Wiley and Sons, New York, NY. 101-128 p.
- 9 • Warcup, J. 1975. Factors affecting symbiotic germination of orchid seed. In F. Sanders,
10 B. Mosse, and P. Tinker. editors, *Endomycorrhizas.* Academic Press, London, England.
11 87-105p.
- 12 • Wells, T. 1967. Changes in a population of *Spiranthes spiralis* Chevall. at Knocking Hoe
13 National Nature Reserve, Bedfordshire, 1962-65. *Journal of Ecology* 55:83-89.
- 14 • Wells, T. 1981. Population ecology of terrestrial orchids. In H. Synge, editor, *The*
15 *biological aspects of rare plant conservation.* John Wiley and Sons, New York, NY. 181-
16 195 p.
- 17 • Williams, L. 1951. *The Orchidaceae of Mexico.* Cieza 2:1-132.
- 18 • van der Pijl, L., and C. Dodson. 1966. *Orchid flowers: Their pollination and evolution.*
19 University of Miami Press, Coral Gables, FL. 214 pp.

**BIOLOGICAL SURVEY
ATTACHMENT B**

(b) (7)(E) STATION WILDLIFE SPECIES LISTS

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
BIRDS				
Blackbirds, Orioles				
Emberizidae				
<i>Agelaius phoeniceus</i>	Red-winged blackbird	G5/S5	---	---
<i>Dolichonyx oryzivorus</i>	Bobolink	G5/S1	---	---
<i>Euphagus cyanocephalus</i>	Brewer's blackbird	G5/S5	---	---
<i>Icterus bullockii</i>	Bullock's oriole	G5/SNRB	---	---
<i>Icterus cucullatus</i>	Hooded oriole	G5/S5	---	---
<i>Icterus parisorum</i>	Scott's oriole	G5/S5	---	---
<i>Molothrus aeneus</i>	Bronzed cowbird	G5/S5	---	---
<i>Molothrus ater</i>	Brown-headed cowbird	G5/S5	---	---
<i>Quiscalus mexicanus</i>	Great-tailed grackle	G5/S5	---	---
<i>Sturnella magna</i>	Eastern meadowlark	G5/S5	---	---
<i>Sturnella neglecta</i>	Western meadowlark	G5/S5	---	---
<i>Xanthocephalus xanthocephalus</i>	Yellow-headed blackbird	G5/S5	---	---
Bushtits				
Aegithalidae				
<i>Psaltirparus minimus</i>	Bushtit	G5/S5	---	---
Caracaras, Falcons				
Falconidae				
<i>Caracara cheriway</i>	Crested caracara	G5/S1S2	---	---
<i>Falco columbarius</i>	Merlin	G5/S4N	---	---
<i>Falco mexicanus</i>	Prairie falcon	G5/S4	---	---
<i>Falco peregrinus</i>	Peregrine falcon	G4T4/S4	SC	WSC
<i>Falco sparverius</i>	American kestrel	G5/S5	---	---
Cormorants				
Phalacrocoracidae				
<i>Phalacrocorax auritus</i>	Double-crested cormorant	G5/S5	---	---
<i>Phalacrocorax brasilianus</i>	Neotropic cormorant	G5/S1N	---	---
Cranes				
Gruidae				
<i>Grus canadensis</i>	Sandhill crane	G5/S3N	---	---
Crows and Jays				
Corvidae				
<i>Aphelocoma californica</i>	Western scrub jay	G5/S5	---	---
<i>Corvus corax</i>	Common raven	G5/S5	---	---
<i>Corvus cryptoleucus</i>	Chihuahuan raven	G5/S4	---	---
<i>Cyanocitta stelleri</i>	Steller's jay	G5/S5	---	---
Cuckoos				
Cuculidae				
<i>Coccyzus americanus</i>	Yellow-billed cuckoo	G3T3Q/S3	C	WSC

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
BIRDS (continued)				
<i>Geococcyx californianus</i>	Greater roadrunner	G5/S5	---	---
Doves				
Columbidae				
<i>Columbina inca</i>	Inca dove	G5/S5	---	---
<i>Columbina passerina</i>	Common ground-dove	G5/S4	---	---
<i>Columbina talpacote</i>	Ruddy ground-dove	G5/S1B,S2N	---	---
<i>Zenaida asiatica</i>	White-winged dove	G5/S5	---	---
<i>Zenaida macroura</i>	Mourning dove	G5/S5	---	---
Ducks, Geese, Swans				
Anatidae				
<i>Aix sponsa</i>	Wood duck	G5/S2B,S3N	---	---
<i>Anas acuta</i>	Northern pintail	G5/S2B,S5N	---	---
<i>Anas americana</i>	American wigeon	G5/S1B,S5N	---	---
<i>Anas clypeata</i>	Northern shoveler	G5/S1B,S5N	---	---
<i>Anas crecca</i>	Green-winged teal	G5/S3B,S5N	---	---
<i>Anas cyanoptera</i>	Cinnamon teal	G5/S5	---	---
<i>Anas discors</i>	Blue-winged teal	G5/S2B,S5N	---	---
<i>Anas penelope</i>	Eurasian wigeon	G5/S2N	---	---
<i>Anas platyrhynchos</i>	Mallard	G5/S5	---	---
<i>Anas strepera</i>	Gadwall	G5/S5	---	---
<i>Anser albifrons</i>	Greater white-fronted goose	G5/S2N	---	---
<i>Aythya affinis</i>	Lesser scaup	G5/S5N	---	---
<i>Aythya americana</i>	Redhead	G5/S4	---	---
<i>Aythya collaris</i>	Ring-necked duck	G5/S5	---	---
<i>Aythya valisneria</i>	Canvasback	G5/S1B,S4N	---	---
<i>Branta canadensis</i>	Canada goose	G5/S1B,S4N	---	---
<i>Bucephala albeola</i>	Bufflehead	G5/S5N	---	---
<i>Chen caerulescens</i>	Snow goose	G5/S3N	---	---
<i>Cygnus columbianus</i>	Tundra swan	G5/S1N	---	---
<i>Dendrocygna autumnalis</i>	Black-bellied whistling-duck	G5/S3	---	WSC
<i>Dendrocygna bicolor</i>	Fulvous whistling-duck	G5/?	---	---
<i>Lophodytes cucullatus</i>	Hooded merganser	G5/S2N	---	---
<i>Mergus merganser</i>	Common merganser	G5/S3S4	---	---
<i>Oxyura jamaicensis</i>	Ruddy duck	G5/S5	---	---
Finches				
Fringillidae				
<i>Carduelis lawrencei</i>	Lawrence's goldfinch	G3G4/S1,S3N	---	---
<i>Carduelis pinus</i>	Pine siskin	G5/S5	---	---
<i>Carduelis psaltria</i>	Lesser goldfinch	G5/S5	---	---
<i>Carduelis tristis</i>	American goldfinch	G5/S1B,S5N	---	---
<i>Carpodacus cassinii</i>	Cassin's finch	G5/S4	---	---

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
BIRDS (continued)				
<i>Carpodacus mexicanus</i>	House finch	G5/S5	---	---
<i>Carpodacus purpureus</i>	Purple finch	G5/S1,S2N	---	---
Gnatcatchers				
Muscicapidae				
<i>Poliophtila caerulea</i>	Blue-gray gnatcatcher	G5/S5	---	---
<i>Poliophtila melanura</i>	Black-tailed gnatcatcher	G5/S5	---	---
Goatsuckers				
Caprimulgidae				
<i>Chordeiles acutipennis</i>	Lesser nighthawk	G5/S5	---	---
<i>Phalaenoptilus nuttallii</i>	Common poorwill	G5/S5	---	---
Grebes				
Podicipedidae				
<i>Aechmophorus occidentalis</i>	Western grebe	G5/S3	---	---
<i>Podiceps nigricollis</i>	Eared grebe	G5/S3B,S5N	---	---
<i>Podilymbus podiceps</i>	Pied-billed grebe	G5/S5	---	---
Grosbeaks and Buntings				
Emberizidae				
<i>Cardinalis cardinalis</i>	Northern cardinal	G5/S5	---	---
<i>Cardinalis sinuatus</i>	Pyrrhuloxia	G5/S5	---	---
<i>Guiraca caerulea</i>	Blue grosbeak	G5/S5	---	---
<i>Passerina amoena</i>	Lazuli bunting	G5/S4	---	---
<i>Passerina ciris</i>	Painted bunting	G5/S2,S3M	---	---
<i>Passerina cyanea</i>	Indigo bunting	G5/S3	---	---
<i>Passerina versicolor</i>	Varied bunting	G5/S3	---	---
<i>Pheucticus ludovicianus</i>	Rose-breasted grosbeak	G5/S3N	---	---
<i>Pheucticus melanocephalus</i>	Black-headed grosbeak	G5/S5	---	---
<i>Spiza americana</i>	Dickcissel	G5/S2M	---	---
Gulls, Terns				
Laridae				
<i>Chlidonias niger</i>	Black tern	G4/S3,S4M	---	---
<i>Larus delawarensis</i>	Ring-billed gull	G5/S5N	---	---
<i>Larus philadelphia</i>	Bonaparte's gull	G5/S3,S4M	---	---
<i>Sterna forsteri</i>	Forster's tern	G5/S2N	---	---
Hawks, Kites, Eagles				
Accipitridae				
<i>Accipiter cooperi</i>	Cooper's hawk	G5/S4	---	---
<i>Accipiter striatus</i>	Sharp-shinned hawk	G5/S4	---	---
<i>Aquila chrysaetos</i>	Golden eagle	G5/S4	---	---
<i>Buteo albonotatus</i>	Zone-tailed hawk	G4/S4	---	---
<i>Buteo jamaicensis</i>	Red-tailed hawk	G5/S5	---	---

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
<i>Buteo lagopus</i>	Rough-legged hawk	G5/SNRN	---	---
BIRDS (continued)				
<i>Buteo nitidus</i>	Gray hawk	G5T4Q/S3	SC	WSC
<i>Buteo regalis</i>	Ferruginous hawk	G5/S3	---	---
<i>Buteo swainsoni</i>	Swainson's hawk	G5/S3	---	---
<i>Buteogallus anthracinus</i>	Common black-hawk	G4G5/S3	---	WSC
<i>Circus cyaneus</i>	Northern harrier	G5/S1S2B,S5N	---	---
<i>Elanus leucurus</i>	White-tailed kite	G5/S2B,S2S3N	---	---
<i>Haliaeetus leucocephalus</i>	Bald eagle	G5/S4N	LT,PDL	WSC
<i>Ictinia mississippiensis</i>	Mississippi kite	G5/S3	---	WSC
<i>Pandion haliaetus</i>	Osprey	G5/S2B,S4N	---	---
<i>Parabuteo unicinctus</i>	Harris's hawk	G5/S5	---	---
Herons, Bitterns, Allies				
Ardeidae				
<i>Ardea herodias</i>	Great blue heron	G5/S5	---	---
<i>Botaurus lentiginosus</i>	American bittern	G4/S1S2	---	---
<i>Bubulcus ibis</i>	Cattle egret	G5/S1B, S4N	---	---
<i>Butorides striatus</i>	Green-backed heron	G5/S4	---	---
<i>Casmerodius albus</i>	Great egret	G5/S1B,S4N	---	---
<i>Egretta caerulea</i>	Little blue heron	G5/S1S2N	---	---
<i>Egretta thula</i>	Snowy egret	G5/S1B,S4N	---	---
<i>Ixobrychus exilis</i>	Least bittern	G5/S3	---	---
<i>Nycticorax nycticorax</i>	Black-crowned night-heron	G5/S3	---	---
<i>Nycticorax violaceus</i>	Yellow-crowned night-heron	G5/?	---	---
Hummingbirds				
Trochilidae				
<i>Archilochus alexandri</i>	Black-chinned hummingbird	G5/S5	---	---
<i>Calypte anna</i>	Anna's hummingbird	G5/S5	---	---
<i>Calypte costae</i>	Costa's hummingbird	G5/S5	---	---
<i>Cynanthus latirostris</i>	Broad-billed hummingbird	G4/S3	---	---
<i>Eugenes fulgens</i>	Magnificent hummingbird	G5/S4	---	---
<i>Selasphorus platycercus</i>	Broad-tailed hummingbird	G5/S5	---	---
<i>Selasphorus rufus</i>	Rufous hummingbird	G5/S5M	---	---
<i>Stellula calliope</i>	Calliope hummingbird	G5/S4M	---	---
Ibises				
Threskiornithidae				
<i>Eudocimus albus</i>	White ibis	G5/?	---	---
<i>Plegadis chihi</i>	White-faced ibis	G5/S?B,S2S3N	SC	---
Kingfishers				

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
Alcedinidae				
BIRDS (continued)				
<i>Ceryle alcyon</i>	Belted kingfisher	G5/S2B,S5N	---	---
<i>Chloroceryle americana</i>	Green kingfisher	G5/S2	---	---
Kinglets and Thrushes				
Muscicapidae				
<i>Catharus guttatus</i>	Hermit thrush	G5/S5	---	---
<i>Regulus calendula</i>	Ruby-crowned kinglet	G5/S5	---	---
<i>Sialia mexicana</i>	Western bluebird	G5/S5	---	---
<i>Turdus migratorius</i>	American robin	G5/S5	---	---
Larks				
Alaudidae				
<i>Eremophila alpestris</i>	Horned lark	G5/S5	---	---
Loons				
Gaviidae				
<i>Gavia immer</i>	Common loon	G5/S2N	---	---
Mockingbirds and Thrashers				
Mimidae				
<i>Dumetella carolinensis</i>	Gray catbird	G5/S1	---	---
<i>Mimus polyglottos</i>	Northern mockingbird	G5/S5	---	---
<i>Oreoscoptes montanus</i>	Sage thrasher	G5/S5	---	---
<i>Toxostoma bendirei</i>	Bendire's thrasher	G4G5/S4	---	---
<i>Toxostoma curvirostre</i>	Curve-billed thrasher	G5/S5	---	---
<i>Toxostoma dorsale</i>	Crissal thrasher	G5/S5	---	---
<i>Toxostoma rufum</i>	Brown thrasher	G5/S1N	---	---
Nuthatches				
Sittidae				
<i>Sitta carolinensis</i>	White-breasted nuthatch	G5/S5	---	---
Owls				
Strigidae				
<i>Asio otus</i>	Long-eared owl	G5/S2B,S3S4N	---	---
<i>Athene cunicularia</i>	Burrowing owl	G4T4/S3	SC	---
<i>Bubo virginianus</i>	Great horned owl	G5/S5	---	---
<i>Micrathene whitneyi</i>	Elf owl	G5/S5	---	---
<i>Otus kennicottii</i>	Western screech owl	G4/S4	---	---
Tytonidae				
<i>Tyto alba</i>	Common barn owl	G5TNR/?	---	---
Pelicans				
Pelicanidae				
<i>Pelecanus occidentalis</i>	Brown pelican	G4/S1N	---	---
Pipits				
Motacillidae				
<i>Anthus rubescens</i>	American pipit	G5/S2B,S5N	---	---

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
BIRDS (continued)				
Plovers				
Charadriidae				
<i>Charadrius vociferus</i>	Killdeer	G5/S5	---	---
Quail, New World				
Phasianidae				
<i>Callipepla gambelii</i>	Gambel's quail	G5/S5	---	---
<i>Callipepla squamata</i>	Scaled quail	G5/S5	---	---
<i>Cyrtonyx montezumae</i>	Montezuma quail	G4G5/S4	---	---
Rails, Gallinules, Coots				
Rallidae				
<i>Fulica americana</i>	American coot	G5/S5	---	---
<i>Gallinula chloropus</i>	Common moorhen	G5/S5	---	---
<i>Porzana carolina</i>	Sora	G5/S4	---	---
<i>Rallus limicola</i>	Virginia rail	G5/S4	---	---
Sandpipers, Phalaropes				
Scolopacidae				
<i>Actitis macularius</i>	Spotted sandpiper	G5/S3S4	---	---
<i>Calidris bairdii</i>	Baird's sandpiper	G5/S4M	---	---
<i>Calidris mauri</i>	Western sandpiper	G5/S1N	---	---
<i>Calidris minutilla</i>	Least sandpiper	G5/S5N	---	---
<i>Calidris pusilla</i>	Semipalmated sandpiper	G5/S2M	---	---
<i>Catoptrophorus semipalmatus</i>	Willet	G5/S4M	---	---
<i>Gallinago delicata</i>	Wilson's snipe	G5/S1B,S4N	---	---
<i>Limnodromus scolopaceus</i>	Long-billed dowitcher	G5/S3S4N	---	---
<i>Numenius americana</i>	Long-billed curlew	G5/S1B,S3S4N	---	---
<i>Phalaropus tricolor</i>	Wilson's phalarope	G5/S1B,S5N	---	---
<i>Tringa flavipes</i>	Lesser yellowlegs	G5/S4M	---	---
<i>Tringa melanoleuca</i>	Greater yellowlegs	G5/S3N	---	---
<i>Tringa solitaria</i>	Solitary sandpiper	G5/S3M	---	---
Shrikes				
Laniidae				
<i>Lanius ludovicianus</i>	Loggerhead shrike	G4/S4	---	---
Silky Flycatchers				
Ptilonotidae				
<i>Phainopepla nitens</i>	Phainopepla	G5/S5	---	---
Sparrows, New World				
Emberizidae				
<i>Aimophila botterii</i>	Botteri's sparrow	G4/S4	---	---
<i>Aimophila cassinii</i>	Cassin's sparrow	G5/S4	---	---
<i>Aimophila ruficeps</i>	Rufous-crowned sparrow	G5/S4	---	---
<i>Ammodramus savannarum</i>	Grasshopper sparrow	G5/S3	---	---

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
BIRDS (continued)				
<i>Amphispiza belli</i>	Sage sparrow	G5/S4	---	---
<i>Amphispiza bilineata</i>	Black-throated sparrow	G5/S5	---	---
<i>Calamospiza melanocorys</i>	Lark bunting	G5/S1B,S5N	---	---
<i>Calcarius ornatus</i>	Chestnut-collared longspur	G5/S3N	---	---
<i>Chondestes grammacus</i>	Lark sparrow	G5/S5	---	---
<i>Junco hyemalis</i>	Dark-eyed junco	G5/S5		
<i>Melospiza georgiana</i>	Swamp sparrow	G5/S2S3N	---	---
<i>Melospiza lincolni</i>	Lincoln's sparrow	G5/S3B,S5N	---	---
<i>Melospiza melodia</i>	Song sparrow	G5/S5	---	---
<i>Passerculus sandwichensis</i>	Savannah sparrow	G5/S5	---	---
<i>Pipilo chlorurus</i>	Green-tailed towhee	G5/S3B,S4N	---	---
<i>Pipilo fuscus</i>	Canyon towhee	G5/S5	---	---
<i>Pipilo maculatus</i>	Spotted towhee	G5/S5	---	---
<i>Pooecetes gramineus</i>	Vesper sparrow	G5/S5	---	---
<i>Spizella atrogularis</i>	Black-chinned sparrow	G5/S5	---	---
<i>Spizella breweri</i>	Brewer's sparrow	G5/S5	---	---
<i>Spizella passerina</i>	Chipping sparrow	G5/S5	---	---
<i>Zonotrichia albicollis</i>	White-throated sparrow	G5/S2S3N	---	---
<i>Zonotrichia atricapilla</i>	Golden-crowned sparrow	G5/S1S2N	---	---
<i>Zonotrichia leucophrys</i>	White-crowned sparrow	G5/S1B,S5N	---	---
Sparrows, Old World				
Passeridae				
<i>Passer domesticus</i>	House sparrow	G5/SNA	---	---
Starlings				
Sturnidae				
<i>Sturnus vulgaris</i>	European starling	G5/SNA	---	---
Stilts, Avocets				
Recurvirostridae				
<i>Himantopus mexicanus</i>	Black-necked stilt	G5/S2	---	---
<i>Recurvirostra americana</i>	American avocet	G5/S2	---	---
Storks				
Ciconiidae				
<i>Mycteria americana</i>	Wood stork	G4/S1N	---	---
Swallows				
Hirundinidae				
<i>Hirundo pyrrhonota</i>	Cliff swallow	G5/S5	---	---
<i>Hirundo rustica</i>	Barn swallow	G5/S5	---	---
<i>Progne subis</i>	Purple martin	G5/S4	---	---

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
<i>Riparia riparia</i>	Bank swallow	G5/S4M	---	---
BIRDS (continued)				
<i>Stelgidopteryx serripennis</i>	Northern rough-winged swallow	G5/S5	---	---
<i>Tachycineta bicolor</i>	Tree swallow	G5/S3	---	---
<i>Tachycineta thalassina</i>	Violet-green swallow	G5/S5	---	---
Swifts				
Apodidae				
<i>Aeronautes saxatilis</i>	White-throated swift	G5/S5	---	---
<i>Chaetura vauxi</i>	Vaux's swift	G5/S4M	---	---
Tanagers				
Emberizidae				
<i>Piranga ludoviciana</i>	Western tanager	G5/S5	---	---
<i>Piranga rubra</i>	Summer tanager	G5/S4	---	---
Titmice				
Paridae				
<i>Parus wollweberi</i>	Bridled titmouse	G5/S4	---	---
Turkeys				
Galliformes				
<i>Meleagris gallopavo</i>	Wild turkey	G5/S5	---	---
Tyrant Flycatchers				
Tyrannidae				
<i>Campostoma imberbe</i>	Northern beardless-tyrannulet	G5/S4	---	---
<i>Contopus cooperi</i>	Olive-sided flycatcher	G4/S4	---	---
<i>Contopus sordidulus</i>	Western wood-peewee	G5/S5	---	---
<i>Empidonax hammondi</i>	Hammond's flycatcher	G5/S1B,S2S3N	---	---
<i>Empidonax oberholseri</i>	Dusky flycatcher	G5/S4	---	---
<i>Empidonax occidentalis</i>	Cordilleran flycatcher	G5/S2S3B	---	---
<i>Empidonax traillii</i>	Willow flycatcher	G5/S1	---	---
<i>Empidonax wrightii</i>	Gray flycatcher	G5/S5	---	---
<i>Myiarchus cinerascens</i>	Ash-throated flycatcher	G5/S5	---	---
<i>Myiarchus tuberculifer</i>	Dusky-capped flycatcher	G5/S4	---	---
<i>Myiarchus tyrannulus</i>	Brown-crested flycatcher	G5/S4	---	---
<i>Myiodynastes luteiventris</i>	Sulphur-bellied flycatcher	G5/S3	---	---
<i>Pachyramphus aglaiae</i>	Rose-throated becard	G4G5/S1	---	---
<i>Pyrocephalus rubinus</i>	Vermilion flycatcher	G5/S5	---	---
<i>Sayornis nigricans</i>	Black phoebe	G5/S5	---	---
<i>Sayornis phoebe</i>	Eastern phoebe	G5/S1N	---	---
<i>Sayornis saya</i>	Say's phoebe	G5/S5	---	---

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
<i>Tyrannus crassirostris</i>	Thick-billed kingbird	G5/S2	---	WSC
<i>Tyrannus melancholicus</i>	Tropical kingbird	G5/S3	---	WSC
BIRDS (continued)				
<i>Tyrannus verticalis</i>	Western kingbird	G5/S5	---	---
<i>Tyrannus vociferans</i>	Cassin's kingbird	G5/S5	---	---
Verdins				
Remizidae				
<i>Auriparus flaviceps</i>	Verdin	G5/S5	---	---
Vireos				
Vireonidae				
<i>Vireo bellii</i>	Bell's vireo	G5/S4	---	---
<i>Vireo gilvus</i>	Warbling vireo	G5/S5	---	---
<i>Vireo huttoni</i>	Hutton's vireo	G5/S5	---	---
<i>Vireo plumbeus</i>	Plumbeous vireo	G5/S5	---	---
<i>Vireo solitarius</i>	Blue-headed vireo	G5/?	---	---
Vultures, New World				
Cathartidae				
<i>Cathartes aura</i>	Turkey vulture	G5/S5	---	---
<i>Coragyps atratus</i>	Black vulture	G5/S1S2	---	---
Waxwings				
Bombycillidae				
<i>Bombycilla cedrorum</i>	Cedar waxwing	G5/S3S4N	---	---
Woodpeckers				
Picidae				
<i>Colaptes auratus</i>	Northern flicker	G5/S5	---	---
<i>Colaptes chrysoides</i>	Gilded flicker	G5/S5	---	---
<i>Melanerpes lewis</i>	Lewis's woodpecker	G4/S4	---	---
<i>Melanerpes uropygialis</i>	Gila woodpecker	G5/S5	---	---
<i>Picoides scalaris</i>	Ladder-backed woodpecker	G5/S5	---	---
<i>Sphyrapicus nuchalis</i>	Red-naped sapsucker	G5/S4	---	---
Wood Warblers				
Emberizidae				
<i>Dendroica caerulescens</i>	Black-throated blue warbler	G5/S1M	---	---
<i>Dendroica coronata</i>	Yellow-rumped warbler	G5/S5	---	---
<i>Dendroica nigrescens</i>	Black-throated gray warbler	G5/S5	---	---
<i>Dendroica occidentalis</i>	Hermit warbler	G4G5/S4M	---	---
<i>Dendroica petechia</i>	Yellow warbler	G5/S4	---	---
<i>Dendroica townsendii</i>	Townsend's warbler	G5/S4M,S1S2N	---	---
<i>Geothlypis trichas</i>	Common yellowthroat	G5/S4	---	---
<i>Icteria virens</i>	Yellow-breasted chat	G5/S4	---	---
<i>Myioborus pictus</i>	Painted redstart	G5/S4	---	---

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
<i>Oporornis formosus</i>	Kentucky warbler	G5/S1M	---	---
<i>Oporornis tolmiei</i>	MacGillivray's warbler	G5/S4	---	---
<i>Seiurus noveboracensis</i>	Northern waterthrush	G5/S2S3M	---	---
<i>Setophaga ruticilla</i>	American redstart	G5/S1	---	---
<i>Vermivora celata</i>	Orange-crowned warbler	G5/S3B,S5N	---	---
<i>Vermivora luciae</i>	Lucy's warbler	G5/S5	---	---
<i>Vermivora ruficapilla</i>	Nashville warbler	G5/S4S5M	---	---
<i>Vermivora virginiae</i>	Virginia's warbler	G5/S5	---	---
<i>Wilsonia pusilla</i>	Wilson's warbler	G5/S5M	---	---
Wrens				
Troglodytidae				
<i>Campylorhynchus brunneicapillus</i>	Cactus wren	G5/S5	---	---
<i>Cistothorus palustris</i>	Marsh wren	G5/S2B,S3S4N	---	---
<i>Salpinctes obsoletus</i>	Rock wren	G5/S5	---	---
<i>Thryomanes bewickii</i>	Bewick's wren	G5/S5	---	---
<i>Troglodytes aedon</i>	House wren	G5/S5	---	---
MAMMALS				
Badgers and Skunks				
Mephitidae				
<i>Mephitis mephitis</i>	Striped skunk	G5/S5	---	---
<i>Spilogale gracilis</i>	Western spotted skunk	G5/S5	---	---
Bats, Free-tailed				
Molossidae				
<i>Nyctinomops macrotis</i>	Big free-tailed bat	G5/S2S3	---	---
<i>Tadarida brasiliensis</i>	Brazilian free-tailed bat	G5/S3S4	---	---
Bats, Leaf-nose				
Phyllostomidae				
<i>Leptonycteris curasoae</i>	Lesser long-nosed bat	G4/S2	LE	WSC
Bats, Plain-nose				
Vespertilionidae				
<i>Antrozous pallidus</i>	Pallid bat	G5/S4S5	---	---
<i>Eptesicus fuscus</i>	Big brown bat	G5/S4S5	---	---
<i>Lasiurus blossevillei</i>	Western red bat	G5/S2	---	WSC
<i>Lasiurus cinereus</i>	Hoary bat	G5/S4	---	---
<i>Lasiurus xanthinus</i>	Western yellow bat	G5/S1	---	WSC
<i>Myotis auriculus</i>	Southwestern myotis	G5/S3	---	---
<i>Myotis californicus</i>	California myotis	G5/S4S5	---	---
<i>Myotis thysanodes</i>	Fringed myotis	G4G5/S3S4	SC	---
<i>Myotis velifer</i>	Cave myotis	G5/S4	SC	---
<i>Pipistrellus hesperus</i>	Western pipistrelle	G5/S5	---	---
<i>Plecotus townsendii</i>	Townsend's big-eared bat	G4/S3S4	---	---

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
Badgers				
Mustelidae				
<i>Taxidea taxus</i>	American badger	G5/S5	---	---
MAMMALS (continued)				
Bears				
Ursidae				
<i>Ursus americanus</i>	Black bear	G5/S5	---	---
Cats				
Felidae				
<i>Lynx rufus</i>	Bobcat	G5/S5	---	---
<i>Puma concolor</i>	Mountain lion	G5/S4	---	---
Coyotes and Foxes				
Canidae				
<i>Canis latrans</i>	Coyote	G5/S5	---	---
<i>Urocyon cinereoargenteus</i>	Gray fox	G5/S5	---	---
Deer				
Cervidae				
<i>Odocoileus hemionus</i>	Mule deer	G5/S5	---	---
<i>Odocoileus virginianus</i>	White-tailed deer	G5/S5	---	---
Javelina				
Tayassuidae				
<i>Pecari tajacu</i>	Collared peccary	G5/S5	---	---
Pocket Gophers				
Geomyidae				
<i>Thomomys bottae</i>	Botta's pocket gopher	G5/S5	---	---
<i>Thomomys umbrinus</i>	Southern pocket gopher	G5/S4	---	---
Pocket Mice and Kangaroo Rats				
Heteromyidae				
<i>Chaetodipus baileyi</i>	Bailey pocket mouse	G5/S5	---	---
<i>Chaetodipus hispidus</i>	Hispid pocket mouse	G5/S5	---	---
<i>Chaetodipus penicillatus</i>	Desert pocket mouse	G5/S5	---	---
<i>Dipodomys merriami</i>	Merriam's kangaroo rat	G5/S5	---	---
<i>Dipodomys ordii</i>	Ord's kangaroo rat	G5/S5	---	---
<i>Dipodomys spectabilis</i>	Banner-tailed kangaroo rat	G5/S5	---	---
<i>Perognathus flavus</i>	Silky pocket mouse	G5/S5	---	---
Porcupines				
Erethizontidae				
<i>Erethizon dorsatum</i>	Porcupine	G5/S4S5	---	---
Rabbits and Hares				
Leporidae				
<i>Lepus californicus</i>	Black-tailed jackrabbit	G5/S5	---	---

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
<i>Sylvilagus audubonii</i>	Desert cottontail	G5/S5	---	---
Raccoon and Ringtail				
Procyonidae				
<i>Nasua narica</i>	White-nosed coati	G5/S4	---	---
MAMMALS (continued)				
<i>Procyon lotor</i>	Common raccoon	G5/S4	---	---
Rats and Mice				
Muridae				
<i>Mus musculus</i>	House mouse	G5/SNA	---	---
<i>Neotoma albigula</i>	White-throated woodrat	G5/S5	---	---
<i>Neotoma mexicana</i>	Mexican woodrat	G5/S5	---	---
<i>Onychomys leucogaster</i>	Northern grasshopper mouse	G5/S5	---	---
<i>Onychomys torridus</i>	Southern grasshopper mouse	G5/S5	---	---
<i>Peromyscus leucopus</i>	White-footed mouse	G5/S5	---	---
<i>Peromyscus maniculatus</i>	Deer mouse	G5/S5	---	---
<i>Reithrodontomys fulvescens</i>	Fulvous harvest mouse	G5/S4	---	---
<i>Reithrodontomys megalotis</i>	Western harvest mouse	G5/S5	---	---
<i>Sigmodon arizonae</i>	Arizona cotton rat	G5/S4	---	---
<i>Sigmodon hispidus</i>	Hispid cotton rat	G5/S5	---	---
<i>Sigmodon ochrognathus</i>	Yellow-nosed cotton rat	G4G5/S3S4	SC	---
Shrews				
Soricidae				
<i>Notiosorex crawfordi</i>	Desert shrew	G5/S4S5	---	---
Squirrels				
Sciuridae				
<i>Ammospermophilus harrisi</i>	Yuma antelope squirrel	G5/S5	---	---
<i>Spermophilus spilosma</i>	Spotted ground squirrel	G5/S4	---	---
<i>Spermophilus variegatus</i>	Rock squirrel	G5/S5	---	---
REPTILES				
Box Turtles				
Emydidae				
<i>Terrapene ornata</i>	Desert box turtle	G5/S3S4	---	---
Mud Turtles				
Kinosternidae				
<i>Kinosternon sonoriense</i>	Sonoran mud turtle	G4/S4	---	---
Alligator Lizards				
Anguidae				

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
<i>Elgaria kingii</i>	Madrean alligator lizard	G5/S5	---	---
Beaded Lizards				
Helodermatidae				
<i>Heloderma suspectum</i>	Gila monster	G4/S4	---	---
REPTILES (continued)				
Collared and Leopard Lizards				
Crotaphytidae				
<i>Crotaphytus collaris</i>	Collared lizard	G5/S5	---	---
<i>Gambelia wislizenii</i>	Long-nosed leopard lizard	G5/S5	---	---
Iguanid Lizards				
Phrynosomatidae				
<i>Holbrookia maculata</i>	Lesser earless lizard	G5/S5	---	---
<i>Phrynosoma cornutum</i>	Texas horned lizard	G4G5/S3S4	----	---
<i>Phrynosoma solare</i>	Regal horned lizard	G5/S5	---	---
<i>Sceloporus clarkii</i>	Clark's spiny lizard	G5/S5	---	---
<i>Sceloporus undulatus</i>	Eastern fence lizard	G5/SNR	---	---
<i>Urosaurus ornatus</i>	Ornate tree lizard	G5/S5	---	---
Whiptail Lizards				
Teiidae				
<i>Cnemidophorus uniparens</i>	Desert grassland whiptail	G5/S5	---	---
Colubrid Snakes				
Colubridae				
<i>Arizona elegans</i>	Glossy snake	G5/S5	---	---
<i>Diadophis punctatus</i>	Ring-necked snake	G5/S4	---	---
<i>Hypsiglena torquata</i>	Nightsnake	G5/S5	---	---
<i>Lampropeltis getula</i>	Common kingsnake	G5/S5	---	---
<i>Masticophis bilineatus</i>	Sonoran whipsnake	G5/S5	---	---
<i>Masticophis flagellum</i>	Coachwhip	G5/S5	---	---
<i>Pituophis catenifer</i>	Gopher snake	G5/S5	---	---
<i>Rhinocheilus lecontei</i>	Long-nosed snake	G5/S5	---	---
<i>Salvadora hexalepis</i>	Western patch-nosed snake	G5/S5	---	---
<i>Sonora semiannulata</i>	Ground snake	G5/S5	---	---
<i>Tantilla hobartsmithii</i>	Southwestern black-headed snake	G5/S5	---	---
<i>Thamnophis eques</i>	Mexican garter snake	G5/S2S3	---	---
<i>Thamnophis marcianus</i>	Checkered garter snake	G5/S5	---	---
Coral Snakes				
Elapidae				
<i>Micruroides euryxanthus</i>	Sonoran coral snake	G5/S5	---	---
Rattlesnakes				

Group / Family / Scientific Name	Common Name	Global and State Rank	State Status	Federal Status
Viperidae				
<i>Crotalus atrox</i>	Western diamondback rattlesnake	G5/S5	---	---
<i>Crotalus scutulatus</i>	Mojave rattlesnake	G5/S5	---	---
AMPHIBIANS				
Spadefoot Toads				
Pelobatidae				
<i>Scaphiopus couchii</i>	Couch's spadefoot	G5/S5	---	---
<i>Spea multiplicata</i>	Mexican spadefoot	G5/S5	---	---
Toads				
Bufo				
<i>Bufo alvarius</i>	Colorado River toad	G5/S5	---	---
<i>Bufo cognatus</i>	Great Plains toad	G5/S5	---	---
<i>Bufo debilis</i>	Green toad	G5/S3	---	---
<i>Bufo punctatus</i>	Red-spotted toad	G5/S5	---	---
<i>Bufo woodhousii</i>	Woodhouse's toad	G5/S5	---	---
True Frogs				
Ranidae				
<i>Rana catesbeiana</i>	Bullfrog	G5/SNA	---	---
<i>Rana chiricahuensis</i>	Chiricahua leopard frog	G3/S3	LT	WSC
<i>Rana yavapaiensis</i>	Lowland leopard frog	G4/S4	SC	WSC
FISH				
Catfish				
Ictaluridae				
<i>Ictalurus pricei</i>	Yaqui catfish	G2/S1	LT	WSC
Minnows				
Cyprinidae				
<i>Campostoma ornatum</i>	Mexican stoneroller	G3/S1	SC	WSC
<i>Cyprinella formosa</i>	Beautiful shiner	G2/S1	LT	WSC
<i>Gila purpurea</i>	Yaqui chub	G1/S1	LE	WSC
<i>Gila robusta</i>	Roundtail chub	G3/S2	---	---
<i>Rhinichthys chrysogaster</i>	Longfin dace	G4/S3S4	SC	---
Suckers				
Catostomidae				
<i>Catostomus berrardini</i>	Yaqui sucker	G4/SX	---	---
Topminnows				
Poeciliidae				
<i>Poeciliopsis occidentalis sonoriensis</i>	Yaqui topminnow	G3T3/S1	LE	WSC

Source: USFWS 2003

(b) (7)(E) Watchable Wildlife List. Global and State Rank from NatureServe 2008.

Federal and State Status from AGFD 2007.